



HAL
open science

Assessing the effect of spatial resolution of regional climate downscaling on the productivity and distribution in three widespread tree species over France

Nicolas Martin-StPaul, Marc Stéfanon, Christophe François, Kamel Soudani, Philippe Drobinski, Julien Ruffault, Serge Rambal, Florent Mouillot, Paul W. Leadley

► To cite this version:

Nicolas Martin-StPaul, Marc Stéfanon, Christophe François, Kamel Soudani, Philippe Drobinski, et al.. Assessing the effect of spatial resolution of regional climate downscaling on the productivity and distribution in three widespread tree species over France. European Geosciences Union General Assembly 2013, Apr 2013, Viena, Austria. hal-02807367

HAL Id: hal-02807367

<https://hal.inrae.fr/hal-02807367v1>

Submitted on 6 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Assessing the effect of spatial resolution of regional climate downscaling on the productivity and distribution of three widespread tree species over France

Nicolas K. Martin-StPaul, Marc Stephanon, Christophe Francois , Kamel Soudani , Eric Dufrêne , Phillipe Drobinski, Alissar Cheaib, Julien Ruffault, Serge Rambal, Florent Mouillot, and Paul Leadley
ESE, CNRS, University Paris-Sud, Orsay, France (nico09.martin@gmail.com)

The recent increases in temperature and water deficit as a result of climate changes have already impaired forest functioning and might trigger tree dieback worldwide in the near future. The assessment of future forest conditions relies on mechanistic models that predict changes in trees and forest functioning as a function of meteorological drivers. Currently, global and regional models (GCM and RCM) are the main providers of climate forcing in impact studies. One large uncertainty when forecasting the forest functioning is associated with the coarse spatial resolution of climate scenarii. In this study we assessed how the spatial resolution in climate forcing provided by the RCM WRF impacted the simulated productivity and distribution of three species (*Fagus sylvatica*, *Quercus ilex*) over France. We ran the forest model CASTANEA over France (that simulates fluxes of carbon and water and forest growth) using the output of WRF at different spatial scales (50 km, 20km, 8km and 1km) as forcing climate entries. The productivity simulated by CASTANEA was used as a surrogate of beech persistence for the reference period of WRF (1988-2008). Because climate variables simulated by WRF exhibited large bias compared to surface observations, WRF was first corrected using a reference dataset (SAFRAN database) upscaled at the WRF resolution (50km and 20 km). Additionally, on 2 specific limited areas (the Languedoc Roussillon and the Bourgogne region) we used a statistical downscaling of the WRF forcing entries in order to increase the spatial resolution up to 1km. Our results showed that simulations at finer resolution had relatively little impact on the mean and variance of beech productivity over France compared to coarser resolutions. However, at the finest resolutions, we observed strong local gradients with important variations in the mean and the variance of forest productivity (up to 60%). These results are particularly noticeable in regions characterized by complex orography such as the Alps or the Mediterranean. Our results highlight the need to consider the importance of spatial scale of climate input when studying climate impacts on tree distributions in areas with steep climatic gradients.