



Root and rhizospheric phosphatase activity of grassland Poaceae grown under contrasted water and phosphorus supplies

Florian Fort, Eric Lecloux, Ciprian Stroia, Pablo Cruz, Claire Jouany

► To cite this version:

Florian Fort, Eric Lecloux, Ciprian Stroia, Pablo Cruz, Claire Jouany. Root and rhizospheric phosphatase activity of grassland Poaceae grown under contrasted water and phosphorus supplies. International society of Root Research (ISRR) “Roots to the future”, Jun 2012, Dundee, United Kingdom. 1 p., 2012. hal-02808767

HAL Id: hal-02808767

<https://hal.inrae.fr/hal-02808767>

Submitted on 6 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Root and rhizospheric phosphatase activity of grassland *Poaceae* grown under contrasted water and phosphorus supplies

Fort F.^{1*}, Lecloux E.¹, Stroia C.², Cruz P.¹, Jouany C.¹

¹ INRA UMR 1248 AGIR ; BP 52627, 31326 Castanet-Tolosan, France.

² USAMVBT, Department of Biology and Plant Protection, Calea Aradului 119, 300645 Timisoara, Romania

* Corresponding author: florian.fort@toulouse.inra.fr

Comparing to the above ground, a few below ground traits are available for a functional characterization of plants. This gap is quite obvious when considering plant functioning for mineral resource acquisition, specially for phosphorus. We hypothesize that a better understanding of relationships between plants functional traits and phosphatase activity (PA) may allowed to build more consistent plant functional types regarding to the phosphorus nutrition.

Objectives


In this work we propose an evaluation of the potential of root PA and rhizospheric PA as a functional trait to indentify plants P limitation and distinguish between acquisitive and conservation species.

Materials & Methods

Experimental design:

Functional Group	Species	Substrate	Water	Phosphorus	Applied treatments
Conservative	<i>Bromus erectus</i> Huds.	Sand 65% Silt 19% Clay 16%	W1 = non limited water supply / W0 = 1/3 W1	P1 = non limited phosphorus / P0 = limited phosphorus	W1 & P0
	<i>Lolium arundinaceum</i> (Schreb.) Darbysh.				W1 & P1
Acquisitive	<i>Dactylis glomerata</i> L.				W0 & P0
	<i>Holcus lanatus</i> L.				W0 & P1

→ 16 plants (10 repeats by treatments) + 12 bulk = 172 pots



* Alkaline phosphomonoesterase activity was measured as described by Johnson *et al.* 1999.

In order to compare the phosphorus stress levels among species and between growth strategies we calculated an index of stress intensity (derivate from Callaway *et al.* 2002) :

$$\text{Phosphorus Stress Intensity Index} = \frac{\text{Mean (Biomass in P1)} - \text{Biomass in P0}}{\text{Max}(\text{Mean}(\text{Biomass in P1}), \text{Biomass in P0})}$$

Results & Discussion




Figure 2: Root phosphatase activity (mean ± se) for crossed phosphorus treatment, water treatment and plant strategy. Different letters highlight significant difference (ANOVA, p<0.05).

We observe a positive and significant effect of the lack of P supply and a positive effect of the water supply on root PA (ANOVA p<0.001). Acquisitive species have higher root PA than conservative ones in case of P starvation (ANOVA, p<0.001). Moreover this difference is higher in the case of non limited water supply, when the plants demand of phosphorus for growth is higher.

Conclusion

Phosphatase activity measured on root and rhizospheric soil is associated to the level of plant phosphorus stress and is also very sensitive to the water supply level. These traits allowed the species separation between capture & conservation strategy, but only in phosphorus stressful conditions.





Figure 1: Root phosphatase activity (µg.g⁻¹.h⁻¹) against the Index of phosphorus stress with water treatment used as a covariate.

When water does not limit the plants growth the root PA increases significantly with the phosphorus stress level (ANCOVA, Fig. 1). However under limited water supply there is not relationship between the phosphorus stress index and the root PA..




Figure 3: Rhizospheric and Bulk soil phosphatase activity (mean ± se, µg.g⁻¹.h⁻¹) for crossed water and phosphorus treatments. Different letters highlight significant difference (ANOVA, p<0.05).

For all treatments PA measured on the rhizospheric soil is significantly higher than the one which is measured for the bulk soil (Fig. 3), revealing a significant positive impact of the root proximity on the soil PA. The rhizospheric soil PA is significantly improved by phosphorus starvation and also by water limited supply in case of phosphorus starvation (Fig. 3).

Références :

Johnson D, Leake JR, Lee JA (1999) The effects of quantity and duration of simulated pollutant nitrogen deposition on root surface phosphatase activities in calcareous and acid grasslands: a bioassay approach. New Phytol 141:433–442.
Callaway RM, Pennings SC, Richards CL (2003) Phenotypic plasticity and interactions among plants. Ecology 84(5) 1115-1128