



**HAL**  
open science

## **Brown trout life history and habitat use in the Kerguelen Islands as revealed by otolith microchemistry**

Jean-Christophe Aymes, Edward Beall, Marc Jarry, Philippe Gaudin

### ► **To cite this version:**

Jean-Christophe Aymes, Edward Beall, Marc Jarry, Philippe Gaudin. Brown trout life history and habitat use in the Kerguelen Islands as revealed by otolith microchemistry. 8. Journées Scientifiques, Comité National Français des Recherches Arctiques et Antarctiques (CNFRAA). FRA., May 2012, Plouzané, France. 1 p. <hal-02808933>

**HAL Id: hal-02808933**

**<https://hal.inrae.fr/hal-02808933v1>**

Submitted on 6 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

## **Brown trout life history and habitat use in the Kerguelen Islands as revealed by otolith microchemistry**

J-C Aymes<sup>1,2</sup>, E. Beall<sup>1,2</sup>, M. Jarry<sup>2,1</sup>, P. Gaudin<sup>1,2</sup>

<sup>1</sup> INRA, UMR ECOBIOP, INRA/UPPA, Pôle d'Hydrobiologie de l'INRA, Quartier Ibarron 64310 Saint Pée sur Nivelle, France

<sup>2</sup> Univ Pau & Pays Adour, UMR ECOBIOP, INRA/UPPA, UFR Côte Basque, Allée du Parc Montaury, 64600 Anglet, France

### **Résumé**

Characterization of the life history traits of introduced brown trout (*Salmo trutta* L.) in the Kerguelen Islands is fundamental to understand the success of the islands invasion by this fish species. The capacity to use marine habitats (Anadromy) by brown trout is one of the most critical traits in the invasion process and its study is essential to understand how marine habitats contribute to dispersal in trout populations. Otoliths are calcified structure (aragonite) located in the inner ear of fishes; they record seasonal growth as scales but are also inert and able to trap trace element present in the fish environment (Campana and Thorrold, 2001). Strontium, particularly the Sr:Ca ratio, is a common indicator used to trace brackish/salt water migrations of salmonids (Limburg et al., 2001; Milton et al., 2008). We studied Sr:Ca ratios on otoliths of brown trout captured in specific habitats (lake, upstream and downstream zones of rivers, marine coastal waters) to characterize life history traits linked to freshwater or salt water migration (resident/migrant status, age at migration, homing, maternal origin). Results showed that the method is relevant because water and otolith present correlated chemical signatures across each selected Kerguelen habitat. A strong diversity of life histories appeared, ranging from pure residency in freshwater to permanent use of saltwater. Intermediate phenotypes were observed in downstream populations taking benefit of the highly productive estuarine habitats whereas specific migratory phenotypes could contribute to colonize new pristine aquatic systems.

### **Références**

Campana, S.E., 1999. Chemistry and composition of fish otoliths: pathways, mechanisms and applications. Marine Ecology Progress Series 188, 263-297.

Limburg, K.E., Landergrén, P., Westin, L., Elfman, M., Kristiansson, P., 2001. Flexible modes of anadromy in Baltic sea trout: making the most of marginal spawning streams. Journal of Fish Biology 59 (3), 682-695.

Milton, D.A., Chenery, Simon R, 2001. Sources and uptake of trace metals in otoliths of juvenile barramundi (*Lates calcarifer*). Journal of Experimental Marine Biology and Ecology 264 (1), 47e65.