



HAL
open science

Insights into Colletotrichum hemibiotrophy from genome and transcriptome sequencing

Richard O'Connell, Stéphane Hacquard, Pamela Gan, Emiel Ver Loren van Themaat, Jochen Kleemann, Stefan Amyotte, Michael R Thon, Li-Jun Ma, Ken Shirasu, Lisa Vaillancourt

► **To cite this version:**

Richard O'Connell, Stéphane Hacquard, Pamela Gan, Emiel Ver Loren van Themaat, Jochen Kleemann, et al.. Insights into Colletotrichum hemibiotrophy from genome and transcriptome sequencing. XI International Fungal Biology Conference, Sep 2013, Karlsruhe, Germany. <hal-02808987>

HAL Id: hal-02808987

<https://hal.inrae.fr/hal-02808987v1>

Submitted on 6 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

Insights into *Colletotrichum* hemibiotrophy from genome and transcriptome sequencing

Richard O'Connell^{1,2}, Stéphane Hacquard¹, Pamela Gan³, Emiel Ver Loren van Themaat¹, Jochen Kleemann¹, Stefan Amyotte⁴, Michael Thon⁵, Li-Jun Ma⁶, Ken Shirasu³, Lisa Vaillancourt⁴

1 Max Planck Institute for Plant Breeding Research, Cologne, Germany

2 INRA-BIOGER, Thiverval-Grignon, France

3 Plant Science Center, RIKEN, Yokohama, Japan

4 Department of Plant Pathology, University of Kentucky, Lexington, Kentucky, USA

5 CIALE, Universidad de Salamanca, Villamayor, Spain

6 University of Massachusetts Amherst, Amherst, Massachusetts, USA

Species of *Colletotrichum* cause devastating diseases on numerous crop plants worldwide. These pathogens use a hemibiotrophic infection strategy that involves the formation of specialized cell-types for initial penetration (appressoria), growth inside living plant cells (biotrophic hyphae) and tissue destruction (necrotrophic hyphae). The genomes of 4 species were sequenced to date: *C. higginsianum* (*Ch*), *C. graminicola* (*Cg*), *C. orbiculare* (*Co*) and *C. fructicola*, which respectively infect *Arabidopsis*, maize, cucurbits and strawberry. All 4 genomes encode extraordinarily large inventories of plant cell wall-degrading enzymes, with the three dicot pathogens having twice as many pectinases as the monocot pathogen *C. graminicola*, reflecting adaptation to host cell wall composition. Genes encoding secondary metabolism (SM) enzymes, especially those involved in the synthesis of polyketides, terpenes and alkaloids, are also expanded compared to other sequenced fungal pathogens, suggesting *Colletotrichum* species are capable of producing great chemical diversity. Similar to biotrophic pathogens, all 4 species encode large, lineage-specific repertoires of putative effector proteins, including an expanded family of chitin-binding LysM proteins that may function in evasion of PAMP-triggered immunity. The transcriptional dynamics underlying hemibiotrophy were examined using RNA-Seq (*Ch*, *Cg*) and microarrays (*Co*). Although appressoria formed *in vitro* resemble those *in planta*, comparison of their transcriptomes showed >1,500 *Ch* genes were induced only upon host contact, suggesting that plant signals sensed by appressoria reprogram fungal gene expression in preparation for host invasion. SM enzymes and secreted effectors were among the most highly up-regulated genes in appressoria, suggesting a role for these infection structures in host manipulation. During the biotrophic phase, transcripts encoding effectors and SM enzymes were also abundant, but there was no specific induction of nutrient uptake transporters, suggesting biotrophic hyphae function primarily as organs for delivering small molecule and protein effectors rather than nutrient acquisition. At the switch to necrotrophy, genes encoding a vast array of wall-degrading enzymes, proteases and membrane transporters were up-regulated, enabling the pathogen to mobilize nutrients from dead and dying cells to fuel its rapid growth and sporulation.