



**HAL**  
open science

## Field measurements of photosynthetic $^{13}\text{C}$ discrimination of *Fagus Sylvatica* branches using laser spectrometry

Lydia Gentsch, Patrick Sturm, Albin Hammerle, Rolf Siegwolf, Lisa Wingate, Jérôme Ogée, Matthias Barthel, Peter Plüss, Thomas Baur, Nina Buchmann, et al.

### ► To cite this version:

Lydia Gentsch, Patrick Sturm, Albin Hammerle, Rolf Siegwolf, Lisa Wingate, et al.. Field measurements of photosynthetic  $^{13}\text{C}$  discrimination of *Fagus Sylvatica* branches using laser spectrometry. JESIUM 2012, Joint European Stable Isotope Users Group Meeting (JESIUM). Saisissez le nom du laboratoire, du service ou du département., Ville service., Sep 2012, Leipzig, Germany. 1 p. hal-02809003

**HAL Id: hal-02809003**

**<https://hal.inrae.fr/hal-02809003v1>**

Submitted on 6 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Field measurements of photosynthetic $^{13}\text{C}$ discrimination of *Fagus Sylvatica* branches using laser spectrometry

LYDIA GENTSCH<sup>1</sup>, PATRICK STURM<sup>2</sup>, ALBIN HAMMERLE<sup>3</sup>, ROLF SIEGWOLF<sup>4</sup>, LISA WINGATE<sup>5</sup>, JÉRÔME OGÉE<sup>5</sup>, MATTHIAS BARTHEL<sup>1</sup>, PETER PLÜSS<sup>1</sup>, THOMAS BAUR<sup>1</sup>, NINA BUCHMANN<sup>1</sup> & ALEXANDER KNOHL<sup>6</sup>

<sup>1</sup>Institute of Agricultural Sciences, ETH Zürich, Zürich, Switzerland, lydia.gentsch@usys.ethz.ch

<sup>2</sup>Laboratory for Air Pollution / Environmental Technology, Empa, Dübendorf, Switzerland

<sup>3</sup>Institute of Ecology, University of Innsbruck, Innsbruck, Austria

<sup>4</sup>Laboratory for Atmospheric Chemistry / Stable Isotopes & Ecosystem Fluxes, PSI, Villigen, Switzerland

<sup>5</sup>INRA, UR1263 Ephyse, Villenave d'Ornon, France

<sup>6</sup>Chair of Bioclimatology, Georg-August University of Göttingen, Göttingen, Germany

Isotopic discrimination against  $^{13}\text{C}_2$  during photosynthesis ( $^{13}\Delta$ ) causes the overall  $^{13}\text{C}$  depletion of the terrestrial biosphere compared to the atmosphere.  $^{13}\Delta$  varies in response to environmental variables that influence photosynthetic gas exchange. For  $\text{C}_3$  plants,  $^{13}\Delta$  mainly reflects the balance between the  $\text{CO}_2$  supply to and the  $\text{CO}_2$  demand by the carboxylation sites, but it is, to a smaller extent, also influenced by carbon isotope fractionations occurring during mitochondrial and photo-respiration. Understanding and predicting  $^{13}\Delta$  variability has gained relevance for  $\text{CO}_2$  flux partitioning on the global and the ecosystem scale, for tree ring analysis or for insights into plant/soil dynamics. Estimates of  $^{13}\Delta$  are commonly obtained by combining carbon isotope measurements of plant-derived organic matter or air profiles with  $^{13}\Delta$ -models. In contrast, direct, gas exchange-based measurements of  $^{13}\Delta$  under field conditions are sparse, due to the technical deployment involved. Hence, we have little experimental verification of diurnal and seasonal variabilities of  $^{13}\Delta$  and their relation to environmental drivers. We further lack a thorough field-based evaluation of the Farquhar et al. (1982)  $^{13}\Delta$ -model. Here, we present continuous, hourly field measurements of  $^{13}\Delta$  of *Fagus sylvatica* L. branches, conducted with three custom-made open branch bags and a laser spectrometer for  $\text{CO}_2$  isotopologue measurements (QCLAS-ISO, Aerodyne Research Inc.). Data from two field campaigns (in total 38 and 60 days) in a mature, temperate mixed-deciduous forest in Switzerland during August / September 2009 and 2010 are shown. We observed a high diurnal variability of  $^{13}\Delta$ , with average diurnal amplitudes of  $\approx 10\text{‰}$  and maximum diurnal amplitudes of  $\approx 20\text{‰}$ . Highest  $^{13}\Delta$  were generally observed during dawn and dusk, and lowest  $^{13}\Delta$  during midday. Morning and afternoon  $^{13}\Delta$  commonly displayed intermediate values. Day-to-day variations were summarized with flux-weighted daily means of  $^{13}\Delta$ , which ranged from  $15\text{‰}$  to  $23\text{‰}$  in 2009 and from  $18\text{‰}$  to  $29\text{‰}$  in 2010. If trees were not water-limited, branch gas exchange, and hence  $^{13}\Delta$ , appeared to be mainly driven by changes in incident PAR. Woody tissue  $\text{CO}_2$  efflux was measured with two independent approaches to assess its potential bias on branch bag measurements. We estimated that its contribution to branch net  $\text{CO}_2$  assimilation was  $< 0.1 \mu\text{mol m}^{-2} \text{s}^{-1}$  per total leaf area during the time of our field campaigns.

Using data from the 2010 field campaign only, we explored the applicability of the comprehensive ( $^{13}\Delta_{\text{comp}}$ ), the simplified ( $^{13}\Delta_{\text{simple}}$ ) and the revised ( $^{13}\Delta_{\text{revised}}$ ) versions of the Farquhar et al. (1982) - model for predicting observed diurnal and day-to-day  $^{13}\Delta$  variabilities.  $^{13}\Delta_{\text{comp}}$  predicted the mean diurnal variability of  $^{13}\Delta$  much better than  $^{13}\Delta_{\text{simple}}$  (RMSE<sub>simple</sub>  $\approx 3.4$ ; RMSE<sub>comp</sub>  $\approx 2.5$ ). Furthermore,  $^{13}\Delta_{\text{comp}}$  was more suitable than  $^{13}\Delta_{\text{simple}}$  for predicting flux-weighted daily means of observed  $^{13}\Delta$ . For model calibrations, a Bayesian inference approach was used. This approach allowed us to reliably quantify uncertainties in the model parameter estimation and to reveal the amount of model-relevant information present in our field dataset.

### Reference

Farquhar GD, O'Leary MH, Berry JA (1982) On the relationship between carbon isotope discrimination and the inter-cellular carbon dioxide concentration in leaves. *Aust. J. Plant Physiol.* 9:121-137