



**HAL**  
open science

## **A dairy vector exclusively fermented by dairy propionibacteria: a new model to study probiotic potentialities in vivo**

Fabien Cousin, Séverine Louesdon, Laurence Le Normand, Gaëlle Boudry, Sandrine Jouan-Lanhouet, Denis Mater, Marie-Thérèse Dimanche-Boitrel, Gwénaél Jan

### ► **To cite this version:**

Fabien Cousin, Séverine Louesdon, Laurence Le Normand, Gaëlle Boudry, Sandrine Jouan-Lanhouet, et al.. A dairy vector exclusively fermented by dairy propionibacteria: a new model to study probiotic potentialities in vivo. International Scientific Conference on Probiotics and Prebiotics (IPC), Jun 2011, Kosice, Slovakia. 1 p., 2011. hal-02809096

**HAL Id: hal-02809096**

**<https://hal.inrae.fr/hal-02809096>**

Submitted on 6 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# **A dairy vector exclusively fermented by dairy propionibacteria: a new model to study probiotic potentialities *in vivo***

Fabien COUSIN<sup>1,2,3</sup>, Séverine LOUESDON<sup>1,2</sup>, Laurence LE NORMAND<sup>4</sup>, Gaëlle BOUDRY<sup>4</sup>, Sandrine JOUAN-LANHOUE<sup>5</sup>, Denis MATER<sup>3</sup>, Marie-Thérèse DIMANCHE-BOITREL<sup>5</sup> and Gwénaél JAN<sup>1,2</sup>

<sup>1</sup> INRA, UMR1253 Science et Technologie du Lait et de l'Œuf, F-35042 Rennes, France

<sup>2</sup> AGROCAMPUS OUEST, UMR1253 Science et Technologie du Lait et de l'Œuf, F-35042 Rennes, France

<sup>3</sup> CNIEL/Syndifrais, 42 rue de Châteaudun, F-75314 Paris 09, France

<sup>4</sup> INRA, UMR1079 SENA, F-35590 Saint Gilles, France

<sup>5</sup> EA SeRAIC, IRSET, Faculté de pharmacie, Université de Rennes 1, IFR 140 GFAS, F-35043, Rennes, France

## **Introduction**

Dairy propionibacteria, commonly used as cheese ripening starters, are also consumed as probiotics in a freeze-dried form. Their probiotic potential includes modulation of the microbiota and epithelial proliferation/apoptosis equilibrium in the gut. They trigger apoptosis of colorectal cancer cells. Short chain fatty acids, propionate and acetate, their major metabolic end-products, are responsible for this programmed knock out of cancer cells, via the mitochondrial (intrinsic) death pathway. Such effects require high populations of live and metabolically active propionibacteria to reach the colon. In this context, appropriate vectors must favour probiotic efficiency. Fermented dairy products protect probiotic bacteria against digestive stresses. Emmental or fermented milks containing dairy propionibacteria may thus deliver live and metabolically active propionibacteria to the colon. Those matrices contain, however, a mixture of other microorganisms, limiting the identification of specific beneficial effects of propionibacteria. We aimed at 1) developing milk exclusively fermented by a dairy propionibacterium; 2) studying propionibacteria survival and tolerance to digestive stress in such a model dairy vector and 3) evaluating the pro-apoptotic potential of the fermented milk on cultured HT-29 human colorectal cancer cells.

## **Methods**

Since dairy propionibacteria do not generally grow in milk, we determined their nutritional requirements with respect to carbon and nitrogen by supplementing milk ultrafiltrate with different concentrations of food grade lactate and casein hydrolysate.

Digestive stress tolerance was assessed *in vitro* towards acid challenge (pH=2) and bile salts challenge (1 g/L). Long term survival during 4°C storage was evaluated by enumerations and a live/dead staining method. Besides, the most tolerant *Propionibacterium freudenreichii* strain was used in an animal trial. Pigs were fed with the model fermented milk (1E+10 cfu/day), then *P. freudenreichii* survival was monitored in faeces and colon contents at slaughtering.

Pro-apoptotic potential of previously neutralised and sterilised fermented milk supernatants was evaluated on HT-29 cells, using usual indicators of apoptosis: cell viability assay, cell cycle analysis, activation of caspases, chromatin condensation and annexin V/7-AAD staining.

## **Results**

Milk supplemented with 50 mM lactate and 5 g/L casein hydrolysate allowed growth of all species and subspecies of the dairy propionibacteria studied, which reached populations of at least  $1E+9$  cfu/mL.

In the model fermented milk, dairy propionibacteria remained viable and stress-tolerant *in vitro* during at least 15 days at 4°C. The population of the *P. freudenreichii* strain used in the *in vivo* trial reached  $1E+7$  cfu/g in the faeces and  $1E+6$  cfu/g in the colon content of pigs.

The model fermented milk was shown to induce apoptosis of HT-29 cells. Typical features of the apoptosis process were observed, including chromatin condensation, formation of apoptotic bodies, DNA laddering, ROS production, caspases activation, cell cycle arrest and emergence of a subG1 population.

## **Discussion**

This work leads to the development of a new food grade vector containing dairy propionibacteria as the sole microorganism, allowing planning of preclinical and clinical trials in human. Such new fermented milks might be of interest in the future as a functional food to prevent colorectal cancer or to potentialize therapeutic treatments.