



HAL
open science

Alternatives to synthetic fungicides using small molecules of natural origin

Christian Chervin

► **To cite this version:**

Christian Chervin. Alternatives to synthetic fungicides using small molecules of natural origin. Plant Defence: Biological Control, 12, Springer Science + Business Media B.V., 412 p., 2012, Progress in Biological Control, 978-94-007-1932-3 978-94-007-1933-0. 10.1007/978-94-007-1933-0_3. hal-02809109

HAL Id: hal-02809109

<https://hal.inrae.fr/hal-02809109>

Submitted on 10 Jan 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Open Archive Toulouse Archive Ouverte (OATAO)

OATAO is an open access repository that collects the work of Toulouse researchers and makes it freely available over the web where possible.

This is an author-deposited version published in: <http://oatao.univ-toulouse.fr/>
Eprints ID: 5334

To link to this article: DOI:10.1007/978-94-007-1933-0_3
http://dx.doi.org/10.1007/978-94-007-1933-0_3

To cite this version:

Chervin, Christian *Alternatives to synthetic fungicides using small molecules of natural origin*. (2012) In: Plant Defence: Biological Control. (Progress in Biological Control). Springer, New York, USA, pp. 55-66. ISBN 978-94-007-1932-3

Any correspondence concerning this service should be sent to the repository administrator: staff-oatao@inp-toulouse.fr

Alternatives to synthetic fungicides using small molecules of natural origin

Christian CHERVIN

Prof. Food and Wine Science, Université de Toulouse,
UMR Génomique et Biotechnologie des Fruits, INRA-INP/ENSAT,
BP 32607, 31326 Catagnet-Tolosan, France.
email : chervin@ensat.fr

Abstract

Several small molecules of natural origin have been reported to act as alternatives to synthetic fungicides. These are reviewed in this chapter, and some ideas of new development are given.

The list, which is not exhaustive, is the following: acetaldehyde, acetic acid, aldehydes (other than acetaldehyde), aminobutyric acids, ascorbic acid, ethanol, ethylene, jasmonic acid and methyl jasmonate, saicylic acid and methyl salicylate, salts (e.g. sodium bicarbonate, calcium chloride, copper sulphate), sorbic acid and sulphur. There is also a paragraph about the potential of combinations (e.g. additive or synergistic effects).

Introduction

There are many grape diseases, among which downy and powdery mildew and gray mold, causing important pre- and post-harvest losses. Most commercial grapevines are susceptible to such fungi. To counteract this disease development a wide panel of synthetic fungicides is in use nowadays. However, solutions to limit pre-harvest treatments with synthetic fungicides are of particular interest as chemical residues are limiting access to many markets, and there is a diminishing number of antifungal compounds that are still registered (Nigro et al., 2006). Moreover, some pathogen strains may develop resistance to some

pesticide, thus alternative strategies are required. One of the most sustainable alternative strategies would be to develop grape cultivars that are naturally resistant to such fungi, and there are available genotypes for this trait (Dry et al., 2010). But it is a long term development and most food industries using grapes are relying on specific cultivars, around which all marketing efforts have been made for decades. So finding alternative treatments with existing cultivars is still of interest for the grape industry.

The purpose of this review is to list various compounds of natural origin that have been tested with or without success in the past. They were ranged by alphabetical order.

Acetaldehyde

As this compound has such a low boiling point (20°C), it is very volatile at ambient temperature, so most trials were performed for postharvest applications using it in the vapour phase (Pesis, 2005). As most aldehydes, it has a strong bactericidal and fungicidal potential, as aldehydes are very oxidative compounds. One must attract attention to users that they are very toxic compounds to manipulators and quite oxidative to many parts of the equipment.




Figure 1 : Acetaldehyde (MW: 44g; BP: 20.2°C, 68°F)

Utama et al. (2002) showed that aldehydes were more effective than alcohols in blocking fungus growth *in vitro*. They stated that acetaldehyde was quite effective against various fungi: *Rhizopus stolonifer*, *Penicillium digitatum*, *Colletotrichum musae*, *Erwinia carotovora*, and *Pseudomonas aeruginosa*, showing germicidal effects at concentrations below 1 mmole/per dish, as the assays were run in Petri dishes; but they did not report trials *in vivo*. These

germicidal concentrations corresponded approximately to 60 μ moles acetaldehyde/ litre of air, in the vapour phase, after one hour of application, and they went down to 20 μ moles/litre of air in 5 days at 25°C. When applied on ‘Sultanina’ and ‘Perlette’ grape berries grapes with low sugar content and high acidity, the acetaldehyde was found to increase total soluble solids, to decrease acidity, and to enhance sensory preference (Pesis and Frenkel, 1989), but these authors did not report effects on fungus. These observations were reported later by Avissar and Pesis (1991) who showed that acetaldehyde was controlling the decay of table grapes in a postharvest trial.

Acetaldehyde is probably present in most plant and fruit, but at very low concentration around a few $\text{nmoles.g}_{\text{FW}}^{-1}$ (Chervin et al., 1999), so whether it is effective or not at this natural concentration is not known. However there was an interesting report by Miyake and Shibamoto (1995) showing that acetaldehyde can be produced in aerobic and relatively mild conditions by oxidation of L-ascorbic acid. Thus during the oxidative stress following a fungus infection, there may be some acetaldehyde produced, and it may be part of the natural defence; this has been shown in the case of resistance of potato plants to *Phytophthora infestans* (Tadege et al., 1998).

Acetic acid

The use of acetic acid fumigation for postharvest control of fungi has been reviewed by Tripathi and Dubey (2004). These authors referenced the fact that acetic acid is a natural metabolite occurring in several fruit, and that fumigation with acetic acid onto grapes has been proven an effective treatment to control gray mold.




Figure 2 : Acetic acid (MW: 60g; BP:118.1°C, 245°F)

A recent paper by Camili et al. (2010) showed that acetic acid might stimulate natural defences in Italia grapes, as the best control of Botrytis was obtained when the fruit was treated with acetic acid vapours 48 hours prior inoculation with Botrytis spores. To my knowledge, there no or very scarce data about a potential control of grape disease by spraying acetic acid in vineyards, but the trials may be worth it.

Interestingly, sprays of indole-acetic acid on tomato seedlings were shown to reduce the symptoms caused by a phytopathogen, *Pythium ultimum*. (Gravel et al., 2007). The indole-acetic acid, one of the auxins, is obviously quite different to acetic acid, however no application has been tested onto grapevines to check if such an effect would be observed with grape specific pathogens.

In a more comprehensive approach, Martin and Maris (2004) tested the antifungal and antibacterial efficacy of seventeen organic and mineral acids against several strains of bacteria and fungi, known as food contaminants. They found interesting inhibitory effects by formic, mandelic and lactic acids.

Aldehydes (other than acetaldehyde)

The remarks regarding their toxicity, outlined in the acetaldehyde paragraph, are valid for the following compounds too.

The cinnamaldehyde is part of the cinnamon aroma. It can be used as a food additive, and has anti-microbial properties at quite low concentration around 10 mM (Smid et al., 1996). There no report of exogenous treatment of cinnamaldehyde on grapes, but a recent report shows an additional potential of such treatments with this aldehyde or other compounds listed in this chapter: Viazis et al. (2011) show that cinnamaldehyde limit the viability of an enterohemorrhagic Escherichia coli strain on leafy vegetables, when sprayed at 0.5% (v/v).

The hexanal and hexenal are two natural compounds, oxidation products of lipids. They have been shown to harbour anti-fungal properties against

Botrytis sp., *Alternaria sp.*, and *Penicillium sp.* among others (Tripathi and Dubey, 2004 and refs herein). This has been confirmed recently by Song et al. (2007), as hexanal vapours gave an excellent control of *Monilinia fructicola* on peaches and a very good control of *Botrytis cinerea* on raspberries at doses around 900 $\mu\text{l.l}^{-1}$.



Figure 3 : Hexanal (MW: 100g; BP: 120°C, 248°F)

In planta experiments have been published recently showing that Arabidopsis over-producing C6 aldehydes (e.g. hexanal and hexenal) were more resistant to *Botrytis* infection, mainly through a direct effect of the aldehydes on the fungus growth rather than through an elicitor role that the aldehydes might have had (Kishimoto et al., 2008). This study also leads to think that the spraying of such aldehydes on grapevines may reduce the development of various fungi, however the cost and the hazard of manipulating such molecules have to be considered.

Utama et al. (2002) tested *in vitro* the bactericidal and fungicidal properties of several aldehydes listed above and also benzaldehyde, which was shown to be slightly more efficient than acetaldehyde.

Aminobutyric acids

These compounds are involved in plant induced resistance to fungi, and most recent studies report the effects of β -aminobutyrate (Dubreuil-Morizi et al., 2010; Walz and Simon, 2009; Slaughter et al., 2008). Its chemical structure is close to the γ -aminobutyrate which is a well known neurotransmitter. For in-field applications to various crops, the toxicity to humans has to be considered.




Figure 4 : β -aminobutyric acid (MW: 103g, solid at ambient temperature)

The β -aminobutyrate was shown to induce resistance to downy mildew in sensitive and resistant grape cultivars, and this was related to increased stilbene synthesis and accumulation (Slaugther et al., 2008). These enhanced protection effect against various fungi have been observed in other plants, like cucumber (Walz and Simon, 2009), and these effects were associated with callose accumulation and rapid cell death around the infected area. Recently, a report stated that this enhanced resistance to mildew due to β -aminobutyrate application could be associated with reactive oxygen species produced by a NADPH oxidase (Dubreuil-Morizi et al., 2010).

Ascorbic acid

It is a natural compound which accumulates in many fruit, but in grapes it does not reach high levels as in citrus, as it is a precursor of tartaric and oxalic acids (DeBolt et al. 2004).




Figure 5: Ascorbic acid (MW: 176g, solid at ambient temperature)

It accumulates at low levels in berries, some μmoles per gram of fresh weight (Cruz-Ruz et al., 2010). Whether it shows some antifungal activity at this level is unknown. Authors have observed antimicrobial effects of ascorbic acid at higher concentration such as 2.5% (Van der Wolf et al., 2008), however no test has been recorded on grapevines. However the role of ascorbic acid on phytopathogens is probably complex, indeed Barth et al. (2004) have shown that Arabidopsis mutants deficient in ascorbic acid are more resistant to bacteria and fungi, may be through an increase in salicylic acid accumulation, thus leading to activation of plant natural defences.

Ethanol

This compound is also naturally present in plant and fruit tissues, and can be found under normal aerobic conditions when the inside of the cells become too acidic or under hypoxic conditions (Sweetman et al., 2009, and refs herein).




Figure 6 : Ethanol (MW: 46g; BP: 78°C, 172°F)

Our work regarding applications of ethanol has been initiated by reading an article by Beaulieu and Saltveit (1997). They observed that exogenous ethanol was stimulating ethylene production and tomato fruit ripening, so we tested it on grapes in order to modulate ripening and anthocyanin accumulation (El Kereamy et al., 2002) and wine colour (Chervin et al., 2001) using hand-held sprayers directed towards the bunches. Later we found that ethanol sprays with commercial sprayers increases mostly the berry diameter (Chervin et al., 2005a).

And then we tested ethanol for its efficacy to limit fungus growth. The first application was post-harvest, we will present pre-harvest applications later. The idea came from a paper by Lichter et al. (2002) showing that dipping grapes at harvest in ethanol solutions was decreasing the *Botrytis cinerea* growth. The ethanol dip has two drawbacks which are the need to promptly dry the grapes after treatment to prevent berry cracking (Karabulut et al., 2004), and the possible cross-contamination with fungus spores from a previously infected grape crate, when working with low ethanol concentrations. We adapted these ethanol treatments to commercial practices using ethanol in the vapour phase (Chervin et al., 2005b). Indeed the industry is already using fumigation with SO₂ in adapted chambers, or in crates with SO₂ pads releasing the SO₂ in contact with air humidity, so if ethanol was going to be efficient and accepted by the industry, a simple change of the active ingredient was possible. The application of ethanol vapours was optimised over two seasons for ‘Chasselas’ table grapes and at a dose rate of 2 ml . kg⁻¹ of grapes, the ethanol vapour was as effective as sulphur dioxide pads to prevent rot development, caused by *Botrytis cinerea*, and stem browning. Further tests with consumer panels showed no significant difference in sensory perception between controls and treated grapes. The application of evenly distributed ethanol vapours is critical, as higher concentrations of ethanol may enhance stem browning. Materials releasing ethanol are already on the market, such as the “ethanol powder” (Suzuki et al., 2004). Postharvest applications of ethanol may also present potentials to reduce berry shatter (Chervin et al., 2005c) but these need further development.

Then we tested pre-harvest applications of exogenous ethanol in the vineyard, to prevent fungus development ahead; the results that are detailed below have been reported recently (Chervin et al., 2009). The idea came from the reading of an article by Karabulut et al. (2003). These authors found that spraying 1 litre per 5 vines of a solution at 50% ethanol, 24 hours prior harvest,

was effective in reducing the rots over the postharvest period. We then adapted this to commercial practices, reducing the amount of solution to 150 litres per hectare (using a mist blower) and no treatment in the last two weeks prior harvest. The treatments were performed every two weeks from veraison, we tested a late harvest, scheduled two months and a half after veraison, was chosen for these trials, so the conditions were optimised for high *Botrytis* development. The fungus development was assessed at harvest and after 4 to 6 weeks cold storage. We always found a higher impact of the treatments after cold storage when *Botrytis* development had occurred at higher rates.


In preliminary trials, we found that even a very low concentration of 2% ethanol was reducing the *Botrytis* growth. It is not likely that ethanol would have had a direct effect on fungus growth at this low percentage, as Lichter et al. (2003) showed that at least 30% EtOH is necessary to prevent *Botrytis cinerea* spore germination. Thus the 2% EtOH dose is more likely to induce plant defence.

The optimal dose of ethanol to reduce *Botrytis* growth by pre-harvest spraying was found to be around 16%. However to match the industry demand, we had to combine it with calcium chloride in order to further reduce the gray mold growth. This was done after reading an article by Nigro et al. (2006) in which the authors reported the efficacy of various salts to reduce the *Botrytis* development. This will be detailed in a paragraph below in this chapter. Thus we reported that preharvest applications of a 16% ethanol solution, containing 1% CaCl₂, reduced gray mold development. At harvest the losses due to rotten clusters dropped from 15% in controls to 5% in grapes treated with ethanol & CaCl₂. Over 6 weeks of cold storage, the losses due to gray mold were reduced by 50% in bunches treated with ethanol & CaCl₂, compared to untreated controls. These treatments did not induce significant changes in fruit quality assessed by sensory analysis of healthy berries.

The ethanol has a much higher boiling point than acetaldehyde and is not oxidative, which renders it safer to use; it is already used by industry as a wetting agent or for its solvent properties. As it is flammable, precautions are necessary. Ethanol is rather cheap to produce, and its worldwide production is increasing, mainly due to its use as ethanol fuel, so its cost will decrease.

Ethylene

It is a gas at ambient temperature and it plays important roles as phytohormone in most plants (Lin et al., 2009). These authors review all aspects of ethylene metabolisms in plant biology, including production and perception.



ture)


Regarding the ethylene role in grape defence, there has been a series of recent works by J.M. Merillon's team showing that 2-chloroethylphosphonic acid, an ethylene precursor also called ethephon, led to a decrease of fungus growth when sprayed onto grapevines, through elicitation of natural defences (Belhadj et al., 2008a). More details will be developed in J.M. Merillon's chapter in this book.

Additionally a recent series of microarray analyses, using mRNAs extracted from berry tissues after exogenous application of ethylene on grape clusters, has been partly published by Chervin et al. (2008): the expression of some genes involved in plant defence was shown to be modulated by such a phytohormone.

Jasmonic acid and methyl jasmonate

Jasmonic acid is present naturally in grape berries, particularly in seeds (Kondo and Fukuda, 2001) up to 50 pmoles/seed, and methyl jasmonate is also present to a lesser extent (about 10 times less).

Data are lacking to check the variability over various cultivars.




Methyl jasmonate has been shown to promote stilbene accumulation (Krisa et al., 1999) and grape natural defenses. There will be more details developed in J.M. Merillon's chapter, in this book.

Salicylic acid and methyl salicylate

The salicylic acid is well-known to be part of the SAR: Systemic Acquired Resistance (Durrant and Dong, 2004).

Park et al. (2007) showed that both conversions of salicylic acid to methyl salicylate and methyl salicylate to salicylic acid were essential for SAR in tobacco, and they concluded that methyl salicylate is a SAR signal in this plant.



However there is still some controversy about the direct involvement of salicylic acid and methyl salicylate in SAR (Durrant and Dong, 2004; Attarana et al., 2009).

Nevertheless, there are a few papers reporting application of salicylic acid on grapes (e.g. Wen et al., 2005), showing that this compound can induce metabolisms known to be involved in plant defence mechanisms, but no report with grapes showing induced resistance. On strawberries, Babalar et al. (2007) showed that pre-harvest sprays of salicylic acid had potential to limit fungal decay. The concentration that gave a rather good control was 2 mM, and several sprays were necessary: at the vegetative growth stage, then the fruit development stage and postharvest. This gives good ideas to set-up an experiment with grapes. Salicylic acid may also interest companies and growers as it seems to be available at low cost (Dr Liliana Martinez, Mendoza, pers. comm.).

In addition to controlling the fungal diseases, salicylic acid and derivatives might have other potentials for grape growers. Indeed methyl salicylate was also used to “recruit” beneficial insects in vineyards (James and Price, 2004). These authors showed that use of controlled-release methyl salicylate in a crop could increase recruitment and residency of populations of certain beneficial insects. This strategy may have the potential to enhance the efficacy and reliability of conservation biological control in crop pest management.


Lastly, salicylic acid was shown to delay ripening in particular conditions (Kraeva et al., 1998). Thus the fruit quality is obviously a parameter to take into account in any experiment dealing with applications of such a compound, and all treatments described in this chapter.

Salts (e.g. sodium bicarbonate, calcium chloride, copper sulfate)

Salts are cheap, accepted by consumers, with minor environmental impact at the effective concentrations, which are non-toxic, and they are already used by the food industry. One of the most recent and most comprehensive study about their use as antifungal agents in vineyards has been published by Nigro et al., (2006). These authors tested 19 different salts, from sodium phosphate dibasic, the most efficient against Botrytis, to potassium carbonate the less efficient in a small trial using artificially infested berries. Then they ran larger field trials in field in which the most efficient salts against bunch rots were calcium chloride, sodium bicarbonate and sodium carbonate at concentrations around 1% (w/v). The spray timing was variable, some treatments performed 90 and 30 days prior harvest in the last trial, but also 20 and 5 days prior harvest in earlier trials. The salts were as efficient as the classical chemical treatments in some field trials.

Another wellknown salt used in viticulture is the copper sulfate pentahydrate, which has a blue colour, and its natural form is called chalcantite. It is one of most common antifungal when it is mixed with lime, particularly against downy mildew, and it is named Bordeaux mixture. It is one of the rare antifungal treatments, with sulphur, allowed in organic vineyards. However its intensive use is known to 'pollute' soils and to render some cultures difficult after several years of vine growing, as other crops are not so tolerant to high Cu concentrations in soils.

Sorbic acid



e)

It is a known food preservative, it can be associated with sodium or potassium, among others. Karabulut et al. (2005) showed that potassium sorbate

when applied on 'Thompson Seedless' grapes after harvest, at concentrations around 1%, reduced the incidence of gray mold over storage. No data was found regarding pre-harvest sprays of sorbate salts to control various fungal or bacterial diseases on grapes, but there is some potential.

Sulphur

Sulphur remains a very efficient and simple alternative to synthetic fungicides against powdery mildew. Recent reports by Crisp et al. (2006) confirmed this fact. They compared the efficacy of several compounds, such as milk, whey, canola oils and potassium bicarbonate to sulphur, and this latter was most often the best blocker of powdery mildew. Whey showed some potential, however in field trials the acceptable yield (i.e. bunches with less than 5% powdery mildew infections) was lower than when treated with sulphur. The whey compounds are mainly lactose and lactoglobulin. Why these compounds or the whey pH might have an inhibitory effect on the fungus is not discussed in the article.

Combinations

These are always interesting as they can generate additive or synergistic effects. One example has been detailed above when combining ethanol with calcium chloride during field sprays gave a better control of gray mold at harvest, and after storage (Chervin et al., 2009). Another paper is reporting such effects with ethanol and potassium sorbate (Karabulut et al., 2005). Belhadj et al. (2008b) have tested with success the combination of methyl jasmonate and sucrose to induce accumulation of polyphenolics in grape cell cultures.

Many combinations have been tested on different fruit, giving ideas for grape future treatments. For example Spadaro et al. (2004) have tested with

some success combinations of hot water, backing soda and ethanol against *Penicillium expansum* and *Botrytis cinerea* over apple storage, but the potential may be extended to grapes and field trials. Wang et al. (2010) showed that the combined treatment with methyl jasmonate and ethanol resulted in a greater control of green mold due to *Penicillium citrinum* and also improved antioxidant capacities of bayberries. These authors reported a clear synergistic effect of the combination of methyl jasmonate and ethanol leading to approximately 10% decay in bayberries when each treatment led to 40 % decay and the controls showing 80% decay. Interestingly the combined treatments led to a better sensory appreciation of the fruit by a sensory panel than for all other treatment.

There is an infinite number of combinations, (i) as the number of potential individual treatments is great, (ii) as the combination order may vary (e.g. treatment A before treatment B, or B before A), and (iii) as the number of repeated applications over the pre- and post-harvest periods (e.g. one treatment every week over the ripening period or one treatment every fortnight). Combinatorial optimisation has an obvious interest in such approaches.

References

- Attarana, E., Zeierb, T.E., Griebela, T., Zeier, J. 2009. Methyl salicylate production and jasmonate signaling are not essential for systemic acquired resistance in Arabidopsis. *Plant Cell* 21, 954-971.
- Avissar, I., Pesis, E., 1991. The control of postharvest decay in table grapes using acetaldehyde vapours. *Ann. Appl. Biol.* 118, 229–237
- Babalar, M., Asghari, M., Talaei, A., Khosroshahi, A. 2007. Effect of pre- and postharvest salicylic acid treatment on ethylene production, fungal decay and overall quality of Selva strawberry fruit. *Postharvest Biol. Technol.* 105, 449-453.

- Barth, C., Moeder, W., Klessig D.F., Conklin, P.L. 2004. The timing of senescence and response to pathogens is altered in the ascorbate-deficient *Arabidopsis* mutant vitamin c-1. *Plant Physiol.* 134, 1784-1792.
- Beaulieu, J.C., Saltveit, M.E. 1997. Inhibition or promotion of tomato fruit ripening by acetaldehyde and ethanol is concentration dependent and varies with initial fruit maturity. *J. Amer. Soc. Hort. Sci.* 122, 392-398.
- Belhadj, A., Telef, N., Cluzet, S., Bouscaut, J., Corio-Costet, M.F., Merillon J.M. 2008a. Ethephon elicits protection against *Erysiphe necator* in grapevine. *J. Agric. Food Chem.* 56, 5781-5787.
- Belhadj, A., Telef, N., Saigne, C., Cluzet, S., Barrieu, F., Hamdi, S., Merillon, J.M. 2008b. Effect of methyl jasmonate in combination with carbohydrates on gene expression of PR proteins, stilbene and anthocyanin accumulation in grapevine cell cultures. *Plant Physiol. Biochem.* 46, 493-499.
- Camili, E.C., Benato, E.A., Pascholati, S.F., Cia, P., 2010. Fumigation of 'Italia' grape with acetic acid for postharvest control of *Botrytis cinerea*. *Rev. Bras. Frut.* 32, 436-443
- Chervin, C., Truett, J.K., Speirs, J. 1999. Alcohol dehydrogenase expression and alcohol production during pear ripening. *J. Amer. Soc. Hort. Sci.* 124, 71-75.
- Chervin, C., Elkereamy, A., Roustan, J.P., Faragher, J.D., Latche, A., Pech, J.C., Bouzayen, M., 2001. An ethanol spray at veraison enhances colour in red wines. *Aust. J. Grape Wine Res.* 7, 144–145.
- Chervin, C., Savocchia, S., Krstic, M., Serrano, E., van Heeswijck, R., 2005a. Enhancement of grape berry weight induced by an ethanol spray four weeks before harvest and effects of a night spray at an earlier date. *Aust. J. Exp. Agric.* 45, 731-734.

- Chervin, C., Westercamp, P., Monteils, G., 2005b. Ethanol vapours limit Botrytis development over the postharvest life of table grapes. *Postharvest Biol. Technol.* 36, 319–322.
- Chervin, C., Bouzambou, N., Latche, A., Pech, J.C., 2005c. Potential for ethanol vapours to limit table grape berry shatter and to limit ethylene evolution from clusters. *Acta Hortic.* 682, 513–518.
- Chervin, C., Tira-umphon, A., Terrier, N. Zouine, M. Severac, D. and Roustan, J.P. 2008. Stimulation of the grape berry expansion by ethylene and effects on related gene transcripts, over the ripening phase. *Physiol. Plant.* 134, 534–546.
- Chervin, C., Lavigne, D., and Westercamp, P. 2009. Reduction of gray mold development in table grapes by preharvest sprays with ethanol and calcium chloride. *Postharvest Biol. Technol.* 54, 115-117.
- Crisp, P., Wicks, T.J., Lorimer, M., Scott E.S. 2006. Novel controls for powdery mildew – greenhouse studies *Aust. J. Grape Wine Res.* 12, 192–202 & 203–211
- Cruz-Rus, E., Botella, M.A., Valpuesta, V., Gomez-Jimenez, M.C. (2010). Analysis of genes involved in L-ascorbic acid biosynthesis during growth and ripening of grape berries. *J. Plant Physiol.* 167, 739-748.
- DeBolt, S., Hardie, J., Tyerman, S., Ford, C.M. (2004). Composition and synthesis of raphide crystals and druse crystals in berries of *Vitis vinifera* L. cv. Cabernet Sauvignon: Ascorbic acid as precursor for both oxalic and tartaric acids as revealed by radiolabelling studies. *Aust. J. Grape Wine Res.* 10, 134–142.
- Dry, I.B., Feechan, A., Anderson, C., Jermakow, A.M., Bouquet, A., Adam-Blondon, A.-F., Thomas, M.R. 2010. Molecular strategies to enhance the genetic resistance of grapevines to powdery mildew. *Aust. J. Grape Wine Res.* 16, 94 -105

- Dubreuil-Maurizi, C., Trouvelot, S., Frettinger, P., Pugin, A., Wendehenne, D., Poinssot, B. 2010. β -aminobutyric acid primes an NADPH oxidase-dependent reactive oxygen species production during grapevine-triggered immunity. *Mol. Plant Microb. Int.* 23, 1012-1021.
- Durrant, W.E., Dong, X. 2004. Systemic acquired resistance. *Ann. Rev. Phytopathol.* 42, 185–209
- El Kereamy, A., Chervin, C., Souquet, J.M., Moutounet, M., Monje, M.C., Nepveu, F., Mondies, H., Ford, C.M., van Heeswijck, R., Roustan, J.P., 2002. Ethanol triggers grape gene expression leading to anthocyanin accumulation during berry ripening. *Plant Sci.* 163, 449–454.
- Gravel, V., Antoun, H., Tweddell, R.J., 2007. Effect of indole-acetic acid (IAA) on the development of symptoms caused by *Pythium ultimum* on tomato plants. *Eur. J. Plant Pathol.* 119, 457–462.
- James, D.G., Price, T.S. 2004. Field-testing of methyl salicylate for recruitment and retention of beneficial insects in grapes and hops. *J. Chem. Ecol.* 30, 1613-1628.
- Karabulut, O.A., Smilanick, J.L., Mlikota Gabler, F., Mansour, M., Droby, S., 2003. Nearharvest applications of *Metschnikowia fructicola*, ethanol, and sodium bicarbonate to control postharvest diseases of grape in central California. *Plant Dis.* 87, 1384–1389.
- Karabulut, O.A., Gabler, F.M., Mansour, M., Smilanick J.L. 2004. Postharvest ethanol and hot water treatments of table grapes to control gray mold. *Postharvest Biol. Technol.* 34, 169–177.
- Karabulut, O.A., Romanazzi, G., Smilanick, J.L., Lichter, A. 2005. Postharvest ethanol and potassium sorbate treatments of table grapes to control gray mold. *Postharvest Biol. Technol.* 37, 129-134.
- Kishimoto, K., Matsui, K., Ozawa, R., Takabayashi, J. 2008. Direct fungicidal activities of C6-aldehydes are important constituents for defense

- responses in *Arabidopsis* against *Botrytis cinerea*. *Phytochem.* 69, 2127–2132
- Kondo, S., Fukuda, K. 2001. Changes of jasmonates in grape berries and their possible roles in fruit development. *Sci. Hort.* 91, 275-288
- Kraeva, E., Andary, C., Carbonneau, A., Deloire, A. 1998. Salicylic acid treatment of grape berries retards ripening. *Vitis*, 37, 143-144.
- Lichter, A., Zutkhy, Y., Sonogo, L., Dvir, O., Kaplunov, T., Sarig, P., Ben-Arie, R., 2002. Ethanol controls postharvest decay of table grapes. *Postharvest Biol. Technol.* 24, 301–308.
- Lichter, A., Zhou, H.W., Vaknin, M., Dvir, O., Zutchi, Y., Kaplunov, T., Lurie, S., 2003. Survival and responses of *Botrytis cinerea* after exposure to ethanol and heat. *J. Phytopathol.* 151, 553–563.
- Lin, Z.F., Zhong, S.L., Grierson, D. 2009. Recent advances in ethylene research. *J. Exp. Bot.* 60, 3311-3336
- Martin, H., Maris, P. 2004. An assessment of the bactericidal and fungicidal efficacy of seventeen mineral and organic acids on bacterial and fungal food industry contaminants. *Sci. Alim.* 25, 105-127.
- Miyake, T., Shibamoto, T., 1995. Formation of acetaldehyde from L-ascorbic acid and related compounds in various oxidation systems. *J. Agric. food Chem.* 43, 1669-1672
- Nigro, F., Schena, L., Ligorio, A., Pentimone, I., Ippolito, A., Salerno, M.G., 2006. Control of table grape storage rots by pre-harvest applications of salts. *Postharvest Biol. Technol.* 42, 142–149.
- Park, S.W., Kaimoyo, E., Kumar, D., Mosher, S., Klessig, D.F. 2007. Methyl salicylate is a critical mobile signal for plant systemic acquired resistance. *Science.* 318 (5847), 113-116.
- Pesis, E., 2005. The role of the anaerobic metabolites, acetaldehyde and ethanol, in fruit ripening, enhancement of fruit quality and fruit deterioration. *Postharvest Biol. Technol.* 37, 1–19.

- Pesis, E., Frenkel, C., 1989. Effects of acetaldehyde vapors on postharvest quality of table grapes. *HortScience* 24, 315–317
- Slaughter A.R., Hamiduzzaman, M.M., Gindro, K., Neuhaus J.M., Mauch-Mani, B. 2008. Beta-aminobutyric acid-induced resistance in grapevine against downy mildew: involvement of pterostilbene. *Eur. J. Plant Pathol.* 122, 185-195.
- Smid, E.J., Hendriks, L., Boerrigter, H.A.M., Gorris, L.G.M., 1996. Surface disinfection of tomatoes using the natural plant compound trans-cinnamaldehyde. *Postharvest Biol. Technol.* 9, 343-350
- Song, J., Hildebrand, P.D., Fan, L.H., Forney, C.F., Renderos, W.E., Campbell-Palmer, L., Doucette, C. 2007. Effect of hexanal vapor on the growth of postharvest pathogens and fruit decay. *J. Food Sci.* 72, M108-M112.
- Spadaro, D., Garibaldi, A., Gullino, M.L. 2004. Control of *Penicillium expansum* and *Botrytis cinerea* on apple combining a biocontrol agent with hot water dipping and acibenzolar-S-methyl, baking soda, or ethanol application. *Postharvest Biol. Technol.* 33, 141-151.
- Suzuki, Y., Uji,T., Terai, H. 2004. Inhibition of senescence in broccoli florets with ethanol vapor from alcohol powder. *Postharvest Biol. Technol.* 31, 177-182.
- Sweetman, C., Deluc L.G., Cramer, G.R., Ford, C.M., Soole, K.L., 2009. Regulation of malate metabolism in grape berry and other developing fruits. *Phytochem.* 70, 1329–1344.
- Tadege M, Bucher M, Stahli W, Suter M, Dupuis I, Kuhlemeier C. 1998. Activation of plant defense responses and sugar efflux by expression of pyruvate decarboxylase in potato leaves. *Plant J.* 16, 661-671.
- Tripathi, P., Dubey, N.K., 2004. Exploitation of natural products as an alternative strategy to control postharvest fungal rotting of fruit and vegetables. *Postharvest Biol. Technol.* 32, 235-245.

- Van der Wolf, J. M., Birnbaum, Y., Van der Zouwen, P.S., Groot S.P.C. 2008. Disinfection of vegetable seed by treatment with essential oils, organic acids and plant extracts. *Seed Sci. Technol.* 36, 76-88.
- Viazis, S., Akhtar, M., Feirtag J., Diez-Gonzalez, F. 2011. Reduction of *Escherichia coli* O157:H7 viability on leafy green vegetables by treatment with a bacteriophage mixture and trans-cinnamaldehyde. *Food Microbiol.* 28, 149-157.
- Utama, I.M.S., Wills, R.B.H., Ben-Ye-Hoshua, S., Kuek, C., 2002. In vitro efficacy of plant volatiles for inhibiting the growth of fruit and vegetable decay microorganisms. *J. Agric. Food Chem.* 50, 6371–6377.
- Walz, A., Simon, O. 2009. β -aminobutyric acid-induced resistance in cucumber against biotrophic and necrotrophic pathogens. *J. Phytopathol.* 157, 356–361.
- Wang, K.T., Jin P., Shang H.T., Zheng Y.H. 2010. Effect of Methyl Jasmonate in Combination with Ethanol Treatment on Postharvest Decay and Antioxidant Capacity in Chinese Bayberries. *J. Agric. Food Chem.* 58, 9597-9604.
- Wen, P.F., Chen, J.Y., Kong, W.F., Pan, Q.H., Wan, S.B., Huang, W.D. 2005 Salicylic acid induced the expression of phenylalanine ammonia-lyase gene in grape berry. *Plant Sci.* 169, 928-934.