

Structural disorder in plant virus proteins

Eugénie Hébrard, Jocelyne J. Walter, Thierry Michon

▶ To cite this version:

Eugénie Hébrard, Jocelyne J. Walter, Thierry Michon. Structural disorder in plant virus proteins. 13. Rencontres de virologie végétale, Jan 2011, Aussois, France. 1 p./102, 2011. hal-02809208

HAL Id: hal-02809208 https://hal.inrae.fr/hal-02809208

Submitted on 6 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.





Aussois du 16 au 20 janvier 2011





53- Structural disorder in plant virus proteins

Eugénie HEBRARD¹, Jocelyne WALTER², Thierry MICHON²

The structure-to-function paradigm has changed with the discovery of intrinsically (or natively) disordered domains in proteins. By contrast to the vast majority of globular proteins, intrinsically disordered proteins (IDP) do not fold as a unique 3D structure. Under physiological conditions, IDPs exist as a dynamic ensemble of conformations containing no or only few secondary structure elements (helices and sheets). About 50% of the proteins coded by higher eukaryotes genomes are predicted to contain disordered domains ((Dunker et al. 2008).). Many of them are involved in cellular regulation processes requiring their interaction with several different partners. Structural disorder is likely to allow the surface plasticity necessary to this multi-partnership. Interestingly many proteins encoded by animal RNA viruses are also predicted to be rich in disordered domains. For instance, there are strong experimental evidences that the measles virus N-tail intrinsic disorder is associated with multipartners interactions, (Bourhis et al. 2004). Up to now, intrinsic disorder has been poorly investigated in plant viruses. VPgs are the only exception to date. VPgs are proteins linked to the 5' end of the genome of many plant and animal RNA viruses. Although sharing this common feature, VPgs are highly diverse in size, sequence and function. They are involved in key steps of the viral cycle such as genome replication, protein translation and viral movement in plant hosts (Sadowy et al. 2001). Using several disorder predictors, we showed that Lettuce mosaic virus (LMV) and Rice yellow mottle virus (RYMV) VPgs are predicted to possess disordered regions. We next extended theses results to 14 VPgs representative of the viral diversity. Disordered regions were predicted in all VPg sequences whatever the genus and the family suggesting that intrinsic disorder is a common feature of VPgs (Hébrard et al, 2009). We will show that intrinsic disorder is also predicted in other major proteins of several phytoviral species.

REFERENCES

Bourhis, J. M., K. Johansson, et al. (2004). "The C-terminal domain of measles virus nucleoprotein belongs to the class of intrinsically disordered proteins that fold upon binding to their physiological partner." Virus Res **99**(2): 157-167.

Dunker, A. K., C. J. Oldfield, et al. (2008). "The unfoldomics decade: an update on intrinsically disordered proteins." <u>BMC Genomics</u> **9 Suppl 2**: S1.

Sadowy, E., M. Milner, et al. (2001). "Proteins attached to viral genomes are multifunctional." Adv Virus Res 57: 185-262.

Hebrard, E., Y. Bessin, et al. (2009). "Intrinsic disorder in Viral Proteins Genome-Linked: experimental and predictive analyses." <u>Virol J</u> 6(1): 23.

¹ UMR 1097, Institut de Recherche pour le Développement (IRD), BP 64501, 34394 Montpellier cedex 5, France

² UMR GDPP, Institut de Biologie Végétale Moléculaire, Villenave d'Ornon, France