



HAL
open science

Genetic adaptation of an avian influenza A virus to swine cells

Vincent Bourret, Jon Lyall, Angélique Teillaud, Jinqi Fu, Guillaume Croville, Jean-Luc Guerin, Laurence Tiley

► To cite this version:

Vincent Bourret, Jon Lyall, Angélique Teillaud, Jinqi Fu, Guillaume Croville, et al.. Genetic adaptation of an avian influenza A virus to swine cells. IX International Congress of veterinary virology, European Society for veterinary virology (ESVV)., Sep 2012, Madrid, Spain. hal-02809684

HAL Id: hal-02809684

<https://hal.inrae.fr/hal-02809684>

Submitted on 6 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

GENETIC ADAPTATION OF AN AVIAN INFLUENZA A VIRUS TO SWINE CELLS

Vincent Bourret^{1,2,3}, Jon Lyall⁴, Angélique Teillaud^{2,3}, JinQi Fu¹, Guillaume Croville^{2,3}, Jean-Luc Guérin^{2,3} and Laurence Tiley⁴

¹Cambridge Infectious Diseases Consortium, Department of Veterinary Medicine, Madingley Road, Cambridge CB3 0ES, U.K.

²INRA, UMR 1225, IHAP, Toulouse, F-31076 France

³Université de Toulouse, INP, ENVT, Toulouse, F-31076 France

⁴Department of Veterinary Medicine, Madingley Road, Cambridge CB3 0ES, U.K.

Background and objectives The main epidemiological reservoir for influenza A viruses is in wild aquatic birds. From this reservoir, some strains can cross the species barrier and evolve as new lineages in other susceptible hosts. Swine are among the sensitive species, with influenza A virus being one of its primary respiratory pathogens. Swine are also considered to be possible intermediary hosts enabling adaptation of avian influenza strains to humans. The objective of this project was to study the adaptation and genetic evolution of an avian influenza strain in swine respiratory cells.

Methods We first adapted an influenza virus field-strain isolated from a duck to swine respiratory cells by repetitive passaging and tested the consequences on growth kinetics in swine, avian, and human cells. We then sequenced the complete parental and adapted viral populations, combining Sanger sequencing and deep-sequencing approaches to study the genetic changes occurring both at the consensus and at the subpopulation levels. We finally set up a prototypic virus rescue system for reverse genetics study of that influenza strain and its swine cell-adapted descendent.

Results As a result of passaging, growth kinetics were improved in swine cells, while they remained mostly unchanged in duck fibroblasts and in a human bronchiolar epithelial cell line; passaging in swine cells also resulted in a marked growth kinetics improvement in chicken cells. The parental field sample was found to contain genes from co-circulating strains present as minority alleles. This enabled reassortment events involving segments 2, 7 and 8 upon culture in swine cells. These reassortments, together with two point mutations on segment 4, resulted in the selection of a near-homogeneous swine cell-tropic virus. Testing rescued viruses carrying combinations of the observed adaptive changes showed that, in the parental virus' genetic background, two non-synonymous point mutations on segment 4 could restore the adapted virus' fast growth phenotype. Reassortment of segment 2 alone, resulting in seven amino acid changes in PB1 and 11 amino acid changes in PB1-F2, seemed to achieve a similar effect.

Conclusions In the context of a naïve duck influenza virus strain, two amino acid changes away from the receptor binding site on the viral haemagglutinin could enhance growth in swine cells. Further, growth kinetics improvement in chicken cells following passaging in swine cells suggested a possible role for swine as an intermediary host for adaptation of an influenza strain from wild waterfowl to domestic chickens.