



HAL
open science

A first assessment of the SMOS data in southwestern France using in situ, airborne and model soil moisture estimates

Clément Albergel, Elena Zakharova, Jean-Christophe Calvet, Mehrez Zribi, Mickaël Pardé, Jean-Pierre Wigneron, Nathalie Novello, Yann H. Kerr

► **To cite this version:**

Clément Albergel, Elena Zakharova, Jean-Christophe Calvet, Mehrez Zribi, Mickaël Pardé, et al.. A first assessment of the SMOS data in southwestern France using in situ, airborne and model soil moisture estimates. 29. Progress in Electromagnetics Research Symposium, PIERS 2011, Mar 2011, Marrakech, Morocco. 1 p. hal-02810056

HAL Id: hal-02810056

<https://hal.inrae.fr/hal-02810056>

Submitted on 6 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A First Assessment of the SMOS Data in Southwestern France Using in Situ, Airborne and Model Soil Moisture Estimates

Clément Albergel¹, Elena Zakharova¹, Jean-Christophe Calvet¹, Mehrez Zribi²,
Mickaël Pardé⁴, Jean-Pierre Wigneron⁴, Nathalie Novello⁴, and Yann Kerr²

¹CNRM-GAME, Météo-France, CNRS, 42 avenue Gaspard Coriolis, Toulouse, France

²CESBIO, CNES/CNRS/IRD/UPS(UMR5126), Toulouse, France

³LATMOS, Paris, France

⁴INRA, EPHYSE, Villenave d'Ornon, France

Abstract— Passive microwave remote sensing of soil moisture has been at the center of attention of many research programs, for several decades. Various airborne and in situ radiometers have been developed, showing the high potential of L-band measurements for the estimation of surface parameters. The Soil Moisture and Ocean Salinity (SMOS) satellite mission, based on an aperture synthesis L-band radiometer was successfully launched on November 2009. In the context of a validation campaign for the SMOS mission, intensive airborne and in situ observations were performed in southern France for the SMOS CAL/VAL, from April to May 2009 and April to July 2010. The CAROLS “Cooperative Airborne Radiometer for Ocean and Land Studies” L band radiometer was designed, built and installed on board a dedicated French ATR42 research aircraft. In spring 2010, soil moisture observations from 12 stations of the SMOSMANIA network of Météo-France were complemented by airborne observations of the CAROLS L-band radiometer, following an Atlantic-Mediterranean transect in southwestern France.

The first CAL/VAL results include a comparison between the surface soil moisture retrieved from the brightness temperatures measured during the CAROLS flights with either in situ (SMOSMANIA network and additional measurements) or synthetic soil moisture simulated by the SIM (Safran-Isba-Modcou) hydrometeorological model. It also includes a comparison between L-band brightness temperatures from CAROLS and SMOS, and between soil moisture values retrieved from CAROLS and SMOS. The latter are compared with the in situ observations of soil moisture. Also, simulated soil moisture estimates produced by the ISBA land surface model are used in the cross-evaluation and the biophysical variables produced by ISBA are used as input to the CMEM microwave emission model to simulate brightness temperatures.