



HAL
open science

Relationship between phosphorus availability and symbiotic n₂ fixation rate in terrestrial ecosystems

Laurent Augusto, Florian Delerue, Anne Gallet-Budynek, D. Achat

► To cite this version:

Laurent Augusto, Florian Delerue, Anne Gallet-Budynek, D. Achat. Relationship between phosphorus availability and symbiotic n₂ fixation rate in terrestrial ecosystems. Agu Fall Meeting 2011, Dec 2011, San Francisco, United States. 1 p., 2011. hal-02810066

HAL Id: hal-02810066

<https://hal.inrae.fr/hal-02810066v1>

Submitted on 6 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Phosphorus availability in terrestrial ecosystems does not constrain symbiotic N₂ fixation rate

Laurent Augusto¹, Florian Delerue^{1,2}, Anne Gallet-Budynek¹, David L. Achat² - (laugusto@bordeaux.inra.fr)

Introduction and Objective

Symbiotic N₂ fixation (SNF) is a major input flux in terrestrial ecosystems. The process has been extensively studied and the factors controlling it, like phosphorus (P), are well identified. Phosphorus is also limiting plant growth in many parts of Earth and is thus of great importance for carbon sequestration in the current context of global change. Although the interactions between P and SNF have been identified as one of the few important research areas on N₂ fixation, the way P controls SNF is still a source of controversy and remains unclear. Considering the importance of SNF for ecosystems, our aim was to elucidate the dependency of this process on P availability in soils of the world.

Material and Methods



We compiled published studies which measured the SNF of plants under different levels of P supply. The results of this compilation were put into a global perspective by comparing them with an extensive analysis of soil phosphorus status throughout the world.

• Compilation of SNF studies under different levels of P supply


We used the ISI Web Of Science database to find published literature on the effect of P supply on SNF of plants grown in natural soils (in the field or in pots) or artificial irrigated substrates (e.g. perlite or hydroponics).

• Worldwide data on P in soils

We used the ISI Web Of Science database to find published literature characterizing water-P (PO₄ soluble in soil water) or labile-P (water soluble PO₄ + soil P fractions available for plants; Hedley fractionation method) in the top layer of the mineral soil or in the forest floor. Data were collected throughout the world. Each continent (North America+West Indies+Central America; South America; Europe; Africa; Asia; Oceania+Pacific) represented 9–33% of our dataset. All biomes, climates (continental to tropical), land uses (croplands, forests, grasslands and savannas) and soil types were represented.



Water-P
(water soluble PO₄)
508 values



Labile-P
(bioavailable P)
631 values

Results and Discussion

