



Overview of the high spatial and temporal resolutions MISTIGRI mission in the Thermal Infrared

Jean-Pierre Lagouarde, Michel Bach, Gilles Boulet, Xavier Briottet, Selma Cherchali, Isabelle Dadou, Gérard Dedieu, Olivier Hagolle, Frédéric Jacob, Françoise Nerry, et al.

► To cite this version:

Jean-Pierre Lagouarde, Michel Bach, Gilles Boulet, Xavier Briottet, Selma Cherchali, et al.. Overview of the high spatial and temporal resolutions MISTIGRI mission in the Thermal Infrared. GlobTemperature - User Consultation Meeting, Jun 2012, Edinburgh, United Kingdom. n.p., 2012. hal-02810946

HAL Id: hal-02810946

<https://hal.inrae.fr/hal-02810946>


Submitted on 6 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Overview of the high spatial and temporal resolutions MISTIGRI mission in the Thermal Infrared

J.-P. Lagouarde ⁽¹⁾, M. Bach ⁽²⁾, G. Boulet ⁽³⁾, X. Briottet ⁽⁴⁾, S. Cherchali ⁽²⁾, I. Dadou ⁽⁵⁾, G. Dedieu ⁽³⁾, O. Hagolle ⁽³⁾, F. Jacob ⁽⁶⁾, F. Nerry ⁽⁷⁾, A. Olioso ⁽⁸⁾, C. Ottlé ⁽⁹⁾, V. Pascal ⁽²⁾, J.L. Roujean ⁽¹⁰⁾, J.A. Sobrino ⁽¹¹⁾



- MISTIGRI = MicroSatellite for Thermal InfraRed Ground Surface Imaging
- Cooperation between France (CNES) and Spain (Univ. Valencia)



- High spatial resolution (~ 50m)
- High revisit capacities (1 day)

- Definition and consolidation of mission specifications during a A phase (sept 2009 - end 2011)

SCIENTIFIC OBJECTIVES


Monitoring of energy and water budgets of continental biosphere

- Agriculture/forestry/natural vegetation productivity/ irrigation
- Biogeochemical cycles / carbon budgets / soil pollution
- Hydrology/ water management



Urban environment monitoring


- Urban climatology and heat waves / pollutants diffusion and air quality / anthropogenic fluxes
- Urban heat island (UHI) / welfare, health of inhabitants
- Urban hydrology



Difference between daytime and nighttime LST over the Madrid city (DESIREX, 2008) retrieved from Airborne Hyperpectral Scanner (AHS) data (Sobrino et al., 2009)

Monitoring of coastal and continental waters

- Monitoring of the coastal areas
- Submesoscale activity in coastal and open ocean
- Monitoring of lakes and rivers / floods
- Air-sea fluxes, GHG, biogeochemical cycles
- Lagunaes / estuaries / deltas
- Biological activity and productivity / discharges / water quality



Modeled SST California Upwelling System (a) with zooms at 6 km (b) and 0.75 km (c) resolutions (Capet et al., 2008)

Other applications

- Vulcanology
- Risk assessment, coalmine and peat fires, epidemic outbreaks...
- ... cf. ESA Fuegosat Synthesis Study, 2010

Ref : Lagouarde et al., IJRS, 2012, in press


REMOTE SENSING CONTEXT

Need of a mission combining **high** resolution and **high** revisit in the TIR


Landsat TM/ETM+	16 days	60 - 120 m
LDCM	16 days	100 m
ASTER	16 days	90 m
Meteosat MSG, GOES	15mn	2.5 - 5 km
AVHRR	2-4/day	1 km
MODIS	1/day	1 km

STRATEGY of the MISSION

Mission based on a network of experimental sites worldwide (similarity with Venus strategy)



SPATIAL RESOLUTION (~ 50 m)



A 60 m resolution allows one to discriminate the contribution to actual evaporation of 2 crops in Iowa (after Kustas et al., 2004)


Upper limit of the resolution imposed by the **mean size of fields** to be observed

Kustas et al., 2004, Garriques et al., 2006, Agam et al., 2007, LDCM...


< 100m

Lower limit of the resolution must cope with temporal fluctuations of surface temperature induced by surface (SBL) and planetary (PBL) boundary layers atmospheric turbulence

> 40m




Cumulative histogram of plot surfaces for 4 crops, in S-W France (statistics based on 80 000 plots): 75 % of surface is made of plots > 5ha



Impact of PBL and SBL turbulence on temporal fluctuations of Ts (pine forest) at 5 and 200 m spatial resolutions
Ref : Lagouarde et al., 2012, IGARSS Munich & Lagouarde et al., RSE submitted

ORBIT and ACCESSIBILITY



561 km, 1 day


720 km, 2 days

REVISIT and OVERPASS TIME

Revisit imposed by both the temporal variability of observed processes (fluxes, drying events) and the cloudiness (data availability)


Revisit also imposed by the performances and accuracy expected from models (and by frequency and duration of stress periods)

Overpass time around 12 UTC best suited for fluxes monitoring




Analysis of hourly solar irradiance data from INRA AGROCLIM network 1993-2009, over 2 locations in France and for 3 revisit assumptions

Analysis of MODIS cloud masks (2000-2009 over Europe for summer (June 1st - August 31st), for 2 revisit assumptions




Analysis of the impact of the revisit on the accuracy of daily AET retrievals (based on a simulated dataset 1950 - 2100, A1B IPCC scenario, with a selection of cloudfree days)



Sensitivity of the efficiency of reconstruction of daily evapotranspiration (from simulation of AET time series) to time of acquisition (Delogu et al., 2012, Gentile, 2007)

Ref : Olioso et al., Lagouarde et al., Boulet et al., RAQRS III, 2012 Valencia; Delogu et al., HESSD 2012

DEVELOPMENT of AET ALGORITHMS



Mapping latent heat flux (LE) over Crau plain (S-E France) from LANDSAT-7 data

Ref : Olioso et al., 2012, TOSCA Conf., Paris

INSTRUMENT

Band	Central wavelength (μm)
TIR-3	10.3
TIR-4	11.5
TIR-1	8.6
TIR-2	9.1

TIR instrument :

- Uncooled ULIS microbolometer 640x480 array (25μm pitch)
- NedT 0.5K (0.2K aimed at)
- Absolute accuracy = 1K

A simulator has been built to specify channels and to evaluate accuracy of Ts derived from TES algorithm. Merging TIR1 and TIR2 for reducing noise currently investigated.

VNIR instrument:


- 1D CCD array
- 4 bands

Use:

- TIR registration
- cloud detection
- vegetation indices
- surface emissivity first guess for TES
- disaggregation of TIR

Band	Central wavelength (μm)
Blue	0.450
Red	0.670
Near-InfraRed	0.865
Near-InfraRed	0.910

Ref : Lesaignoux et al., 2012, RSE submitted; Briottet, 2012, TOSCA Conf., Paris



MISSION ARCHITECTURE

Instrumental concept

- Satellite 1.5 slow down (pitch rotation of the platform)
- On-ground TDI-like binning (30 lines)
- On-board blackbodies for TIR calibration (283 and 313K)

Platform

- Compatibility of the MYRIADE CNES platform with the mission
- Myriade data storage 16 Gbits

Feasibility study

- Evaluation of performances of microbolometers and of the TIR instrument currently being performed at CNES
- First instrumental design proposed by TAS

Ground segment

- Already largely existing
- Venus developments

⁽¹⁾ INRA-EPHYSE, 71 av. E. Bourlaux BP 81, F-33883 Villenave d'Ornon

⁽²⁾ CNES, 18 av. E. Belin, F-31401 Toulouse Cedex 9

⁽³⁾ CESBIO (CNRS, UPS, CNES, IRD), 18 av. E. Belin, F-31401 Toulouse cedex 9

⁽⁴⁾ ONERA-DOTA, BP 74025, 2, av. E. Belin, F-31055 Toulouse cedex 4

⁽⁵⁾ LEGOS, OMP, 14 Avenue E. Belin, F-31400 Toulouse cedex

⁽⁶⁾ IRD-LISAH, 2 place P. Viala, F-34060 Montpellier cedex 1

⁽⁷⁾ LSIIT/TRIO, Bd Sébastien Brant, BP 10413, F-67412 Illkirch cedex

⁽⁸⁾ INRA-EMMAH, UMR 1114, Domaine St Paul, Site Agroparc, F-84 914 Avignon

⁽⁹⁾ LSCE, Centre d'Etudes de Saclay, Orme des Merisiers, F-91191 Gif-sur-Yvette

⁽¹⁰⁾ CNRM-GAME, Météo-France, 42, av. G. Coriolis, F-31057 Toulouse

⁽¹¹⁾ UCG/ IPL, Univ.Valencia P. O. Box 22085, E-46071 Valencia (Spain)