



First evaluation of the SMOS observations over the VAS site: comparing different retrieval approaches

Jean-Pierre Wigneron, Nathalie Novello, Christophe Moisy, Dominique Guyon, Heather Lawrence, Mike Schwank, Ernesto Lopez-Baeza, Cristina Millan, Maciej Miernecki, Yann H. Kerr, et al.

► To cite this version:

Jean-Pierre Wigneron, Nathalie Novello, Christophe Moisy, Dominique Guyon, Heather Lawrence, et al.. First evaluation of the SMOS observations over the VAS site: comparing different retrieval approaches. IEEE International Geoscience and remote Sensing Symposium - IGARSS 2012, IEEE, Jul 2012, Munich, Germany. n.p. hal-02811221

HAL Id: hal-02811221

<https://hal.inrae.fr/hal-02811221>

Submitted on 6 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

First evaluation of the SMOS observations over the VAS site: comparing different retrieval approaches

J-P Wigneron, N. Novello, H. Lawrence, C. Moisy, D. Guyon, Inra Ephyse, Bordeaux France
wigneron@bordeaux.inra.fr, tel 33 5 57 12 24 19, FAX 33 5 57 12 24 20

M. Schwank, Helmholtz Centre Potsdam, GFZ German Research Centre for Geosciences, Germany

E. Lopez-Baeza, M. Mierneck, C. Millan, , UV, Valencia, Spain

Y H. Kerr, A. Al Bitar, F. Cabot, S. Juglea, D. Leroux, A. Mialon, P. Richaume, CESBIO, Toulouse France,

J. P. Grant & T. Casals, ESA/ESTEC, Noordwijk, The Netherlands

P. Waldteufel, IPSL-LATMOS, Paris, France

K. Saleh, CDTI, Madrid, Spain

A. Mahmoodi, Array Systems, USA

S. Delwart, S. Mecklenburg, ESA, ESRIN, Frascati, Italy

Abstract

The SMOS (Soil Moisture and Ocean Salinity) mission was launched on November 2, 2009. Over the land surfaces, simultaneous retrievals of surface soil moisture (SM) and vegetation characteristics are made from the multi-angular and dual polarization SMOS observations by inverting the L-MEB (L-band Microwave Emission of the Biosphere) model [1-2]. Preliminary analyses evaluating the SMOS observations in terms of Brightness Temperatures (TB) and Level-2 products (SM and vegetation optical depth TAU) were carried out over several sites of all continents.

Here, the study is based on SMOS observations and in situ measurements carried out in 2010 over one of the main SMOS calibration/validation sites in Europe: the VAS (Valencia Anchor Station) site in the region of Utiel-Requena, close to Valencia, Spain. The main vegetation types in the region are vineyards (cover fraction ~60%), natural Mediterranean vegetation (pine trees and shrubs ~30%) and orchards. The SMOS observations were analysed in conjunction with those carried out by the L-band ELBARA II radiometer over a vineyard which is representative of the main land use of the VAS site [3]. Retrievals of SM and TAU based on the SMOS observations (Level-2 products) and the ELBARA II instrument were compared and evaluated against in situ measurements and simulations of SM by atmospheric models.

Even though there is a very large difference in the spatial scale of the observations made from ELBARA II (a plot within a vineyard field) and SMOS (~a circle with a 40 km diameter), a good agreement was found between the TB observations ($R^2 > 0.88$) at both polarizations.

ICAR 88 2012

Paper 1425 - 17-27 July, invited sessions.

A good agreement was found too between the retrieved time variations in SM over the site computed from both the SMOS and ELBARA II observations. Retrievals computed from the SMOS observations were obtained a) from the official ESA SM products computed with the Level 2 algorithm and b) from a simplified retrieval process based on the L1C Tb products and assuming homogeneous soil/vegetation conditions over the SMOS pixel and neglecting the effects due to the SMOS synthetic lobe patterns (referred to as the WEF function).

It was found the SMOS Level-2 SM products underestimated the SM values retrieved from ELBARA II by a bias of about -0.20% (Fig. 1a). Conversely, retrievals computed from the SMOS data using a simplified L-MEB approach, that did not consider the pixel heterogeneity, produced very satisfying results (Fig. 1b) with a much lower bias (Bias \sim -0.06 m³/m³, R²=0.76) when compared to retrievals computed from the ELBARA II observations. Current studies investigate this result which could provide interesting hints to explain the dry bias which is obtained with the Level- 2 products over some specific geographical areas.

The time variations in the retrieved TAU computed from the ELBARA II observations could be closely related to the time variations in the NDVI vegetation index, obtained from the MODIS observations in the optical domain (TAU = 0.61. NDVI -0.11, with R² = 0.61). Conversely, the time variations in the SMOS Level-2 TAU product did not provide clear patterns that could be related to the vegetation dynamics and seemed to be strongly affected by the effects of the microwave interferences (RFI) over the VAS site.

The comparison between the SM and TAU retrievals obtained from (i) the Level-2 algorithm and (ii) the simplified L-MEB approach allowed making recommendations about potential future improvements of the Level-2 algorithms.

References:

- [1] Kerr Yann H., Philippe Waldteufel, Philippe Richaume, Jean Pierre Wigneron, Paolo Ferrazzoli, Ali Mahmoodi, Ahmad Al Bitar, François Cabot, Claire Gruhier, Silvia Juglea, Delphine Leroux, Arnaud Mialon, Steven Delwart, 'The SMOS soil moisture retrieval algorithm', IEEE Trans. Geosc. Remote Sens., in press
- [2] Wigneron J-P, Y. Kerr, P. Waldteufel, K. Saleh, M.-J. Escorihuela, P. Richaume, P. Ferrazzoli, P. de Rosnay, R. Gurney, J.-C. Calvet, J.P. Grant, M. Guglielmetti, B. Hornbuckle, C. Mätzler, T. Pellarin, M. Schwank, 'L-band Microwave Emission of the Biosphere (L-MEB) Model: Description and calibration against experimental data sets over crop fields', Remote Sens. Env., 107, p. 639-655, 2007.
- [3] Schwank M, E. Lopez-Baeza, J.-P. Wigneron, I. Völksch, Y. Kerr, C. Mätzler, 'L-Band Radiative Properties of Vine-Vegetation at the SMOS Cal/Val Site MELBEX III', IEEE Trans. Geosc. Remote Sens. , in press

Figures




Fig. 1 ab Comparison between retrieved Soil Moisture (SM) values obtained from the *in situ* ELBARA-II TB observations with SM values retrieved from the SMOS observations over the VAS site in 2010. These latter were obtained a) using official ESA SM products computed with the Level 2 algorithm b) using a simplified retrieval process based on the L1C TB products assuming homogeneous soil/vegetation conditions over the SMOS pixel.

