



**HAL**  
open science

## Comparison of spring $^{15}\text{N}$ uptake in oak and beech saplings submitted to shading and summer defoliation

Pascale Maillard, Luis Valenzuela Nunes, Dominique Gerant, Claude C. Brechet, Rana El-Zein, Nathalie Bréda

### ► To cite this version:

Pascale Maillard, Luis Valenzuela Nunes, Dominique Gerant, Claude C. Brechet, Rana El-Zein, et al.. Comparison of spring  $^{15}\text{N}$  uptake in oak and beech saplings submitted to shading and summer defoliation. Joint European Stable Isotope User Meeting, JESIUM 2008, Aug 2008, Presqu'île de Giens, France. 1 p. hal-02811721

**HAL Id: hal-02811721**

**<https://hal.inrae.fr/hal-02811721>**

Submitted on 6 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Session 3-Poster**

**Comparison of spring  $^{15}\text{N}$  uptake in oak and beech saplings submitted to shading and summer defoliation**

Pascale Maillard<sup>1</sup>, Luis Valenzuela Nunes<sup>2</sup>, Dominique Gérard<sup>1</sup>, Claude Brechet<sup>1</sup>, Rana El Zein<sup>1</sup> and Nathalie Breda<sup>1</sup>

<sup>1</sup>IFR 110, "Génomique Ecophysiologie et Ecologie fonctionnelles", INRA-UHP Nancy1, Faculté des Sciences et Techniques, 54506 Vandoeuvre, France

<sup>2</sup>Centro Nacional de Investigación Disciplinaria en la Relación Agua-Suelo-Planta-Atmósfera. Gómez Palacio, Durango, México.

Tree species are characterized by a long-term and cyclic growth. The storage of reserves and their remobilisation is one of the major functions ensuring permanence facing a variable environment. A part of reserves will be consumed allowing trees to face to various abiotic and biotic stresses such as defoliation and to regenerate new structures. In this context, storage of reserves will be a fundamental process that depends on both nutrient availability and environmental factors as light. Light availability will be determinant for carbon and nitrogen uptake and growth. However, light demand vary with species: beech is tolerant to shading whereas oak prefers full light.

A summer defoliation (July 2004; removal of 70% of leaves) was simulated on two-year-old oak and beech trees grown either in full light or in shading conditions (40% of incident light) in a nursery at INRA (Champenoux, France). The following spring (year 3) at bud bursting, trees were watered twice a week with a complete nutrient solution enriched in  $^{15}\text{N}$  ( $^{15}\text{NH}_4^{15}\text{NO}_3$ ,  $^{15}\text{N} > 99\%$ ; Spectra Stable Isotopes, Columbia, M.D., USA). Two hypotheses were tested 1) defoliation applied in summer would lead to a major N loss imposing an increase of  $^{15}\text{N}$  uptake next spring to ensure new growth and 2) responses to defoliation would vary according to light tolerance of species.

Shading did not alter N accumulation (year 2) but increased  $^{15}\text{N}$  uptake in spring (year 3) in both oak and beech. In contrast, responses to defoliation differed according species. Nitrogen storage was decreased in oak but not in beech (year 2) and N uptake (year 3) was increased in beech only.

As a consequence, defoliation created a N imbalance not compensated by an increase of uptake in oak trees whereas N stock of defoliated beech was restored at the end of the year 2 and  $^{15}\text{N}$  uptake was increased the next spring. In forest conditions, changes in  $^{15}\text{N}$  uptake due to silvicultural practices such as thinning that modifies light environment are investigated too on 40-year-old oak trees.