



HAL
open science

Modelling climate change impacts on production, carbon sequestration and water demand of livestock systems

Anne-Isabelle Graux, Romain Lardy, David Hill, Jean-François Soussana

► To cite this version:

Anne-Isabelle Graux, Romain Lardy, David Hill, Jean-François Soussana. Modelling climate change impacts on production, carbon sequestration and water demand of livestock systems. Climate Change Congress: Global Risks, Challenges and Decisions, Mar 2009, Copenhagen, Denmark. 2009. hal-02811933

HAL Id: hal-02811933

<https://hal.inrae.fr/hal-02811933v1>

Submitted on 6 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Modelling climate change impacts on production, carbon sequestration and water demand of livestock systems

Graux A.I.⁽¹⁾, Lardy R.⁽¹⁾, Hill D.⁽²⁾, Soussana J.F.⁽¹⁾

⁽¹⁾ INRA, UR874, UREP, Grassland Ecosystem Research, Clermont-Ferrand (France) aigraux@clermont.inra.fr

⁽²⁾ ISIMA/LIMOS, UMR CNRS 6158, Université Blaise Pascal, Aubière (France)

Introduction

Although the increase of atmospheric CO₂ concentration is expected to enhance plant growth and to increase carbon sequestration in grassland ecosystems, an increase in future drought and heatwave events could turn grasslands into carbon sources, contributing to positive carbon-climate feedbacks, and could also lead to a decrease of yields, as it has already been observed in 2003. The aims of this work are: a) to simulate the impacts of climate change on production, carbon sequestration and water demand; b) to test the sustainability of contrasted grassland systems; c) to identify sources of uncertainty and of variability in climate projections.

Materials and Methods

To simulate the impacts of climate change on livestock systems, we used a biogeochemical grassland ecosystem “PASIM” [1] that simulates fluxes of carbon, nitrogen, water and energy at the soil-plant-animal-atmosphere interface (Fig. 1). We compared impact projections on production, carbon sequestration and water sensitivity of livestock systems, for a range of IPCC SRES scenarios (A1B, A2 & B1) of the ARPEGE model and of downscaling methods (anomalies, variable correction and statistical disaggregation [2]) for the A1B scenario, at 2 French sites: Colmar and Toulouse (Fig. 2). To separate climate change and CO₂ concentration effects: i) the A1B scenario was simulated with a constant CO₂ concentration (325ppm) corresponding to 1970, ii) a constant climate control was simulated by sampling years between 1970 and 2000.




Figure 1 : The PASIM model




Figure 2 : Grassland sites for projections of impacts : a) location, b) evolutions of i) mean air temperature (T), ii) precipitations (P) and atmospheric CO₂ concentration

Name	Management	Grassland type	Fertilization [kg N.ha-1]	Irrigation [% of needs]	Cutting dates				Grazing periods		Stocking density [LSU.ha-1]
					15-apr.	30-jun.	15-aug.	15-oct.	20/07-05/08	15/10-01/11	
GI	Sown	100 % Festuca aru. or Lolium per.	320	80	15-apr.	30-jun.	15-aug.	15-oct.	-	-	-
G			200	0	15-apr.	30-jun.	15-aug.	15-oct.	-	-	-
PPP	Permanent	20% of Trifolium rep.	200	0	15-apr.	01-jun.	-	-	20/07-05/08	15/10-01/11	1,5
PPE		5% of Trifolium rep.	0	0	15-apr.	01-jun.	-	-	20/07-05/08	15/10-01/11	0,8

Figure 3 : Simulated livestock systems

Moreover, we compared 6 livestock systems, differing by grassland type (permanent or sown, grass species, legume fraction) and management (fertilization, irrigation, cutting and/or grazing) (Fig. 3). The agricultural practices were set at their current level. Soil organic matter was initialized at equilibrium with the climate in the 1950's.

Results




Figure 4: Projected climate change impacts of a) water stress, b) forage dry matter (DM) c) nitrogen yields and d) soil carbon at Toulouse for the “PPE system”, comparing 3 IPCC SRES scenarios climate scenarios obtained with the variable correction downscaling method (A), or separating the climate change and CO₂ concentration effects (B). Error bars show the interannual variability during the period

According to the PASIM model, when averaged over 30 years and compared to the reference period (1970-2000):

- Water requirements for grassland ecosystems tend to increase in far future, especially for the more pessimistic A2 scenario,
- Production does not vary significantly. Results emphasize a large effect of interannual climate variability for grassland production and there is no indication of a change in interannual yield variability,
- Soil organic carbon (SOC) content increases markedly through time, indicating that the increased atmospheric CO₂ concentration combined with warming leads to carbon sequestration.

These evolutions are observed for the 2 sites and the different livestock systems, with differences in mean values.

The separation of climate change and CO₂ concentration effects underlines the high weight of atmospheric CO₂ increase on PASIM results.

Conclusions

These results should be taken with caution, in regard to current model limitations for simulations of high temperatures and of severe droughts. Model tests and parametrisation against data from climate change experiments are planned and should allow more realistic projections of climate change impacts. Simulations at 10 other French sites are also planned and will complete these first results.