



HAL
open science

Modelling as a tool to identify key measures to reduce Salmonella prevalence in slaughter pigs

Amandine Lurette, Suzanne Touzeau, Pauline Ezanno, Thierry Hoch,
Christine Fourichon, Henri H. Seegers, Catherine C. Belloc

► **To cite this version:**

Amandine Lurette, Suzanne Touzeau, Pauline Ezanno, Thierry Hoch, Christine Fourichon, et al..
Modelling as a tool to identify key measures to reduce Salmonella prevalence in slaughter pigs. 20.
International Porcine Veterinary Society Congress, Jun 2008, Durban, South Africa. hal-02812297

HAL Id: hal-02812297

<https://hal.inrae.fr/hal-02812297>

Submitted on 6 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

MODELLING AS A TOOL TO IDENTIFY KEY MEASURES TO REDUCE *SALMONELLA* PREVALENCE IN SLAUGHTER PIGS

*Lurette A, **Touzeau S, *Ezanno P, *Hoch T, *Fourichon C, *Seegers H, *Belloc C
 *INRA, ENVN, UMR1300 Bio-agression Epidémiologie et Analyse de Risque, BP40706, F-44307 Nantes, France
 INRA, UR341 Mathématiques et Informatique Appliquées, F-78350 Jouy-en-Josas, France

• Introduction

Since the control of *Salmonella* carriage is a major public concern, a new European regulation dealing with zoonosis aimed at reducing *Salmonella* in the pork food chain is required. At the herd level, several control measures are available to reduce the seroprevalence (Farzan *et al.*, 2006, Roesler *et al.*, 2006). Additional potential measures can be derived from what is used in other species, for instance vaccination. These control measures are characterized (i) by which mechanisms they modify, (ii) by which animals are targeted, (iii) during which period, and (iv) how efficiently they act. The aim of this study was to demonstrate how a modelling approach can help to apply a measure in an optimal way.

Material and method

In this study, a stochastic mathematical model representing the farrow-to-finish herd dynamics (Lurette *et al.*, 2008) as well as the *Salmonella* spread was used. We tested three reduction levels for two parameters that are assumed to be affected by the potential control measures: the infection probability and the *Salmonella* quantity shed. The measures were applied either to sows or pigs during their whole lifetime, or only to lactating sows, post-weaning pigs and finishing pigs. The assessment criterion used to compare the effect of the measures implemented was the percentage of groups of delivered pigs with a seroprevalence lower than 5% ($p_{5\%}$).

Results

Scenarios which led to more than 50% of groups of

delivered pigs with a low seroprevalence differed according to the parameter considered, its level of reduction and the animals targeted. Indeed, reducing the infection probability by 50% for all pigs induced a $p_{5\%}$ higher than 50%. This value was only obtained with a 90% reduction when applied to finishing pigs (Fig. 1). A $p_{5\%}$ higher than 50% was reached by reducing the quantity of *Salmonella* shed only when applied to all sows and for a reduction of at least 50% (Fig. 2).




Figure 1 Percentage of groups of delivered pigs with *Salmonella* seroprevalence lower than 5% according to the level of reduction of the infection probability, the duration of the measure and the type of animal concerned. Values with the same letter are not significantly different




Figure 2 Percentage of groups of delivered pigs with *Salmonella* seroprevalence lower than 5% according to the level of reduction of the quantity of *Salmonella* shed, the duration of the measure and the type of animal concerned. Values with the same letter are not significantly different

Discussion

These results show that a modelling approach is a helpful tool to define the key features of potential *Salmonella* control measures at the herd level: which animals need to be targeted, during which period, and which efficiency the measure should reach to obtain a given *Salmonella* prevalence improvement.

Scenarios identified by the model results can be experimented to assess which level of parameter reduction can be reached by a measure implementation. Moreover, experimentations have shown that measures act on several mechanisms at once (for example vaccination). These measures could be therefore assessed with the model by testing the combined effect of several parameter reductions.

References

Farzan, A., et al.(2006) *Prev. Vet. Med.*, 73(4), 241-254. Lurette, A. et al (2008) *Animal*, 2(1), 105-116. Roesler, U., et al (2006) *J. Vet. Med.*, 53(5), 224-228.