



HAL
open science

Modelling Salmonella spread in a pig farm under three biosecurity strategies

Amandine Lurette, Catherine C. Belloc, Suzanne Touzeau, Thierry Hoch, Henri H. Seegers, Christine Fourichon

► **To cite this version:**

Amandine Lurette, Catherine C. Belloc, Suzanne Touzeau, Thierry Hoch, Henri H. Seegers, et al.. Modelling Salmonella spread in a pig farm under three biosecurity strategies. International Symposium of Salmonella and Salmonellosis (I3S), May 2006, Saint-Malo, France. 2006. hal-02812895

HAL Id: hal-02812895

<https://hal.inrae.fr/hal-02812895v1>

Submitted on 7 Aug 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Modelling *Salmonella* spread in a pig farm under three biosecurity strategies

A. Lurette¹, C. Belloc¹, S. Touzeau², T. Hoch¹, H. Seegers¹, C. Fourichon¹

¹Unit of Animal Health Management, Veterinary School – INRA, BP 40706, 44307 Nantes Cedex 3, France.

²Unit of Applied Mathematics and Computer Science, INRA, 78352 Jouy-en-Josas Cedex, France.



INTRODUCTION

The production level is a critical point of the contamination of the whole pork supply chain by *Salmonella*.

Objectives : To estimate the number of shedder and carrier pigs at slaughter age under three biosecurity strategies.

MATERIAL & METHODS

1. Development of a **stochastic mathematical model** to simulate:

- the pig **population dynamics (Fig. 1)** in a farrow-to-finish herd with batch farrowing management;
- the **transmission of *Salmonella*** depending on the contamination of the pen floor (**Fig. 2**).




Figure 1. Simplified farrow-to-finish herd production system.
 → flows linked to demographic processes
 - - - → flows controlled by producers




Figure 2. Epidemiological model.

Contact structure influenced by:

- the variability in reproduction and in growth,
- the producers' management,
 → leading to variations in the transmission.

Groups of slaughtered pigs were issued from several batches depending on their weight.

Parameters: from published literature and expert opinion.

2. The three **biosecurity strategies** differ only for finishing pigs:

- AIAO** strict all-in/all-out management with complete cleaning, disinfection and drying period between batches;
- NDP** all-in/all-out allowing the suppression of the drying period;
- BM** management allowing the suppression of the drying period and batch mixing.

3. **Simulations** were run after the introduction of a shedder gilt in a AIAO herd. After equilibrium, the 3 strategies were implemented.

RESULTS

The number of batches in a group of slaughtered pigs was significantly higher under the BM strategy (**Tab. 1**).

Table 1. Proportion of groups of pigs at slaughter per number of batches under the three biosecurity strategies.

Biosecurity strategy	Number of batches			
	1	2	3	>3
AI/AO	0.18	0.34	0.30	0.18
NDP	0.21	0.35	0.27	0.17
BM	0.15	0.20	0.31	0.34

Prevalence of shedder and carrier pigs varied widely between batches, within and between strategies (**Fig. 3**).




Figure 3. Prevalence of shedders & carriers at slaughter age.

Mean prevalence differed significantly between strategies.

AIAO: 0.27 ± 0.04 , **NDP**: 0.45 ± 0.02 , **BM**: 0.53 ± 0.01 .

DISCUSSION & CONCLUSION

A less strict management increased dramatically the number of shedder and carrier pigs at slaughter age.

Further validation of the model based on field data is considered.

Both the heterogeneity of the contact structure and the biosecurity strategies play a major role in pathogen transmission in a farrow-to-finish herd.