



HAL
open science

Modelling as a tool to identify key measures to reduce Salmonella prevalence in slaughter pigs

Amandine Lurette, Suzanne Touzeau, Pauline Ezanno, Thierry Hoch,
Christine Fourichon, Henri H. Seegers, Catherine C. Belloc

► **To cite this version:**

Amandine Lurette, Suzanne Touzeau, Pauline Ezanno, Thierry Hoch, Christine Fourichon, et al..
Modelling as a tool to identify key measures to reduce Salmonella prevalence in slaughter pigs. 1.
International Conference on Infectious Disease Dynamics (Epidemics), Dec 2008, Asilomar, United
States. 2008. hal-02813018

HAL Id: hal-02813018

<https://hal.inrae.fr/hal-02813018>

Submitted on 7 Aug 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Modelling as a tool to identify key measures to reduce *Salmonella* prevalence in slaughter pigs

A. Lurette¹, S.Touzeau², P. Ezanno¹, T. Hoch¹, C. Fourichon¹, H. Seegers¹, C. Belloc¹

¹INRA, ENVN, UMR1300 Bio-agression Epidémiologie et Analyse de Risque, BP 40706, F-44307 Nantes, France,

²INRA, UR341 Mathématiques et Informatiques Appliquées, F-78350 Jouy-en-Josas, France




INTRODUCTION

The control of *Salmonella* in pigs is a major public health concern.

The objective of this study was to demonstrate how the modelling approach developed allows to determine better ways of applying control measures: animals targeted, periods of application and the measure of efficiency required; in order to significantly reduce the *Salmonella* prevalence in pigs.

MATERIALS & METHODS

A stochastic mathematical model representing both the pig and sow populations within a farrow-to-finish herd and the *Salmonella* spread was used (Lurette *et al.*, 2008) (Fig. 1).



Measures to test :

- Optimize the implementation of existing measures → acidified food Farzan *et al.*, 2006
- Identify new potential measures and their implementation → vaccination Roesler *et al.*, 2006

Representation in the model :

- Hypothesis on implied mechanisms → effect on three epidemiological parameters: infection probability (p_i), quantity shed (Q) and re-activation probability (β_r) → 3 reduction levels (25, 50 & 90%) from their default value.
- Selection of animals and duration → 2 types of measure
 - Measure 1) either on sows or pigs during their lifetime
 - Measure 2) on lactating sows, on post-weaning or on finishing pigs

RESULTS

Reduction %	Infection probability p_i	Quantity shed Q					
		25	50	90			
Animal Targeted	Reference	25	50	90	25	50	90
Measure 1)							
Sows	19.8	10.1	1.9	0.2	15.2	13.4	4.1
Pigs	19.8	16	13.1	7.7	19.5	16.8	10.8
Measure 2)							
Lactating sows	19.8	19.8	19.6	18.7	17.2	15.4	7.1
Finishing pigs	19.8	15.9	10.9	6.3	19.8	19.8	19.8


Table 1. Mean seroprevalence in groups of delivered pigs according to reduction in the probability of infection (p_i) or of the quantity shed (Q). Reference : default values for all parameters



Scenarios decreasing the mean seroprevalence differ according to the parameter considered, its level of reduction and the animal targeted (Tab. 1 & Fig. 2). A mean seroprevalence in groups of delivered pigs equal or less to 5% can be obtained by:

- Reducing the quantity of bacteria shed by the sows by 90%,
- Reducing both the probability of infection and the quantity shed by 25% and 50% respectively.

Moreover, when also combined with a reduction of the reactivation probability, the scenario can lead to a mean seroprevalence under 1%.



DISCUSSION

Preliminary results that determine the required effect of a control measure on infection parameters. Further experimental studies are needed to confirm such effects are attainable. Moreover, our model allows to assess the effects of combinations of control measures.

Farzan *et al.*, (2006) *Prev. Vet. Med.*, 73, 241-254
 Lurette *et al.*, (2008) *Vet. Res.*, 49, 39:01
 Roesler *et al.*, (2006) *J. Vet. Med. B.*, 53, 224-228