



How strict selection of the raw eddy-covariance data influences gpp estimation of a temperate beech forest

Bernard B. Longdoz, Patrick Gross, André A. Granier

► To cite this version:

Bernard B. Longdoz, Patrick Gross, André A. Granier. How strict selection of the raw eddy-covariance data influences gpp estimation of a temperate beech forest. 5. Réunion du projet Carbo Europe, Oct 2007, Poznan, Poland. 1 p., 2007. hal-02815801

HAL Id: hal-02815801

<https://hal.inrae.fr/hal-02815801>

Submitted on 6 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

HOW STRICT SELECTION OF THE RAW EDDY-COVARIANCE DATA INFLUENCES GPP ESTIMATION OF A TEMPERATE BEECH FOREST

LONGDOZ Bernard, GROSS Patrick, GRANIER André
UMR Ecologie et Ecophysiologie forestière, Centre INRA Nancy, 54280 Champenoux
longdoz@nancy.inra.fr

1. Abstract

The **data quality** and the **selection** of the correct Eddy Covariance (EC) records become an important step in the CO_2 flux determination procedure. An **innovative combination** of existing assessment **tests** is used to give a relatively complete evaluation of the **net ecosystem exchange** measurements. For the 2005 full-leaf season at the Hesse site, the percentage of **bad quality** data is relatively high (**59.6%**) especially during night-time (68.9%). This result strengthens the importance of the data gap filling method. The **filtering** used does **not lead** to a real **improvement** of the accuracy of the relationship between the CO_2 fluxes and the climatic factors. The **soil respiration spatial heterogeneity** (on a site with relatively homogenous vegetation pattern) seems to be too important to allow this improvement. However, the **data rejected** present some common characteristics. Their removal lead to a **10% increase** in the total amount of CO_2 respired (**Reco**) and photosynthesised (**GPP**) during the 2005 full-leaf season. Consequently the application of our combination of multiple quality tests is able improve the inter-annual analysis. The question of a systematic application on the large database like the CarboEurope and FLUXNET is legitimate.


2. Hesse site

- Hesse beech (90%) plot
- Age: 39 years
- Height: 17 m
- Precipitation: 950 mm yr⁻¹
- Mean annual temp. : 10.1°C
- Soil type : Luvisol/Stagnic luvisol
- Soil texture: Sable 7%; Limon 64%; Argile 29%



Fig. 1

Fig. 2



3. Material

EDDY-COVARIANCE SYSTEM

- Tower (Fig. 2)
- IRGA LI6262 at soil level (Fig. 3)
- Sonic anemometer Gill R2 (Fig. 4)
- Tubing from sonic to IRGA (30 m)



Fig. 3



Fig. 4

4. Method

Raw data tests (on high frequency [CO_2] and wind measurements)

- Anemometer or IRGA problems: spikes, discontinuities of mean or variance, unrealistic data, large skewness, kurtosis and standard deviation (Vickers & Mahrt, 1997)
- Mass Flow Controller (MFC) anomalies: constant airflow rate in the EC tubing

CO_2 EC fluxes (F_c , computed each half-hour) tests

- Flux stationarity during the half-hour (Foken & Wichura, 1996)
- Footprint, F_c flagged if 10% comes out of Hesse beech forest (Soegaard et al., 2003)
- U^* threshold to dismiss horizontal advection periods

5. Results

Quality test	% flagged
Anemometer problems	1.6
IRGA problems	33.3
MFC problem	0.1
Stationarity	23.9
Footprint	1.5
U^*	9.9
Total	59.6

Table 1

Quality test	% flagged
Spikes	0.1
Mean discontinuities	1.9
Variance discontinuities	2.1
Absolute limits	0.1
Skewness	14.5
Kurtosis	31.9
Standard Deviation	3.4
Total	33.3

Table 2

Table 1 : Percentages of half-hours flagged for the different tests
Table 2 : Percentages of half-hours flagged for the different IRGA problems

- Two tests appear to be highly **restrictive**: the CO_2 **IRGA kurtosis** anomalies and the **stationarity**
- The stationarity flag percentage is close to the one estimated by Rebmann et al. (2005) for the Hesse summer 2000 (28%).
- In Vickers and Mahrt (1997), the kurtosis test is also the more efficient among the IRGA problems ones (with % higher than here)
- The sum of all the % of the different tests exceeds the total value because of multi-flagged data.

- The data elimination changes the relationship between the CO_2 fluxes and the environmental factors
⇒ **Reco**, **GPP** and **NEE** for the 2005 full-leaf season vary all by about **10%**, with **more** important **photosynthesis** exchanges, **more** CO_2 produced by **respiration** processes and a **higher net sequestration** when **quality tests** are applied
- The selection of the non flagged data doesn't really improve the fit of Reco with soil t° function (R^2 stable) and water content function (R^2 gains less than 5%) except for the u^* test
⇒ **Unexplained temporal Reco variations** after u^* filtering don't come from problems by the tests but rather from **different footprint area combined with large soil respiration heterogeneity**