

First results on inducing quiescence in Trichogramma chilonis development

Elisabeth Tabone, Hong Do Thi Khanh, Etty Colombel, - Goebel R., - Roux E.

▶ To cite this version:

Elisabeth Tabone, Hong Do Thi Khanh, Etty Colombel, - Goebel R., - Roux E.. First results on inducing quiescence in Trichogramma chilonis development. 7 ISSCT Entomology Workshop, Apr 2009, San Miguel de Tucumán, Argentina. 1 p. hal-02815945

HAL Id: hal-02815945 https://hal.inrae.fr/hal-02815945

Submitted on 6 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

First Results on Inducing Quiescence in Trichogramma chilonis Development

Elisabeth Tabone¹, Hong Do Thi Khanh¹, Etty Colombel¹, Régis Goebel³, and Estelle Roux²

¹INRA Sophia Antipolis - Unité de lutte biologique, 1382 route de Biot, 06560 Valbonne, France (tabone@sophia.inra.fr)

²FDGDON La Réunion. - 23, rue Jules Thirel-Savannah, 97460 Saint-Paul, La Réunion ³CIRAD La Valette Agropolis, Unité de Recherches Systèmes canniers, Avenue Agropolis 34398 Montpellier, France

Since 2000, a collaborative project between INRA, CIRAD and FDGDON has successfully developed a program for biocontrol of the sugarcane stem borer using Trichogramma chilonis on Reunion Island. Previous research has demonstrated interest in biological control as a control strategy in sugarcane. However, to implement this strategy on a large scale, some important technical problems needed addressing, mainly how to lower the overall costs associated with the program. To develop a cold storage technology will bring us a step closer to reaching that goal. This objective corresponds to the current demand by the industry community involved in the promotion and trading of agricultural products. Since 2008 a new research program has been initiated to study the induction of diapause or quiescence in T. chilonis development. The resulting knowledge is expected to greatly improve both mass rearing and releasing labour. This project is supported for three years by the French Ministry of Agriculture (CASDAR fund) (from mid 2008 to mid 2011). Optimal conditions (temperature, photoperiod and stage) to induce and maintain diapause or quiescence are being studied on the strain of T. chilonis currently used. The following parameters will be recorded: emergence rate, fecundity, sex-ratio and development time. A range of temperature lower than 18°C will be tested. The effects of storage at low temperature on the physiology of T. chilonis will be monitored. Initial results have already defined biotic and abiotic parameters required for successful storage for up to 2 months and these findings allow us to be optimistic for the future.