



HAL
open science

Les territoires d'exploitations agricoles qui pratiquent des systèmes de production différents présentent-ils une biodiversité différente? L'exemple des oiseaux nicheurs

Sébastien Bonthoux, Jean-Philippe Choisis, Annick A. Gibon, Sylvie Ladet,
Gerard Balent

► To cite this version:

Sébastien Bonthoux, Jean-Philippe Choisis, Annick A. Gibon, Sylvie Ladet, Gerard Balent. Les territoires d'exploitations agricoles qui pratiquent des systèmes de production différents présentent-ils une biodiversité différente? L'exemple des oiseaux nicheurs. 3. Journée Recherche 2010 de l'ENSAT, Jun 2010, Auzeville Tolosan, France. 18p. hal-02816458

HAL Id: hal-02816458

<https://hal.inrae.fr/hal-02816458>

Submitted on 6 Jun 2020


HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Les territoires d'exploitations agricoles qui pratiquent des systèmes de production différents présentent-ils une biodiversité différente ?

L'exemple des oiseaux nicheurs

BONTHOUX S., CHOISIS JP., GIBON A., LADET S., BALENT G.



Ce travail est financé par une bourse ANR, BIODIVAGRIM (ANR-07-BDIV-002) "La conservation de la biodiversité dans les agro-écosystèmes : une approche de modélisation des paysages spatialement explicite".

Aim of the study

Agri-environmental policies are most often applied at the individual farm level. An improved effectiveness of these policies could result from an it is then necessary to study the ecological patterns by considering the level of agricultural organisation.

However, many studies are currently lead at the field scale or the landscape scale without considering the whole farm territory.

Our study focus on the relationships between bird diversity in farmland and spatial patterns of farming-system types regarded as a level of agricultural organisation.


We test **2 hypotheses**:

- The farming systems can discriminate bird diversity.
- Landscape areas with extensive farming systems have an higher bird diversity than the ones with intensive farming systems because they include more semi-natural elements.

Our study area



11 km




'Vallées et Coteaux de Gascogne'
Long Term Ecological Research site




- A hilly region
- A combination of arable crops, (maize, oilseed rape, sorghum, sunflower, forage), meadows, grazed pastures, and small woodlands

Method **Global approach**



Ecological data
A grid of point counts to record birds



Agronomical data
34 interviewed farms



A typology of farming systems
in 5 modalities



Results


Bonthoux S. et al.

Method Global approach

Ecological data


A grid of point counts to record birds

1




2


Agronomical data
34 interviewed farms



A typology of farming systems
in 5 modalities



3



Results


Ecological data

A grid of point counts on the whole study area:

A systematic sample

One point count = 20 minutes

All birds seen or heard are recorded




Agronomical data

34 interviewed farms

Farming system data

- history of farm
- farm productions
- technical operations
- economical data



Spatial arrangement of farm territory (computed by GIS)

- field size, field shape, field slope
- farm territory fragmentation


Co-inertia analysis

A typology in 5 types of farming system


Adapted from Choisis et al. 2010

Bonthoux S. et al.

A typology in 5 types of farming system



Production intensity





Some characteristics of the farm types:

Production intensity

	Less favored extensive livestock	Extensive livestock system in a cessation process	Mixed crop-livestock system	Intensive livestock system	Crop system
Farm number	4	5	15	6	4
Averaged Agricultural area (ha)	82	23	61	155	69
Averaged Field size (ha)	1.8	1.8	2.5	3.9	4.8
% of sloping fields (>15%)	58	26	20	25	6

Land uses



→ semi-natural elements

Some pictures of the farming systems



An open landscape with arable crops




A landscape with sloping fields and permanent pastures



An intermediate landscape with a combination of crops and pastures

Linking farmland bird diversity to farming system



→ Each point count is related to one farming system

- 5 GLMs to link the bird species richness to the types of farming system

Statistical unit = point count, n=363

5 Response variables

Species Richness

of

All bird species
Nesting on ground species
Nesting in trees species
Omnivorous species
Insectivorous species

Explanatory variable

Type of farming system (5 modalities)

43 farmland species (passerines) were recorded in the 363 point counts.

Occurrence of the 10 most frequent species

English name	Latin name	Occurrence
Eurasian Blackcap	<i>Saxicola atricapilla</i>	234
Common Blackbird	<i>Turdus merula</i>	209
Common Nightingale	<i>Luscinia megarhynchos</i>	176
Corn Bunting	<i>Emberiza calendra</i>	156
Common Whitthroat	<i>Sylvia communis</i>	132
Great Tit	<i>Parus major</i>	128
Blue Tit	<i>Parus caeruleus</i>	119
Common Chiffchaff	<i>Phylloscopus collybita</i>	110
Skylark	<i>Alauda arvensis</i>	101
Red Robin	<i>Erithacus rubecula</i>	87

Two species of open habitat



Skylark



Corn Bunting

Three species of intermediate habitat



Eurasian Blackcap



Common Nightingale

Common Blackbird



Linking farmland bird diversity to farming system

Farming systems

Production intensity →

		Less favored extensive livestock	Extensive livestock system	Mixed crop-livestock system	Intensive livestock system	Crop system
Species richness	Total	ns	ns	ns	ns	-
	Avged SR / point	7.6	7.0	7.3	7.5	6.3
	Nesting ground	+	ns	ns	ns	-
	Nesting in tree	ns	ns	ns	ns	ns
	Omnivorous	+	ns	ns	ns	-
	Insectivorous	ns	ns	ns	ns	ns

+ = positive effect **-** = negative effect **ns** = non significant

Discussion

Our results show that bird diversity differs among farming systems which are extremely opposed along a gradient of production intensity.

The crop system which has few semi-natural elements on its territory, has the less bird diversity. In contrast, the less favored extensive livestock system has the higher diversity.

However, the difference of bird diversity is very weak between these farming systems. This is probably because the studied landscape is globally heterogeneous. The different farming systems are spatially mixed. Then, birds which are very mobile move between the different farming systems.

Results would be probably more contrasted if we would consider these farming systems in different homogeneous landscapes.

Perspectives

Mettre en lien la **dynamique** du territoire des exploitations agricoles avec la dynamique des communautés d'oiseaux



1992



2006

Remembrement sur la
commune d'Eoux



Merci pour votre attention !