



HAL
open science

Reducing conditions varied with diets and bacterial communities in the rumen on dairy cows

Christine Julien, Jean-Philippe Marden, E. Auclair, Laurent L. Cauquil, -
Moncoulon, Corine Bayourthe

► To cite this version:

Christine Julien, Jean-Philippe Marden, E. Auclair, Laurent L. Cauquil, - Moncoulon, et al.. Reducing conditions varied with diets and bacterial communities in the rumen on dairy cows. Gut Microbiology, 7. Joint Symposium, Jun 2010, Aberdeen, United Kingdom. pp.1. hal-02817627

HAL Id: hal-02817627

<https://hal.inrae.fr/hal-02817627v1>

Submitted on 6 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Reducing conditions varied with diets and bacterial communities in the rumen of dairy cows. By C. JULIEN¹, J.P. MARDEN², E. AUCLAIR², L. CAUQUIL¹, R. MONCOULON¹ AND C. BAYOURTHE¹, ¹*Université de Toulouse, INRA, UMR 1289 INRA/INPT/ENVT TANDEM, 31326 Castanet-Tolosan, France,* ²*Lesaffre Feed Additives, 59520 Marquette-Lez-Lille, France.*

This study aimed at characterizing both the change in ruminal bacterial population and physico-chemical conditions of four dairy cows fed two maize silage (67 %) and soybean meal (17%)–based diets completed with either 16 % of ground wheat (WD) or 16% of crushed maize (MD), on a DM basis.

Ruminal pH and redox potential (E_h) were recorded (Marden et al., 2005) and the Clark's Exponent (rH) was calculated. Ruminal content was sampled for bacterial profiling by means of Capillary Electrophoresis Single-Strand Conformation Polymorphism (CE-SSCP).

Average ruminal pH was significantly lower with WD (6.12) than MD (6.25). Ruminal E_h and rH were also lower with WD than MD: – 186 mV, 6.04 and – 152 mV, 7.43, respectively. Results suggested that dietary composition influenced ruminal reducing conditions. CE-SSCP profiles showed that the bacterial communities varied with diet composition (R-ANOSIM = 0.173, $P = 0.051$): there was 20 % of difference between the average profiles of the 2 diets. Furthermore, there was a correlation between CE-SSCP profiles and both pH and rH ($P \leq 0.05$, $r^2 = 0.19$ and $r^2=0.13$, respectively). It confirmed the well-known link between bacterial communities and ruminal pH and also put forward a relationship between bacterial communities and ruminal reducing conditions. It seemed that ruminal rH and E_h were specifically correlated to the richness of bacterial communities ($P \leq 0.05$, $r^2 = 0.25$ and $r^2=0.25$, respectively).

In conclusion, this study addresses the possibility to modulate ruminal E_h via feeding in ruminants to stimulate the adequate microflora for better use of dietary material.

Marden, J. P., C. Bayourthe, F. Enjalbert, and R. Moncoulon. 2005. A new device for measuring kinetics of ruminal pH and redox potential in dairy cows. *J. Dairy Sci.* 88:277-281.