



# Transcriptomic analysis of intestinal genes following acquisition of pea enation mosaic virus by the pea aphid *Acyrthosiphon pisum*

Catherine Reinbold, Sylvie Tanguy, Denis Tagu, Véronique Brault, . Société Entomologique de France, . Université Lyon 1, . Institut National Des Sciences Appliquées

## ► To cite this version:

Catherine Reinbold, Sylvie Tanguy, Denis Tagu, Véronique Brault, . Société Entomologique de France, et al.. Transcriptomic analysis of intestinal genes following acquisition of pea enation mosaic virus by the pea aphid *Acyrthosiphon pisum*. 16. Colloque de Biologie de l'Insecte, Oct 2010, France. 1 p. hal-02818371

HAL Id: hal-02818371

<https://hal.inrae.fr/hal-02818371>

Submitted on 6 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## P.34 Transcriptomic analysis of intestinal genes following acquisition of pea enation mosaic virus by the pea aphid *Acyrthosiphon pisum*

C. Reinbold<sup>1</sup>, S. Tanguy<sup>2</sup>, D. Tagu<sup>2</sup>, V. Brault<sup>1</sup>

catherine.reinbold@colmar.inra.fr

<sup>1</sup> INRA, UMR 1131 SVQV, 28 rue de Herrlisheim, F-68021 COLMAR – <sup>2</sup> INRA, UMR 1099 BiO3P, Domaine de la Motte, F-35653 LE RHEU

Viruses in the *Luteoviridae* family are strictly transmitted by aphids in a non-propagative circulative and persistent mode. Virions ingested by aphids, successively cross the gut and the accessory salivary gland epithelia before being released, together with saliva, in the plant vasculature. Virions transport through aphid cells occurs by a transcytosis mechanism. We conducted a transcriptomic analysis of intestinal genes of the pea aphid *Acyrthosiphon pisum* following uptake of *Pea enation mosaic virus-1* (a member of the *Luteoviridae* family). Among the 7166 transcripts analysed, 128 were significantly regulated (105 genes down-regulated and 23 up-regulated). 5% of the identified genes were involved in intracellular trafficking, endocytosis and signal transduction, three important steps in the internalization and transport of virions. The limited levels of down- (maximum of 3.45 fold) and up-regulation (maximum of 1.37 fold) suggest that the virus hijacks a constitutive endocytosis-exocytosis mechanism without heavily perturbing cell metabolism. Although limited to about 20% of the pea aphid genes, this work represents the first large scale analysis of aphid gene regulation following virus acquisition.

- Brault et al. (2010) *J. Gen. Virol.* 91, 802-808