HAL
open science

Intercroppping, an application of ecological principles, to increase yield and durum wheat grain protein in low nitrogen input systems

Laurent Bedoussac, Eric E. Justes

To cite this version:

Laurent Bedoussac, Eric E. Justes. Intercroppping, an application of ecological principles, to increase yield and durum wheat grain protein in low nitrogen input systems. Farming systems design. Methodologies for integrated analysis of farm production systems, Aug 2009, Monterey, United States. , 1p., 2009. hal-02819302

HAL Id: hal-02819302

https://hal.inrae.fr/hal-02819302v1

Submitted on 20 Jun 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

INTERCROPPING, AN APPLICATION OF ECOLOGICAL PRINCIPLES TO INCREASE YIELD AND DURUM WHEAT GRAIN PROTEIN IN LOW NITROGEN INPUT SYSTEMS

L. BEDOUSSAC and E. JUSTES - INRA, UMR 1248 AGIR, F-31326 Castanet-Tolosan, France
E-mail: Laurent.Bedoussac@toulouse.inra.fr – Eric.Justes@toulouse.inra.fr

Farming Systems Design 2009
Methodologies for Integrated Analysis of Farm Production Systems
August 23-26 2009 - Monterey - USA

OBJECTIVES

- **Nitrogen acquisition** is often a major concern, particularly in low input systems where mineral **N is a limited resource**.
- Intercropping (IC) can improve the use of environmental resources (light, nutrients and water) resulting **in yield and quality advantages compared to sole cropping** (SC) (e.g. Willey, 1979).
- No reference on winter crops IC was available, despite winter sole crops seems more adapted to Southern Europe conditions.
- *Aim of our study:* **Propose innovative Durum wheat – Winter pea intercropping managements to optimize the use of available resources**
  - ***i)*** **Understanding competition between durum wheat and winter pea for different wheat cultivars**;
  - ***ii)*** **Analyzing the consequences of N availability on the performance of IC (grain protein, yield and species proportion)**.

CONCLUSIONS

- The '**Durum wheat - Winter pea intercropping**' seems well adapted to the Southern France conditions because it allows:
  - ***i)*** **A better use of N resources (and light) during winter and early spring due to the complementarities of the 2 species**
  - ***ii)*** **A higher grain protein concentration of durum wheat at harvest due to:**
    - *a) High pea N_2 fixation rate in IC making available for the IC wheat almost as much soil mineral N per square meter as in the SC;*
    - *b) Fewer wheat ears, grains and yield per area in IC compared to SC due to interspecific competitions (Bedoussac and Justes 2009).*
- IC advantages were greater for the unfertilized or late N-fertilized treatment **confirming the interest of intercropping in low-input farming**.
- Optimal choices in N supply and wheat cultivar depend on the target of the intercrop. Two directions are possible:
  - ***i)*** **Increasing N availability and/or choosing a tall wheat cultivar that could increase wheat proportion**;
  - ***ii)*** **Reducing N supply and/or choosing a short wheat cultivar that could increase wheat grain protein and pea proportion.**

MATERIAL AND METHODS

- An experiment was carried out in Auzeville (SW France) in 2006-2007 on a clay loamy soil. The two species were sown in the 9th of November 2006 in **row-intercropping**. The experiment was based on a split-split-plot design with 3 replicates.
- **Three main treatments were compared:**
  - ***i)*** **W-SC:** Durum wheat (sown at 336 seeds/m²);
  - ***ii)*** **P-SC:** Winter pea (cv. Lucy sown at 72 seeds/m²);
  - ***iii)*** **IC:** Durum wheat-winter pea IC, **each specie sown at half of SC density**.

- **Four wheat cultivars of different height:** ***i)*** **Ac:** Acalou (89 cm); ***ii)*** **Nf:** Nefer (96 cm) ; ***iii)*** **Nd:** Neodur (98 cm) and ***iv)*** **Oj:** Orjaune (116 cm).
- **Four fertilizer-N sub-treatments (Pea SC only in N0) :** ***i)*** **N0:** No fertilizer ; ***ii)*** **N60:** 60 kg N.ha^{-1} (at **FLV** 'flag leaf visible' to increase **wheat grain protein content**) ; ***iii)*** **N80:** 80 kg N.ha^{-1} (at 'ear 1cm' to increase **wheat yield**) and ***iv)*** **N140:** corresponding to N80 and N60.
- **Measurements :** ***i)*** Wheat grain protein content (**GPC**); ***ii)*** Grain yield; ***iii)*** Land Equivalent Ratio (**LER,** see Fig.1 for formula) (Willey 1979) and ***iv)*** Percentage of N derived from N_2 fixation (**%Ndfa**) of pea calculated using the ^{15}N natural abundance method (Amarger et al. 1979).

RESULTS

Wheat Grain Protein in IC vs. SC



- **GPC** was 12% **higher in IC** than in SC
- The lowest the **GPC in SC** the highest the increase in **IC**
- IC reduced the gap in cultivars **GPC**
- **Ac** and **Nd** have greater GPC in SC & IC

Land Equivalent Ratio



- **LER** greater than 1 for N0 and N60

→ **IC more efficient than SC in N0 and N60**

- **Wheat LERp** always higher than 0,5

→ **Wheat took more advantage of N than pea**

- **Pea LERp** strongly reduced with N supply
- **Nd** & **Oj** Wheat LERp greater than **Nf** & **Ac**
- **Pea LERp** lower for **Nd** & **Oj** than **Nf** & **Ac**

→ **Pea yield is more reduced with tall cultivars**

→ **Complementary use of N resources**

Plant N derived from air (%Ndfa) and N derived from soil (QNdfs)

		%Ndfa (% N uptake)	QNdfs (kg N ha^{-1})
SC	N0	52 ± 4	83
IC	N0	84 ± 5	13
	N60	85 ± 7	7
	N80	60 ± 9	19
	N140	70 ± 9	12

- **Pea %Ndfa higher in IC** than in SC
- **Pea %Ndfa** reduced with N80 & N140
- **Pea QNdfs** only 14kg N ha^{-1} in IC

→ **Same N available for wheat in IC&SC**

- Amarger, N., A. Mariotti, F. Mariotti, J. Durr, C. Bourguignon, and B. Lagacherie. 1979. Estimate of symbiotically fixed nitrogen in field grown soybeans using variations in 15N natural abundance. Plant Soil 52:269-280.
- Bedoussac, L. and E. Justes. 2009. The efficiency of a durum wheat-winter pea intercrop to improve yield and wheat grain protein concentration depends on N availability during early growth. Plant Soil DOI: 10.1007/s11104-009-0082-2
- Willey, R. 1979. Intercropping - its importance and research needs. 1. Competition and yield advantages. Field Crop Abstr. 32:1-10.