HAL open science

Volatile soil carbon: joint project PROCOPE

Bernard B. Longdoz, Martin Maier, Helmer Schack-Kirchner, Caroline C. Plain

To cite this version:

Bernard B. Longdoz, Martin Maier, Helmer Schack-Kirchner, Caroline C. Plain. Volatile soil carbon: joint project PROCOPE. Réunion dans le cadre du partenariat Hubert Curien "PROCOPE", Feb 2009, Freiburg, Germany. 22 p. hal-02820864

HAL Id: hal-02820864

https://hal.inrae.fr/hal-02820864v1

Submitted on 6 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

VOLATILE SOIL CARBON: JOINT PROJECT PROCOPE

Bernard Longdoz, Martin Maier, Helmer Schack-Kirchner, Caroline Plain

Bilateral project between :

Inst. Fur Bodenkunde Und Waldernahrungslehre

&

UMR Forest Ecology and Ecophysiology

+

Nancy-Université
Université Henri Poincaré

ALBERT-LUDWIGS-UNIVERSITÄT FREIBURG

INRA

UMR FOREST ECOLOGY and ECOPHYSIOLOGY

(Director A. Granier)

Plant and ecosystem functioning under environmental changes (50 permanents, 75 people)

Forest Phyto-ecology
(J.-L. Dupouey)

Tree and Ecosystem Functioning
(D. Epron)

Physiology
(E. Dreyer)

Technical support (Isotope, microscope, mineral analysis (C. Brechet)

GIS (D. Maurice)

Nancy-Université
Université Henri Poincaré

ALBERT-LUDWIGS-UNIVERSITÄT FREIBURG

INRA

UMR INRA-UHP 1137
Ecologie et Ecophysiologie Forestières

Team : Tree and Ecosystem Functioning

- Flux and budget of carbon, water and nitrogen in interaction with climate and edaphic conditions
  - Ecosystem carbon cycle :
    - ✓ Ability of the forest to store the atmospheric carbon :

      Climatic changes

      Forest management (harvest)
  - Tree carbon and nitrogen reserves
  - Carbon and nitrogen allocation in the tree
  - Hydraulic tree and canopy functioning

Experimental Site : Hesse



Location : 48°40'N, 7°05'E

65 km from Nancy (East)
10 km from Sarrebourg (South)

Mean annual air Temp. : 10°C

Mean annual Precip. : 950 mm

Climate : Temperate

Experimental Site : Hesse



Experimental Site : Hesse

Equipped since 1996

GIP ECOFOR

Main instrumentation : Eddy covariance system

Experimental Site : Hesse

Eddy Covariance system

Sonic anemometer

3 wind speed components measurements at high ν (>10 Hz)

IRGA

[CO_2] and [H_2O] at high ν (>10 Hz)

Tower



Air transport

Data record and storage system

Net Fluxes : CO_2 (NEE), Evapo-transp.

Sensible and Latent heat

every 30 min

Experimental Site : Hesse



NEE

GPP (Photosynthesis)

TER (Total ecosystem respiration)

Estimation of TER during daytime by extrapolation of night EC data

GPP = NEE –TER

Annual

carbon sequestration (NEE)

GPP

TER



Experimental Site : Hesse

Automatic :

- Eddy covariance (Net CO_2 Ecosystem Exchange)
- Climate (T°, radiation, humidity, precipitations)
- Edaphic conditions (soil T° and water content)
- Tree diameter (dendrometer) ⇨ C biomass

Campaigns :

- LAI
- Soil respiration (Rs)
- Aerial biomass (trunks, leaves,…)
- Below ground biomass (roots)
- Soil composition,…

Temporal variability

Partitioning

Nancy-Université
Université Henri Poincaré

ALBERT-LUDWIGS-UNIVERSITÄT FREIBURG

INRA

UMR INRA-UHP 1137
Ecologie et Ecophysiologie Forestières

Why Soil Respiration ?

- Partitioning NEE between components fluxes

  (Rs second flux, Rs = 60-70% of TER)
- Explain temporal variability of TER
- Explain spatial variability of TER

Why Volatile Carbon Soil ?

- Solve eddy covariance measurement problems
- Improve soil carbon modelling
- Explain stable isotope fluctuations of soil CO_2 efflux

Eddy Covariance Problems

When low turbulence = quite nights

➔ EC fluxes ≠ Total ecosystem respiration (30% of the nighttimes)

(Longdoz et al. 2008)



Eddy Covariance Problems

Correction CO_2 stored in the canopy air and in the soil

Measurements of CO2 profile in canopy air below EC system

➔ We need $[CO_2]_{soil}$ measurements for soil CO_2 storage estimation



Modelling (carbon, water, energy)

(Longdoz et al. 2004)

Simulate impact of

- Climate changes (extreme events)
- Management
- Differences between species behaviour

on carbon sequestration

Two « black boxes » : Soil carbon budget & photosyntates allocation

Soil carbon model :

- “Century” soil carbon pools
- Site specific soil CO_2 efflux function Fc (t°, SWC)

➔Multi-layers model :

- CO_2 production profile depending on layer features

Modelling

➔Multi-layers model :

CO_2 production profile depending on layer features

➔ Heterotrophic – Autotrophic partitioning



➔ **We need $[CO_2]_{soil}$ measurements in addition to efflux**

Nancy-Université Université Henri Poincaré

ALBERT-LUDWIGS-UNIVERSITÄT FREIBURG

INRA

UMR INRA-UHP 1137 Ecologie et Ecophysiologie Forestières

Temporal variability of isotopic signature of soil efflux

- Carbon has two main isotopes (^{12}C and ^{13}C)
- Isotopic composition $\delta^{13}C$ (‰)

$$\delta^{13}C = \frac{R_{13/12}^{mes}}{R_{13/12}^{st}} - 1$$

$\delta^{13}C_{atm} \simeq -8$ ‰

$\delta^{13}C_{SOM} \simeq -21 \rightarrow -30$ ‰

- Discrimination (modification of the $\delta^{13}C$ value) during the biophysical processes (photosynthesis, CO_2 diffusion)

Nancy-Université Université Henri Poincaré

ALBERT-LUDWIGS-UNIVERSITÄT FREIBURG

INRA

UMR INRA-UHP 1137 Ecologie et Ecophysiologie Forestières

Measurements of $\delta^{13}C$ soil respiration

- ➔ Contribution of the heterotrophic respiration sources (litter/old SOM)
- ➔ Contribution of the plant organs (soluble sugar and amino acids, storage carbohydrate and proteins) in autotrophic soil respiration
- ➔ Separation of NEE between GPP-Reco

$$NEE = GPP + Réco$$

$$\delta^{13}C_{NEE} \cdot NEE = \underbrace{\delta^{13}C_{atm}}_{(\ -8‰)} \cdot GPP + \underbrace{\delta^{13}C_{Réco}}_{(\ -26‰)} \cdot Réco$$

- ➔ Input for inverse carbon cycle model

Determination of terrestrial & oceanic fluxes from $[CO_2]_{atm}$ and δ_{atm}

$\delta = -26‰$ $\delta = -12‰$

Nancy-Université Université Henri Poincaré

ALBERT-LUDWIGS-UNIVERSITÄT FREIBURG

INRA

UMR INRA-UHP 1137 Ecologie et Ecophysiologie Forestières

Up to now low frequency measurements (Keeling plots)

but temporal variability of $\delta^{13}C_{Rs}$!!!!

➔ Daily and seasonal $\delta^{13}C_{Rs}$ variations ?

➔ Which factors control these variations ?

High frequency measurements of fluxes of $^{13}CO_2$ and $^{12}CO_2$ with a Tunable Diode Laser Spectrophotometers (TDLS) as a promising tool for these purposes

Tunable Diode Laser Spectrophotometer



UMR INRA-UHP 1137
Ecologie et Ecophysiologie Forestières

Nancy-Université
Université Henri Poincaré

ALBERT-LUDWIGS-UNIVERSITÄT FREIBURG

INRA

UMR INRA-UHP 1137
Ecologie et Ecophysiologie Forestières

Chamber description



Seasonal $\delta^{13}C_{Rs}$ variations



Daily $\delta^{13}C_{Rs}$ variations



!!! Same range of fluctuation (up to 2‰) !!!

➔ Too large variations to avoid them

➔ What causes these variations ? (toward prediction of them)

Biological process during:

- ✓ Photosynthesis
- ✓ Carbon Transport
- ✓ CO2 respiration (Production)

Physical process during:

- ✓ CO_2 diffusion through the soil (from production point to the surface)

To separate the impact of physical from biological processes

➔ **We need measurements of $[CO_2]_{soil}$ and its $\delta^{13}CO_2$**

Nancy-Université
Université Henri Poincaré

ALBERT-LUDWIGS-UNIVERSITÄT FREIBURG

INRA

UMR INRA-UHP 1137
Ecologie et Ecophysiologie Forestières

Isotope workpackage of the project

Two/one time(s) per month

➞ air sampling (≠ depths)➡IRMS analyses for $^{13}CO_2$ in soil

➞ Keeling plot ➡ $^{13}CO_2$ of soil CO2 efflux