



HAL
open science

Hen oocyte quality and fertility

Sébastien Elis, Marina Govoroun, Joëlle Dupont, Philippe Monget, Isabelle Couty, Elisabeth Blesbois, Florence Batellier

► **To cite this version:**

Sébastien Elis, Marina Govoroun, Joëlle Dupont, Philippe Monget, Isabelle Couty, et al.. Hen oocyte quality and fertility. International Chick Meeting, Apr 2007, Barcelone, Spain. 1 p., 2007. hal-02821070

HAL Id: hal-02821070

<https://hal.inrae.fr/hal-02821070>

Submitted on 6 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Hen oocyte quality and fertility


S. Elis, M. Govoroun, J. Dupont, P. Monget, I. Couty, E. Blesbois, F. Batellier
INRA, PRC - 37380 Nouzilly

Introduction

The reproductive physiology of the hen shows numerous specificities that include oviparity and telolecithe oocyte. Moreover, the intensive selection on conformation parameters leads to a dramatic decrease in reproductive ability. This leads to investigate the regulation of the expression and function of genes involved in the maturation of the oocyte, with consequences on the development of the early embryo. This last point is very important in hen because of the late embryonic genome reactivation. The aim of the present study was to explore the expression of two genes of the TGF- β family suggested to be implicated in fertility, *gdf9* and *bmp15*. Biological effects of BMP15 were also investigated.


Materials and Methods

mRNA was extracted from Granulosa Cells, Germinal Disc Region and embryo with tri-reagent for quantitative RT PCR on candidate genes.





Tissular specificity and pattern of expression were explored during follicular maturation and early embryo development. *In situ* hybridization was performed on hen ovary to localize these genes. Effects of one of these genes, BMP15, were investigated on granulosa cells of F3/4, F2 and F1 follicles

Results



Preferentially expressed in the ovary



mRNA localised in the oocyte Both of these genes are expressed in the oocyte, and their expression decrease after the ovulation

BMP15 inhibits progesterone LH- or FSH-induced secretion

bmp15 and *gdf9* are expressed in the hen oocyte
Their expression decreased after the ovulation, even after the embryonic genome activation. They should be used during the follicular maturation or early embryo development
BMP15 inhibits progesterone induced secretion

BMP15 may have a key role in the female fertility in birds