



**HAL**  
open science

## Water and energy fluxes above an Eucalyptus plantation in Brazil: environmental control and comparison with two eucalypt plantations in Congo

Yann Nouvellon, Jose Luiz Stape, Jean-Paul Laclau, Jean-Marc J.-M. Bonnefond, Humberto da Rocha, Otávio Camargo Campoe, Claire Marsden, Jean-Pierre Bouillet, Rodolfo Araujo Loos, Antoine Kinana, et al.

### ► To cite this version:

Yann Nouvellon, Jose Luiz Stape, Jean-Paul Laclau, Jean-Marc J.-M. Bonnefond, Humberto da Rocha, et al.. Water and energy fluxes above an Eucalyptus plantation in Brazil: environmental control and comparison with two eucalypt plantations in Congo. IUFRO Workshop Canopy processes in a changing climate, Oct 2010, Hobart, Australia. 1 p. hal-02821072

**HAL Id: hal-02821072**

**<https://hal.inrae.fr/hal-02821072>**

Submitted on 6 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Water and energy fluxes above an Eucalyptus plantation in Brazil: environmental control and comparison with two eucalypt plantations in Congo

Yann Nouvellon<sup>1,2,3</sup>, José Luiz Stape<sup>3</sup>, Jean-Paul Laclau<sup>1,4</sup>, Jean-Marc Bonnefond<sup>5</sup>, Humberto da Rocha<sup>2</sup>, Otavio Camargo Campoe<sup>4</sup>, Claire Marsden<sup>1</sup>, Jean-Pierre Bouillet<sup>1,4</sup>, Rodolfo Araujo Loos<sup>6</sup>, Antoine Kinana<sup>7</sup>, Gueric le Maire<sup>1</sup>, Laurent Saint-André<sup>1</sup>, Olivier Roupsard<sup>1</sup>

1 CIRAD, Persyst, UPR80, 34398 Montpellier Cedex 5, France; Email: [yann.nouvelon@cirad.fr](mailto:yann.nouvelon@cirad.fr)

2 Departamento de Ciências Atmosféricas/IAG/Universidade de São Paulo, São Paulo, Brasil

3 North Carolina State University, Raleigh, 27695, NC-USA, [jlstape@ncsu.edu](mailto:jlstape@ncsu.edu)

4 USP, ESALQ, Av. Pádua Dias, 11, Piracicaba, SP, 13418-900, Brazil

5 INRA, UR1263 EPHYSE, F-33140, Villenave d'Ornon, France

6 Fibria, Aracruz, Esperitu Santo, Brazil

7 CRDPI, BP 1291, Pointe-Noire, Republic of Congo

Tropical eucalypt plantations provide an increasing share of the global wood supply. High yields on weathered tropical soils are one of the main reasons for their success. However, there are often suspected of high water-use and their productivity may often be constrained by water availability. Our main objectives in this study were: 1) to quantify water-use (actual evapotranspiration, AET) by a 5-6 yr-old eucalypt plantation in Southeast Brazil; 2) to investigate the main environmental factors controlling canopy stomatal conductance and the partitioning of available energy between latent (LE) and sensible heat (H) fluxes; 3) to compare the AET measured in this highly productive plantation to the AET measured in two eucalypt plantations in Congo and identify the main factors that explained the observed differences in water-use. AET measured by eddy-covariance in the Brazilian plantation ( $\approx 1300$  mm/yr) represented nearly all of the annual rainfall ( $\approx 1380$  mm/yr). Most of the available energy (82%) was used for AET. The bowen ratio (H/LE) remained very low most of the year, but increased sharply when the “effective” relative extractable water content (REW) in the root zone (0-10 m) decreased below a threshold value. The diurnal and seasonal variability in canopy stomatal conductance was accurately explained by a simple model accounting for the effects of VPD, REW, LAI, and incoming PAR. Despite similar annual rainfall ( $\approx 1200$  mm/yr), AET measured in the less productive Eucalypt plantations in Congo was low ( $\approx 700$  mm/yr) compared to that measured in Brazil. These differences were mostly explained by differences in net radiation ( $\approx 2740$  MJ/m<sup>2</sup>/yr in Congo versus  $\approx 3830$  in Brazil), site fertility and LAI (1.4 and 2.4 for the two clones in Congo versus 3.2 in Brazil). In Brazil, AET decreased sharply after clear-cutting and replanting allowing deep drainage (below 10 m) a few months after clear-cutting.

## Key Words

Eucalypt plantation, Actual evapotranspiration, eddy-covariance, canopy stomatal conductance