



## QSAR/QSPR approach of aroma compounds behaviour in food products

Anne Tromelin, Samuel Lubbers, Daniel Martinez, Elisabeth Guichard

### ► To cite this version:

Anne Tromelin, Samuel Lubbers, Daniel Martinez, Elisabeth Guichard. QSAR/QSPR approach of aroma compounds behaviour in food products. 14. European Conference on Food Chemistry (Euro Food Chem XIV), Aug 2007, Paris, France. 2007. hal-02821418

**HAL Id: hal-02821418**

**<https://hal.inrae.fr/hal-02821418v1>**

Submitted on 6 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Introduction

- ❖ Improved standard of living and lifestyle → demand for food products with less or no fat, sugars
- Equilibrium flavour retention/flavour release depends:
  - on the components of the food matrix
  - on the physicochemical properties of the flavour compound
- Reducing the fat content of a food system → modifications in flavour release and perception
- To retain the properties of a product when the fat is removed, fat-substitutes are used (starch, pectin...)

Influence of texturing agents is unclear

Use QSPR approach to evaluate the influence of texturing agent and the chemical structure of aroma compounds on the partition coefficient of aroma compounds between food product and vapour phase

Quantitative Structure-Activities Relationships (QSAR)  
Quantitative Structure-Properties Relationships (QSPR)

assumption:

changes in molecular properties elicit different responses

dependant variable:  
"activity" or "property":  
Experimental Response

QSAR

QSPR

links

independent variable:  
molecular features  $p_1, p_2, \dots, p_n$   
Molecular Descriptors

Generates a relationship

$$ER = \text{« activity »} = f(p_1, p_2, \dots, p_n) \quad f \text{ linear or non-linear function}$$

Molecular descriptors are intrinsic molecular features

Ex: Molar Weight (MW)      Hydrophobicity (AlogP98)  
Polarisability (Apol)      Dipolar Moment (Dipole -X, Y, Z)

Surface Descriptors:

« Jurs » descriptors surface and charged surface descriptors

Determination of the partition coefficients for yogurt

- headspace analysis at equilibrium using solid-phase micro extraction.


QSPR study:

- molecular descriptors calculations
- equations constructions

carried out using Cerius<sup>2</sup> (Accelrys Inc.)

$$K_{ret} = \frac{\text{chromatographic area reference}}{\text{chromatographic area thickeners}}$$

$$JursWNSA-1 = \sum_s \text{surfaces areas of all negatively charged atoms} \times \frac{\text{total molecular solvent-accessible area}}{1000}$$



20 descriptors selected → present at least a dependable correlation with the partition coefficient values  $K_{ret}$  (Starch 0.1%)  
For higher amounts of starch (0.4 and 0.6%) and pectin (0.04, 0.1, and 0.15%)

all of the previous correlations are severely disturbed

no reliable correlation between the descriptors and  $\log K_{ret}$  values

surface descriptor Jurs WNSA-1 is a particular interest

Jurs WNSA-1 encodes the surface negatively charged area weighted by the total surface area

For the same chemical class negatively charged areas similar


- lower values → smallest molecules → low hydrophobicity
- higher values → biggest molecule → high hydrophobicity

only true for Starch 0.1%

Increase in the amount of starch disrupts "hydrophobic effect" in favour of other natures of interactions

Jurs WNSA-1 values highlights the existence of two groups:

- ethyl hexanoate, hexyl acetate, ethyl octanoate, and ethyl decanoate
- butanol, hexanol, octanol, decanol and methyl 2-methylbutanoate



**Conclusion : QSPR study** shows an intricate role due to adding both starch at concentrations higher than 0.1% and pectin puts forward the power of negatively charged surface areas

suggests the existence of competitive interactions, hydrophobic and polar binding

QSPR approach constitutes a promising tool for characterizing the interactions involved in complex food matrices

Experiments are now in progress to obtain true partition coefficient values involving a larger range of chemical structures