



HAL
open science

Potential indirect impact of the invasive leaf-miner, *Cameraria ohridella*, on native leaf-miners

Christelle Péré, Sylvie Augustin, Rumen Tomov, Ted Turlings, Marc Kenis

► To cite this version:

Christelle Péré, Sylvie Augustin, Rumen Tomov, Ted Turlings, Marc Kenis. Potential indirect impact of the invasive leaf-miner, *Cameraria ohridella*, on native leaf-miners. 23. International Congress of Entomology; ICE 2008, Jul 2008, Durban, France. p. 843. hal-02821649

HAL Id: hal-02821649

<https://hal.inrae.fr/hal-02821649>

Submitted on 6 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Potential indirect impact of the invasive leaf-miner, *Cameraria ohridella*, on native leaf-miners

Christelle Péré¹, Sylvie Augustin³, Rumen Tomov⁴, Ted Turlings², Marc Kenis¹

¹CABI Europe-Switzerland, Rue des Grillons 1, 2800 Delémont, Switzerland, ²University of Neuchâtel, rue Emile Argand 11, CP 158, 2009 Neuchâtel, Switzerland, ³INRA, 2163 av. de la Pomme de Pin, BP 20619, Ardon, 45166 Olivet Cedex, France, ⁴University of Forestry, Bul. Kliment Ohridski 10, 1756 Sofia, Bulgaria

Introduction: The horse-chestnut leaf-miner, *Cameraria ohridella*, is a moth of unknown origin that was first observed in Macedonia in 1984 and has now invaded most of Europe. It is attacked by several native leaf-miners' parasitoids. Despite the low parasitism rates observed in *C. ohridella*, populations are so high that an unusual amount of polyphagous parasitoids are produced in the vicinity of infested horse-chestnut trees two or three times per year. In spring, the bulk of the parasitoids emerge at least six weeks before the first suitable *C. ohridella* larvae or pupae are available. These parasitoids could massively attack the first indigenous leaf-miners developing in spring. We tested the hypothesis that the presence of *C. ohridella* has a negative effect on populations of native leaf-miners through shared natural enemies (apparent competition).

Methods: Species richness and abundance of indigenous leaf-miners were assessed in presence and absence of *C. ohridella*, in Switzerland, France and Bulgaria.

Results: In spring, the species richness of indigenous leaf-miners was significantly lower in the presence of *C. ohridella* than in its absence. Some native leaf-miners, such as the beech and oak leaf-mining weevils, *Orchestes fagi* and *O. quercus*, sharing their parasitoid complex with *C. ohridella*, were significantly less abundant at sites with horse-chestnuts attacked by *C. ohridella* compared to control sites.

Conclusion: Parasitoids, and possibly predators, are the only link between *C. ohridella* and native leaf-miners feeding on other host-plants and are thus probably responsible for the decrease.