



HAL
open science

Annotation of the immunity-related genes in the pig genome

. Immune Response Annotation Group (irag), Claire Rogel Gaillard, Christopher K. Tuggle, Jane Loveland, Jennifer Harrow, Elisabetta Giuffra, Joan Lunney, David Hume, Alan Archibald, Harry Dawson, et al.

► To cite this version:

. Immune Response Annotation Group (irag), Claire Rogel Gaillard, Christopher K. Tuggle, Jane Loveland, Jennifer Harrow, et al.. Annotation of the immunity-related genes in the pig genome. 32nd Conference of the International Society of Animal Genetics (ISAG), Jul 2010, Edinburgh, United Kingdom. 2010. hal-02821666

HAL Id: hal-02821666

<https://hal.inrae.fr/hal-02821666>

Submitted on 6 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

P5053 Whole genome re-sequencing of Japanese Black cattle for SNP discovery in critical regions for *Marbling-2*, *CW-2* and *FMA*

Takashi Hirano¹, Toshio Watanabe¹, Tetsuo Kunieda², Yoshikazu Sugimoto¹ and Akiko Takasuga¹.

¹Shirakawa Institute of Animal Genetics, Nishi-shirakawa, Fukushima, Japan.

²Graduate School of Natural Science and technology, Okayama University, Okayama, Japan.

Marbling-2 and *CW-2* are QTL for marbling and carcass weight, and *FMA* is a genetic disease for Forelimb-girdle Muscular Anomaly. *Marbling-2* and *CW-2* have been mapped with progenies of Bull A, while *FMA* has been mapped with a progenies of Bull B. The critical regions for *Marbling-2*, *CW-2* and *FMA* were 4.6-Mb (BTA7), 0.9-Mb (BTA6) and 2.4-Mb (BTA26), respectively. To identify responsible/causative genes or SNPs from these relatively large regions, we performed whole genome re-sequencing of Bull A and B that are maternal half-sib.

A single-end library and three size-different paired-end libraries were prepared according to the manufacture's protocol, followed by sequencing with 36-bases reads using Genome Analyzer II (Illumina).

More than 3×10^9 reads were generated from Bull A and B, respectively, of which 11.5-M reads (412.3-Mb; x 52.03) and 11.7-M reads (415.3-Mb; x 58.96) were mapped to the target regions. Nearly 90% of the target regions were covered by ≥ 10 depth in both Bull A and B, and 21,136 (Bull A) and 19,598 SNPs (Bull B) were detected in comparison with Btau4.0 reference sequence. Based on criteria that Bull A is heterozygous for *Marbling-2* and *CW-2*, and Bull B is heterozygous for *FMA*, responsible or causative SNP will be searched among the SNPs.

P5054 Annotation of the immunity-related genes in the pig genome

Immune Response Annotation Group (IRAG).

Coordination: Claire Rogel-Gaillard¹ and Christopher K Tuggle².

Participants: Jane Loveland³, Jennifer Harrow³, Elisabetta Giuffra⁴, Joan Lunney⁵, David Hume⁶, Alan Archibald⁶, Harry Dawson⁷, Jim Reecy⁸, Hirohide Uenishi⁹, Takeya Morozumi⁹, Hiroki Shinkai⁹, Dennis Prickett¹⁰, Shuhong Zhao¹¹, Yongming Sang¹², Frank Blecha¹², Guolong Zhang¹³, Bertrand Bed'hom¹, Ryan Cheng², Yasiara Rodriguez², Anna Anselmo⁴, Bouabid Badaoui⁴, Celine Chen⁷, Hongbo Chen⁶, Samuel Abrams⁵, Dario Beraldi⁶, Ronan Kapetanovic⁶, Zhi-Liang Hu⁶, and the Swine Genome Sequencing Consortium.

¹INRA, UMR de Génétique Animale et Biologie Intégrative, Jouy-en-Josas, France.

²Department of Animal Science and Center for Integrated Animal Genomics, Iowa State University, Ames, IA, USA.

³Wellcome Trust Sanger Institute, Hinxton, Cambridgeshire CB10 1HH, UK.

⁴Parco Tecnologico Padano, Livestock Genomics Unit, Polo Universitario, 26900 Lodi, Italy.

⁵USDA, ARS, BARC, Animal Parasitic Diseases Laboratory, Beltsville, MD USA.

⁶The Roslin Institute and Royal (Dick) School of Veterinary Studies, University of Edinburgh, Roslin, Midlothian, EH25 9PS UK.

⁷Diet, Genomics and Immunology Laboratory, Beltsville Human Nutrition Research Center, ARS, USDA, Beltsville, MD.

⁸Department of Animal Science and NRSP-8 Bioinformatics Coordination project, Iowa State University, Ames, IA, USA.

⁹Animal Genome Research Program, NIAS/STAFF, Tsukuba, Ibaraki, Japan.

¹⁰Institute for Animal Health, Compton, nr Newbury, RG20 7NN, UK.

¹¹Key Lab of Agricultural Animal Genetics, Breeding, and Reproduction of Ministry of Education, Huazhong Agricultural University, Wuhan 430070, PR China.

¹²College of Veterinary Medicine, Kansas State University, Manhattan, KS USA.

¹³Department of Animal Science, Oklahoma State University, Stillwater, OK 74078, USA.

Corresponding authors: Christopher K Tuggle: cktuggle@iastate.edu;

Claire Rogel-Gaillard: claire.rogel-gaillard@jouy.inra.fr

Resistance to disease and control of immune responses are complex processes that may be studied at several levels including individuals, populations, species, and phylogenetic groups. Indeed, there are mechanisms and responses shared between species and others that are species-specific, partly due to co-evolution of hosts and pathogens. Large scale sequencing of complex genomes provides unique opportunities to annotate large sets of genes and their structural variations for further comparative analyses. An automatically annotated draft assembly of the pig genome was released with Ensembl 56 in October 2009. The Immune Response Annotation Group (IRAG), comprising scientists working on resistance to disease and immunity in swine, was established to identify shared and species-specific

immune responses and refine the annotation of immunity-related genes. A list of close to 1700 genes was drawn using information from gene ontology annotation (GO: 0002376 for immune system process), a core set of genes involved in host pathogen interplay (Jenner and Young, 2005), and gene sets under positive selection in humans (Barreiro and Quintana-Murci, 2010) and cattle. Manual annotation of these genes has begun using the WTSI Otterlace software. Alignments of genomic contigs with publicly available mRNAs, ESTs and proteins are analyzed to delineate exon and intron boundaries, thus providing a refined functional annotation of genes. Preliminary results have identified gene duplications and many new alternative splice variants for known genes. Our future challenge will be to identify the underlying biology specific to the pig species that can be identified from this major annotation effort.

P5055 Gene expression and pathways analysis of white blood cells from cattle orally challenged with Bovine Amyloidotic Spongiform Encephalopathy (BASE) one year post-infection

Caterina Cambuli¹, Simona Panelli², Francesco Strozzi², Rossana Capoferri¹, Ilaria Barbieri³, Nicola Martinelli³, Lorenzo Capucci³, Guerino Lombardi³ and John L. Williams².

¹IDRA-LAB, Istituto Sperimentale Italiano "L. Spallanzani", Via Einstein, 26900-Lodi, Italy.

²IDRA-LAB, Parco Tecnologico Padano, Via Einstein, 26900-Lodi, Italy.

³Istituto Zooprofilattico Sperimentale della Lombardia e dell'Emilia Romagna, via Bianchi 9, 25124, Brescia, Italy.

Corresponding author: john.williams@tecnoparco.org

Bovine Amyloidotic Spongiform Encephalopathy (BASE) is a recently discovered atypical form of BSE, transmissible to primates, and has been proposed as an equivalent of sporadic Creutzfeldt-Jacob Disease in man. Although transmissible, it is unknown whether BASE is acquired through infection or arises spontaneously. In the present study, the gene expression patterns of white blood cells (WBCs) from five cattle one year after oral BASE challenge were compared with negative controls using a custom microarray containing 43,768 unique gene probes.

A total of 140 genes were found to be differentially expressed between BASE and control animals with a log fold change of 1.5 or greater. Of these, 70% were up-regulated in the infected animals. Microarray data were then confirmed by qRT-PCR. The majority of the differentially expressed genes are related to immune functions, and several belong to the same pathways as revealed by KEGG analysis. In particular, BASE animals appear to have significantly modified expression of genes linked to the differentiation of subsets of immune cells, T and B cell development and activation, Natural Killer activity and inflammatory response. Samples from the same animals were examined by qRT-PCR at different time-points for the most interesting pathways to define when changes in gene expression are first observed and their kinetics.

The potential impacts of these gene expression changes will be discussed, together with their potential use as biomarkers and their contribution to better define PrP related gene functions in immune cells.