

**Viral determinant implicated in the transmission of the  
Grapevine fanleaf virus by its nematode vector,  
Xiphinema index**

Pascale Schellenberger, Gerard Demangeat, Peggy Andret-Link, Corinne Keichinger, Marc Bergdoll, Patrick Bron, Bernard Lorber, Olivier Lemaire, Christophe Ritzenthaler

► **To cite this version:**

Pascale Schellenberger, Gerard Demangeat, Peggy Andret-Link, Corinne Keichinger, Marc Bergdoll, et al.. Viral determinant implicated in the transmission of the Grapevine fanleaf virus by its nematode vector, Xiphinema index. Genetic Control of Plant Pathogenic Viruses, and their Vectors, in European Crops, Nov 2008, Puerto de Santa María, Spain. ResistVir, n.p., 2008. hal-02822312

**HAL Id: hal-02822312**

**<https://hal.inrae.fr/hal-02822312>**

Submitted on 6 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**(P2-17) Viral determinant implicated in the transmission of the *Grapevine fanleaf virus* by its nematode vector, *Xiphinema index***

Pascale Schellenberger<sup>1,2</sup>, Gérard Demangeat<sup>1</sup>, Peggy Andret-Link<sup>1</sup>, Corinne Keichinger<sup>2</sup>, Marc Bergdoll<sup>2</sup>, Patrick Bron<sup>3</sup>, Bernard Lorber<sup>4</sup>, Olivier Lemaire<sup>1</sup> and Christophe Ritzenthaler<sup>2</sup>

<sup>1</sup>Institut National de la Recherche Agronomique, UMR 1131 Santé de la Vigne et Qualité du Vin, INRA/ULP, Laboratoire de Virologie et Vecton, 28 rue de Herrlisheim 68021 Colmar (France).  
E-mail: [pschellenberger@colmar.inra.fr](mailto:pschellenberger@colmar.inra.fr)

<sup>2</sup>Institut de Biologie Moléculaire des Plantes (IBMP), UPR 2357 du CNRS, 12 rue du Général Zimmer 67084 Strasbourg Cedex (France)

<sup>3</sup>Université de Rennes I, UMR 6026 du CNRS, 263 avenue du Général Leclerc 35042 Rennes Cedex (France)

<sup>4</sup>Institut de Biologie Moléculaire et Cellulaire (IBMC), UPR 9002 du CNRS, 15 rue R. Descartes 67084 Strasbourg Cedex (France)

*Grapevine fanleaf virus* (GFLV; genus *Nepovirus*, family *Comoviridae*) is an icosahedral virus with a positive sense bipartite RNA. GFLV is specifically transmitted from grapevine to grapevine by the ectoparasitic nematode *Xiphinema index* (Andret-Link *et al.*, 2004, *J. Plant Path.* 86, 183-195). Previous experiments replacing the viral GFLV coat protein (CP) gene by the CP gene of *Arabidopsis mosaic virus* (ArMV), a closely related nepovirus specifically transmitted by *Xiphinema diversicaudatum*, indicates that the specificity of transmission is solely determined by the CP (Andret *et al.*, 2004, *Virology* 320, 12-22). The objective of our study is to identify the determinants of transmission specificity on CP. We hypothesized that amino acids involved in the specificity of transmission should be different between both viruses and located at the external surface of the capsids in order to interact with potential receptors in the food canal of nematodes.

To identify divergent external amino acids, a 3D-structural model of GFLV capsid has been deduced from the 3.5 Å resolution structure of *Tobacco ringspot virus* (Chandrasekar & Johnson 1998, *Structure* 6, 157-171). Based on this model, five potential external domains, from 6 to 12 amino acids, were selected. Sixteen mutants in which these five domains were substituted either in single or multiple combinations by their ArMV counterpart domains, have been engineered. Two mutants were able to induce a systemic infection in plants. Only one led to the loss of transmission by *X. index*, indicating a potential function of this domain in GFLV transmission.

To fully validate our model, structural analyses (cryo-electron microscopy and X-ray crystallography) are performed in parallel. For cryo-electron microscopy, images of frozen-hydrated purified viral particles of GFLV and ArMV were recorded and are analysed using the polar Fourier transforms method developed by Baker. The crystallographic study is done with GFLV purified particles. The size homogeneity of GFLV particles have been analysed by dynamic light scattering (DLS) and revealed to be highly homogeneous. Several crystallization conditions have also been tested and have resulted in viral crystals which will be used for X-ray diffraction and modelling studies.