



**HAL**  
open science

## Does pea/cereal intercropping have an effect on Ascochyta blight epidemic development ?

Alexandra Schoeny, Stéphane Jumel, Francois Rouault, A. Dibet, Joëlle Fustec, J. Kinane, M. Lyngkjaer, M. Fernandez-Aparicio, Diego Rubiales,  
Bernard B. Tivoli

### ► To cite this version:

Alexandra Schoeny, Stéphane Jumel, Francois Rouault, A. Dibet, Joëlle Fustec, et al.. Does pea/cereal intercropping have an effect on Ascochyta blight epidemic development?. 6. European Conference grain legumes, Nov 2007, Lisbon, Portugal. 2 p. hal-02823352

**HAL Id: hal-02823352**

**<https://hal.inrae.fr/hal-02823352>**

Submitted on 6 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Posters 269, 270 & 271

## Assessment of winter legume-cereals intercrops in France

Marie-Hélène Jeuffroy

INRA UMR Agronomie (211) INRA-AgroParisTech, BP01, 78 850 Thiverval-Grignon, France

Poster 160

## Comparison of spring and winter cereal-legume intercrops in organic farming by analysis of interactions between species

Christophe Naudin<sup>1\*</sup>, Anne Aveline<sup>1</sup>, Guénaëlle Corre-Hellou<sup>1</sup>, Audrey Dibet<sup>1</sup>, Marie-Hélène Jeuffroy<sup>2</sup> and Yves Crozat<sup>1</sup>.

1. Laboratoire d'Ecophysiologie Végétale et Agroécologie, Ecole Supérieure d'Agriculture, 55 rue Rabelais, BP 30748, 49007 Angers cedex 01, France.

2. UMR 211 Agronomie INRA AgroParisTech, BP01, 78850 Thiverval-Grignon, France.

\* correspondence: c.naudin@groupe-esa.com

Keywords: intercropping, sowing date, competition, complementarity, pea, wheat, barley.

Cereal-legume intercrops are gaining increasing interest in Europe in low input farming systems. In numerous previous studies, intercropping has been shown to result in a higher or equal yield, compared to respective sole crops (1) (2). The comparison between spring and winter intercrops has rarely been studied. Field trials were carried out to compare the performance of spring and winter intercrops in organic farming over three years. Spring pea (*Pisum sativum* L., cv Baccara), winter pea (*Pisum sativum* L., cv Lucy), spring barley (*Hordeum vulgare* L., cv. Scarlett) and winter wheat (*Triticum aestivum* L., cv. Apache) were grown in sole crops and in additive and in substitutive intercrops, in 2003, 2004, and 2005. Analysis of net biodiversity effect ( $\Delta$ ) of intercrops on grain yields in comparison with sole crops was performed using partitioning selection effect (competition) and complementarity effect (both facilitation and niche differentiation) (3).

The yields were higher for winter than spring crops. A yield gain was observed for intercrops compared to sole crops in both winter and spring crops but the types of interaction involved were not similar. The net biodiversity effect ( $\Delta$ ) was mainly correlated with a complementarity effect in spring crops and with a selection effect in winter crops. The variability of these effects was largely explained by cereal density observed after emergence, whereas pea plant density was not determinant. Crop accidents during crop establishment were frequent in winter crops in our study and affected plant densities of pea or wheat. Finally, partitioning methodology proved useful for post-harvest analysis of intercrop performances in comparison with that of sole crops.

(1) Corre-Hellou, G., Fustec, J. and Crozat, Y., 2006. Interspecific Competition for Soil N and its Interaction with N<sub>2</sub> Fixation, Leaf Expansion and Crop Growth in Pea-Barley Intercrops. *Plant and Soil* 282, 195-208.

(2) Jensen, E.S., 1996. Grain yield, symbiotic N<sub>2</sub> fixation and interspecific competition for inorganic N in pea-barley intercrops. *Plant and Soil* 182, 25-38.

(3) Loreau, M. and Hector, A., 2001. Partitioning selection and complementarity in biodiversity experiments. *Nature* 412, 72-76.

Posters 244 & 311

## Effects of intercropping on Orobanche control

Mónica Fernández-Aparicio and Diego Rubiales

Institute of Sustainable Agriculture, CSIC, Apdo. 4084, 14080 Córdoba, Spain.

Poster 303

## Does pea/cereal intercropping have an effect on Ascochyta blight epidemic development?

A. Schoeny<sup>\*,\*\*</sup>, S. Jumel<sup>\*</sup>, F. Rouault<sup>\*</sup>, A. Dibet<sup>\*\*\*</sup>, J. Fustec<sup>\*\*\*</sup>, J. Kinane<sup>\*\*\*\*</sup>, M. Lyngkjær<sup>\*\*\*\*</sup>, M. Fernández-Aparicio<sup>\*\*\*\*\*</sup>, D. Rubiales<sup>\*\*\*\*\*</sup> and B. Tivoli<sup>\*</sup>

\* INRA, UMR1099 Biologie des Organismes et des Populations appliquée à la Protection des Plantes, Le Rheu (France), alexandra.schoeny@avignon.inra.fr

\*\* INRA, UR407 Pathologie Végétale, Montfavet (France)

\*\*\* Ecole Supérieure d'Agriculture - Laboratoire d'Ecophysiologie végétale et Agroécologie, Angers (France)

\*\*\*\* RISØ National Laboratory, DTU - Biosystems Department, Roskilde (Denmark)

\*\*\*\*\* CSIC - Institute of Sustainable Agriculture, Córdoba (Spain)

Keywords: Ascochyta blight, cereal, disease development, intercropping, microclimate, pea

Ascochyta blight (*Mycosphaerella pinodes*/*Phoma medicaginis* var. *pinodella*/Ascochyta pisi) is one of the most damaging diseases of field peas world-wide. Varieties with robust resistance are not available. Fungicide use and agronomic practices

such as burial or destruction of infected stubble, adoption of a suitable crop rotation, spatial separation of fields from past seasons' infected stubble are not always practiced and may not be suitable for many farm situations particularly in organic farming. Alternative practices have thus to be developed. Using plant and canopy architectural features could be a possible lead to impact disease development through modification of pathogen movement within the canopy and/or modification of microclimate.

Intercropping is the agricultural practice of cultivating two or more crops in the same space at the same time (1). The two or more crops used in an intercrop may be from different species and different plant families, or they may simply be different varieties or cultivars of the same crop species. Growing legume and cereal species simultaneously in the same field can increase the use efficiencies of growth resources and reduce fertilizer N requirements (2). Considering that it also modifies drastically plant and canopy architecture, we investigated if pea/cereal intercropping could contribute to *Ascochyta* blight control.

Field experiments were conducted in three countries (Denmark, France and Spain) between 2004 and 2006. Row intercrops involving either winter pea/winter wheat (F and SP) or spring pea/spring barley (F and DK) were compared to pea sole crops according to a coordinated protocol. Disease levels in pea sole crops were variable according to sites and years. They were low in several experiments even with artificial inoculation causing difficulties in determining the effect of cropping method. Disease progression on stipules was on average slightly reduced in intercrop compared to pea sole crop but effects were rarely significant. Disease reduction was stronger on pods and stems late in the season. Monitoring microclimate within canopies showed variable results according to the sensor type, sensor location and monitoring period. Late in the season, leaf wetness duration appeared to be slightly shorter within the intercrop compared to the pea sole. This could explain the reduction in disease level on pods and stems.

Finally, the evaluation of the relevance of pea/cereals intercropping for the management of *Ascochyta* blight has to be placed in the wider framework of sustainable farming taking in consideration all the multifunctional roles of intercropping (addition and recycling of organic material, water management, protection of soil from erosion and pest or disease suppression).

(1) Andrews D.J. and Kassam A.H. (1976) The importance of multiple cropping in increasing world food supplies. pp. 1-10 in R.I. Papendick, A. Sanchez, G.B. Triplett (Eds.), *Multiple Cropping*. ASA Special Publication 27. American Society of Agronomy, Madison, WI.

(2) Francis C.A. (1986). *Multiple Cropping Systems*. MacMillan. New York. 383 p.