



The first finding of Wolbachia in thelytokous seed chalcids (Hymenoptera: Torymidae, Megastigmus sp)

Thomas Boivin, Jean-Noël Candau, Alain Roques, Marie-Anne Auger-Rozenberg

► To cite this version:

Thomas Boivin, Jean-Noël Candau, Alain Roques, Marie-Anne Auger-Rozenberg. The first finding of Wolbachia in thelytokous seed chalcids (Hymenoptera: Torymidae, Megastigmus sp). 23. International Congress of Entomology; ICE 2008, Jul 2008, Durban, South Africa. 1 p. hal-02823868

HAL Id: hal-02823868

<https://hal.inrae.fr/hal-02823868>

Submitted on 6 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



THOMAS BOIVIN¹, JEAN-NOEL CANDAU¹, ALAIN ROQUES² and MARIE-ANNE AUGER-ROZENBERG²

¹INRA, UR 629 Ecologie des Forêts Méditerranéennes, Domaine Saint Paul, Site Agroparc, 84914 Avignon Cedex 9, France

²INRA, UR 0633 Zoologie forestière, 2163 Avenue de la Pomme de Pin, CS40001 Ardon, 45075 Orléans Cedex, France

In haplodiploid insect species, thelytokous parthenogenesis may be associated with bacterial endosymbionts that manipulate their host reproduction. Such an association was tested in the highly invasive genus *Megastigmus*, within which thelytokous parthenogenesis is much less frequent than arrhenotoky, which is the classical way of reproduction in chalcid insects.

SCREENING FOR REPRODUCTION-ASSOCIATED ENDOSYMBIOTS AMONG *MEGASTIGMUS* SPECIES

Specific primers were used for the screening of *Wolbachia* (*wsp*), *Cardinium* (16S rDNA), *Rickettsia* (16S rRNA), and *Arsenophonus* (23S rDNA) endosymbionts in seven thelytokous and three arrhenotokous *Megastigmus* species. PCR amplification with *Wolbachia* primers was successful in all of the seven thelytokous *Megastigmus* species while no reaction occurred using *Cardinium*, *Rickettsia* and *Arsenophonus* primers. None of the three arrhenotokous species tested allowed amplification whatever the primer (Table 1).



ASSOCIATION BETWEEN THELYTOKY AND WOLBACHIA ENDOSYMBIOTS

Both *Wolbachia* infection and progeny sex ratios were compared between non- and antibiotic-treated females of *M. pinsapinis*. Seven-day tetracycline treatments of *M. pinsapinis* females resulted in a loss of *Wolbachia* infection and in a return to an arrhenotokous mode of reproduction as they mainly produced a male progeny using asexual reproduction (Table 2).



TABLE 1. Detection tests of endosymbionts associated with host reproduction manipulation in 8 thelytokous and 3 arrhenotokous *Megastigmus* species.

Species (N)	Reproduction	Infection %			
		<i>Wolbachia</i>	<i>Cardinium</i>	<i>Arsenophonus</i>	<i>Rickettsia</i>
<i>M. rosae</i> (30)	thelytoky	100	0	0	0
<i>M. aculeatus</i> (43)	thelytoky	100	0	0	0
<i>M. brevicaudis</i> (1)	thelytoky	100	not tested	not tested	not tested
<i>M. pinsapinis</i> (60)	thelytoky	100	0	0	0
<i>M. suspectus</i> (65)	thelytoky	100	0	0	0
<i>M. pictus</i> (30)	thelytoky	100	0	0	0
<i>M. pistaciae</i> (30)	thelytoky	100	0	0	0
<i>M. alba</i> n.sp. (5)	thelytoky	100	not tested	not tested	not tested
<i>M. spermotrophus</i> (30)	arrhenotoky	0	0	0	0
<i>M. rafni</i> (30)	arrhenotoky	0	0	0	0
<i>M. schimitscheki</i> (30)	arrhenotoky	0	0	0	0

TABLE 2. Effects of antibiotic treatments on the infection of *M. pinsapinis* females by *Wolbachia* and their male production following asexual reproduction. (Control = sugared water ; Antibiotic = Tetracycline (2%) + sugared water)

Treatment mode	Treatment duration (in days)	Infection % by <i>Wolbachia</i> after treatment (N females)		Frequency of males in the progeny \pm S.E. (N larvae)
		Control	Antibiotic	
Control	3	100 (38)		0 (127)
	7	100 (44)		0 (112)
Antibiotic	3	100 (42)		0 (109)
	7	0 (45)		95.4 \pm 2.6 (132)

PHYLOGENY OF WOLBACHIA STRAINS INFECTING THELYTOKOUS *MEGASTIGMUS* SPECIES

The sequences of the *Wolbachia* *wsp* fragments found in the different thelytokous *Megastigmus* species were used to assess the evolutionary trajectories of their association. Molecular phylogeny of *Wolbachia* variants paralleled the observed radiation of *Megastigmus* species with respect to host-plant families (Pinaceae, Rosaceae and Anacardiaceae), but no significant divergence in *Wolbachia* sequences was found within host-plant families (Figure 1).



CONCLUSIONS

Thelytokous parthenogenesis in *Megastigmus* species may be associated with *Wolbachia* infection. Phylogenetic analyses indicate a possible partition of the studied *Wolbachia* strains in three clades that fit the host-plant structuration of their insect host. This suggests that horizontal transfers of *Wolbachia* may have occurred between *Megastigmus* species exploiting closely related host-plants. This may raise critical questions within the context of biological invasions in this insect genus, which frequently generate novel cases of sympatry and thus novel ecological connexions that may influence the spread of *Wolbachia*.

Figure 1. Phylogenetic relationships between sequences of the *Wolbachia* *wsp* fragments found in thelytokous *Megastigmus* species.

