



Mating success diminished by isolation in space and time

Christelle Robinet, David Gray, Andrew Liebhold

► To cite this version:

Christelle Robinet, David Gray, Andrew Liebhold. Mating success diminished by isolation in space and time. Réunion annuelle de la société d'entomologie du Canada et de la société d'entomologie du Québec;Joint annual meeting of the entomological society of canada and the société d'entomologie du Québec, Nov 2006, Montréal, Canada. 1 p., 2006. hal-02824064

HAL Id: hal-02824064

<https://hal.inrae.fr/hal-02824064>

Submitted on 6 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



MATING SUCCESS DIMINISHED BY ISOLATION IN SPACE AND TIME

Lost in time : temporal asynchrony

OBJECTIVES

To explore the effects of
1) dispersion of sexual maturation times,
and
2) time-lag between male and female mean
maturation (protandry)
on mating success.

METHODS

1) **Generalized model:** Distributions of ♂ and ♀ sexual maturation times were generated using Gaussian density functions with varying standard deviation and varying lags between the mean time of ♂ and ♀ maturation. Mating probability was derived from a negative exponential model (eq. 1) where P_d is the probability that a female is mated on day d given the number of mature males M_d on this day (depending on the daily rate of males becoming sexually mature, male longevity and the total number of males):

$$P_d = 1 - \exp(-s M_d) \quad (\text{eq. 1})$$

Population-level mating success MS was based upon the temporal distribution of females calculated from the daily rate of female sexual maturation and female longevity.

2) **Gypsy moth case study:** emergence curves were estimated using a Life Stage (GLS) Model (Gray 2004) based upon daily temperature. For each day, we estimated the probability that each female was mated using a model derived from Sharov *et al.* (1995) (eq. 1 with $s=0.15$). Abundance of males was described by the season-long trap capture of males (TC) and it was set constant over space for simplification. Mating success was calculated using weather station data from 7,591 locations across North America from 1960 to 2000.

RESULTS

1) **Generalized model:** both dispersion in developmental time and lags in male and female maturation affected mating success (Fig. 1). Females that matured late, or near the mean time, had a negligible mating probability (due to lack of males) whereas the females maturing earliest were the main contributors to population growth.

2) **Gypsy moth case study:** in addition to the mating failure in extreme northern and southern areas caused by low or no emergence of adults, mating success appears non uniformly distributed across North America (Fig. 2). A central zone where mating is most successful stretches across the USA.

The Rocky Mountain area is clearly unfavorable for mate-finding (long time lag between males and females: 8.1 days, and short dispersion of emergence curves: $sd=1.9$ days in Jackson, WY). Surprisingly, coastal California was also relatively unfavorable (small time lag of 2.2 days but extremely low dispersion of emergence times: $sd>1$ day in Santa Barbara, CA). However, Fig. 1 indicates that a slight increase in developmental variability would enable much higher mating success in such a case.




Figure 2: Average mating success of gypsy moth over 1960-2000, based on a single oviposition date (August 1), a season-long trap capture of 100 males, emergence curves predicted by the GLS phenology model and the mating model developed in this study.

C. Robinet^{1,2}, D. Gray³ & A. Liebhold²

¹ INRA – Station de Zoologie Forestière – Avenue de la Pomme de Pin – Ardon – 45166 Olivet – France

² Northern Research Station – USDA Forest Service – 180 Canfield St. – Morgantown – WV26505 – USA

³ Natural Resources Canada, Canadian Forest Service, Landscape Disturbances, 1350 Regent St., Fredericton, NB E3B 2G6 – Canada
contacts: robinet@orelans.inra.fr; dgray@nrcan.gc.ca; aliebhold@fs.fed.us

INTRODUCTION

Various processes affecting low density populations may generate **Allee effects** and profoundly affect population persistence. Understanding these effects could improve conservation strategies and invasion control. While mate-finding is often mentioned in the literature as a cause of Allee dynamics, the underlying mechanisms are still unclear. Mating failure can result from both **spatial and temporal isolation of males and females**. Here we address both issues and more particularly the temporal issue since it has received little attention up to now.

Lost in space : stochastic diffusion

OBJECTIVES

- 1) Investigate the effects of stochastic diffusion in mate-finding in low density populations
- 2) Determine a threshold number of introduced individuals that enable a population to persist.

METHODS

Assuming that n egg masses (each composed of 100–250 eggs of each sex) are introduced in a new area, we coupled the following stochastic processes to model the mating success of the Gypsy Moth:

- **Diffusive dispersal** ($D=0.003$ km²/generation, Shigesada & Kawasaki 1997) in two spatial dimensions, and for each larva, we identified the cell to which they dispersed according to the probability distribution.
- **Mortality rate of 95%** was used to simulate whether each individual survived.
- **Emergence time**: for each individual, we simulated the emergence day from the emergence probability distribution predicted by the GLS model (see "lost in time"), for example in Farmville, VA, in 1997.
- **Mating success**: we constructed a regression model based on experimental data of release-recapture (D. Lance, USDA) to determine the probability that a male mates a female at a given distance and at a given day after its emergence.

For each iteration, we calculated the number of females effectively mated and the growth rate of the population.

RESULTS

Since each process was modeled using stochastic simulation, we observed a high variability in the mating success but globally a positive relationship between population density and population growth. When there were initially 250 eggs of each sex per egg mass (common in low density populations), a strong Allee effect was not detected (growth rate always >1). But when there were 150 eggs of each sex per egg mass (typical of high density populations), we could identify a threshold of 3 egg masses to enable the population to persist (Fig. 3).




Figure 3: Effect of the number of egg masses introduced on the population growth rate (150 eggs of each sex per egg mass, 50 simulations).

DISCUSSION & CONCLUSION

Even though reproductive asynchrony has been mainly considered as an advantage for populations, especially in presence of strong environmental stochasticities, it also may enhance Allee effects and be responsible of population declines. This temporal isolation of males and females, coupled with the difficulty males to find each other over space, is a fundamental mechanism underlying the failure of populations to establish.

Allee effects are known to affect the establishment and spread of invading gypsy moth populations (Liebhold and Bascompte 2003, Tobin *et al.* 2006). Our study identifies more precisely the possible causes and describes a spatial gradient in mating success due to a climatic heterogeneity. In comparison to climatic suitability for completion of gypsy moth development (Gray 2004), mating heterogeneity is quite different and suggests a complementary factor in establishment failure.

References:

- Gray DR (2004) The gypsy moth life stage model: landscape-wide estimates of gypsy moth establishment using a multi-generational model. *Ecol. Model.* 176:155-171.
- Liebhold A. & Bascompte J. (2003) The Allee effect, stochastic dynamics and the eradication of alien species. *Ecology Letters* 6:133-140.
- Sharov AA, Liebhold AM & Ravlin FW (1995) Prediction of gypsy moth (Lepidoptera: Lymantriidae) mating success from pheromone trap counts. *Environ. Entomol.* 24:1239-1244.
- Shigesada N & Kawasaki K (1997) *Biological Invasions: Theory and Practice*, in: Oxford Series in Ecology and Evolution, Oxford University Press, Oxford.
- Tobin PC, Whitmire SL, Johnson DM, Bjornstad ON & Liebhold AM. Invasion speed is affected by geographic variation in the strength of Allee effects. *Ecology Letters* (in press)