



**HAL**  
open science

## Carpel development

Cristina Ferrandiz, Chloé Fourquin, Nathanael Prunet, C.P. Scutt, Eva Sundberg, Christophe Trehin, Aurélie Vialette-Guiraud

► **To cite this version:**

Cristina Ferrandiz, Chloé Fourquin, Nathanael Prunet, C.P. Scutt, Eva Sundberg, et al.. Carpel development. *Advances in Botanical Research*, 55, Academic Press, 2010, *Advances in Botanical Research*, 978-0-12-380868-4. 10.1016/S0065-2296(10)55001-8 . hal-02824413

**HAL Id: hal-02824413**

**<https://hal.inrae.fr/hal-02824413v1>**

Submitted on 11 Feb 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Elsevier Editorial System(tm) for Advances in Botanical Research  
Manuscript Draft

Manuscript Number:

Title: Carpel Development

Article Type: 2009

Section/Category:

Keywords: carpel, gynoecium, angiosperm, flower, development, evolution

Corresponding Author: Dr. Charlie P SCUTT,

Corresponding Author's Institution:

First Author: Cristina Ferrandiz

Order of Authors: Cristina Ferrandiz; Chloe Fourquin; Nathanael Prunet; Charlie P SCUTT; Eva Sundberg; Christophe Trehin; Aurelie C Vialette-Guiraud

**Abstract:** The carpel is the female reproductive organ which encloses the ovules in the flowering plants, or angiosperms. The origin of the carpel and its subsequent morphological modifications were probably of vital importance to the evolution of the angiosperms, and the carpel is also very important as the precursor organ to the fruit. Here we describe the general attributes of the angiosperm carpel, and several hypotheses for its evolutionary origin. As carpels share many developmental processes with leaves, we describe these processes in the leaf, and then detail the regulation of carpel and fruit development in the model angiosperm *Arabidopsis thaliana*. We also describe the relationship between carpel formation and the arrest of organ proliferation which occurs at the centre of the *Arabidopsis* floral meristem. We then provide a brief overview of carpel development in angiosperms occupying important phylogenetic positions, including ANA grade angiosperms, monocots, basal eudicots and core eudicots, focussing on the probable ancestral state of the carpel in each case, and on the available molecular and genetic data. We end with a brief discussion of future research directions relating to carpel and fruit development.

## **Carpel Development**

Submitted by invitation to *Advances in Botanical Research*

### Authors

Cristina Ferrándiz<sup>1</sup>, Chloé Fourquin<sup>1</sup>, Nathanael Prunet<sup>2</sup>, Charlie P. Scutt<sup>2</sup>, Eva Sundberg<sup>3</sup>, Christophe Trehin<sup>2</sup> and Aurélie C.M. Vialette-Guiraud<sup>2</sup>

### Author addresses

1.

Instituto de Biología Molecular y Celular de Plantas,  
CSIC-Universidad Politécnica de Valencia,  
CPI Ed. 8E, Camino de Vera s/n. 46022 Valencia,  
Spain.

2.

Reproduction et Développement des Plantes,  
UMR 5667, CNRS-INRA-Université de Lyon,  
Ecole Normale Supérieure de Lyon,  
46, allée d'Italie,  
69364 Lyon Cedex 07,  
France.

3.

Uppsala BioCenter,  
Department of Plant Biology and Forest Genetics,  
Swedish University of Agricultural Sciences,  
PO Box 7080,  
SE-75007 Uppsala,  
Sweden.

### Manuscript information

Text pages (excluding references): 69

Word count (excluding references): 20 550

Figures: 11.

Please note that Figures 2A and 2B are supplied in separate files. All figures are in colour, except for Figure 2B and Figure 9.

### Contact for editorial purposes

E-mail: [Charlie.Scutt@ens-lyon.fr](mailto:Charlie.Scutt@ens-lyon.fr)

Tel +33 4 72 72 86 03

Fax +33 4 72 72 86 00

## **Abstract**

The carpel is the female reproductive organ which encloses the ovules in the flowering plants, or angiosperms. The origin of the carpel and its subsequent morphological modifications were probably of vital importance to the evolution of the angiosperms, and the carpel is also very important as the precursor organ to the fruit. Here we describe the general attributes of the angiosperm carpel, and several hypotheses for its evolutionary origin. As carpels share many developmental processes with leaves, we describe these processes in the leaf, and then detail the regulation of carpel and fruit development in the model angiosperm *Arabidopsis thaliana*. We also describe the relationship between carpel formation and the arrest of organ proliferation which occurs at the centre of the *Arabidopsis* floral meristem. We then provide a brief overview of carpel development in angiosperms occupying important phylogenetic positions, including ANA grade angiosperms, monocots, basal eudicots and core eudicots, focussing on the probable ancestral state of the carpel in each case, and on the available molecular and genetic data. We end with a brief discussion of future research directions relating to carpel and fruit development.

## **I. Introduction**

### **A. The carpel and gynoecium**

The carpel is the female reproductive organ that encloses the ovules in the flowering plants, or angiosperms. By contrast, ovules in the remainder of the seed plants, or gymnosperms, occur as naked structures, often borne in the axils of leaf-like organs such as the cone scales of conifers. Indeed, the terms “angiosperm” and “gymnosperm” describe this difference, as they respectively refer to enclosed and naked seeds (from the Greek *angeion* = vessel, *gymnos* = naked and *sperma* = seed). Carpels typically occur in the fourth and innermost whorl of the angiosperm flower, which is termed the gynoecium. These organs may occur separately, in which case the gynoecium is said to be apocarpic, or may be fused together into a gynoecium which is then termed syncarpic. Both individual carpels and syncarpic gynoecia are divided longitudinally into tissues which perform distinct roles in reproduction. Thus, the stigma at the apex of these structures is specialised for the capture and germination of pollen grains, and below this the style conducts pollen tubes to the ovary, which houses the ovules and in which fertilisation takes place.

### **B. The advantage of having carpels**

The carpel is thought to confer a number of major advantages on the flowering plants. Firstly, carpels protect the ovules within them, in part through the expression of genes associated with defence against insects and micro-organisms (Scutt et al., 2003). Secondly, systems have evolved to enable pollen capture and pollen tube guidance in carpel tissues, which may represent considerable improvements over equivalent mechanisms operating in gymnosperms. Thirdly, during the phase of pollen germination and growth, the carpel provides a site for the operation of self- and inter-specific incompatibility mechanisms: self-incompatibility prevents

close inbreeding and thereby conserves the capacity for evolutionary change, while inter-specific incompatibility prevents wide hybridisations that may lead to the production of unviable offspring. Fourthly, carpel tissues undergo developmental changes after fertilisation to form fruits, which protect the developing seeds within them and, at maturity, contribute to the dissemination of these.

### C. The added benefits of carpel fusion

More than 80% of angiosperm species are syncarpic: their carpels are fused into a single female structure in the centre of the flower, and this trait has probably arisen over 20 times independently in the angiosperms (Armbruster et al., 2002). Carpel fusion confers numerous advantages on syncarpic species (Armbruster et al., 2002, Endress, 1982), of which one of the most important is the provision of a compitum: a tissue which acts as an interchange between the entire stigmatic surface and the ovary, thus allowing any pollen tube to access any ovule. Another potentially important advantage of syncarpy results from the enhanced competition that this produces between pollen tubes: an effect which may select for vigorous male parents. Syncarpy also allows for the production of larger fruits, with potentially more complex and efficient seed dispersal mechanisms. Finally, a syncarpic gynoecium may require a lesser energy input for cell wall production, compared to an apocarpic gynoecium of similar size.

Syncarpy can be divided into two types based on the timing of the fusion event involved: where carpels are fused from the earliest emergence of their primordia, the fusion is termed “congenital”, whereas fusion that takes place during development is termed “post-genital”. Congenital carpel fusion is the most common type, with post-genital fusion occurring in only a few families (Lolle and Pruitt, 1999). The molecular basis for congenital carpel fusion has not been investigated in detail, though a large number of *Arabidopsis* mutants are known that

disrupt congenital carpel fusion (Vialette-Guiraud and Scutt, 2010). Postgenital carpel fusion has been studied most fully in *Catharanthus roseus* (Apocynaceae), in which two separate carpel primordia are initiated and then grow until their inner surfaces come into contact (Siegel and Verbeke, 1989, Verbeke, 1992, Walker, 1978). The already differentiated epidermal cells of these surfaces then begin to interlock and redifferentiate into parenchyma by a process which is dependent on diffusible, water-soluble agents produced by the carpels (Siegel and Verbeke, 1989). For all of the reason given above, both the origin of the carpel and the multiple origins of syncarpy were almost certainly major factors in the evolutionary success of the angiosperms. This group arose from an unknown common ancestor, believed to have lived in the Lower Cretaceous Period, to generate an estimated 300 000 or more species alive today.

## **II. The Origin of the Carpel**

### **A. An “abominable mystery”**

This is how Charles Darwin famously referred to the recent apparition and rapid diversification of the angiosperms (see Friedman, 2009). The rapidity of early angiosperm diversification, as suggested by the fossil record, was in contradiction with Darwin’s viewpoint as an evolutionary gradualist, and though this feature of angiosperm evolution may seem less mysterious to present day biologists, we still lack understanding of many aspects of the origin of the flowering plants. For example, little is currently known of the molecular changes that were responsible for the highly novel anatomical features of the first flowers, including the carpel. Additionally, we lack a clear picture of the non-flowering progenitor of the flowering plants, and knowledge of which fossil gymnosperms, such as *Corystospermales*, *Caytoniales*, *Glossopteridales*, *Bennettitales* or *Schmeissneria* (Taylor and Taylor, 2009, Wang et al., 2007), might be sister or ancestral to the angiosperms. Further unanswered

questions relate to where the flowering plants originated and to the date of this event: though a Lower Cretaceous origin is widely cited, the carpel and other key floral features may have originated earlier than that. Though much further work is thus necessary if we are to piece together the early evolution of the flower and carpel, the current lack of firm evidence has not prevented the construction of numerous hypotheses on this subject, as described below.

### **B. Hypotheses of carpel origin**

A hypothesis proposed by the writer and philosopher Goethe (von Goethe, 1790), and which is now well supported by molecular and genetic evidence (Honma and Goto, 2001), regards all plant lateral organs, including carpels, as variants of a basic leaf-type developmental ground plan. Though the carpel may thus be homologous to leaves, this floral reproductive organ is almost certainly more directly related to the leaf-like structures present in the reproductive axes of the angiosperms' sister group, the gymnosperms. On this subject, hypotheses for flower origin divide conceptually into two types, depending on whether they regard the carpel as derived by the modification of male or female structures in a presumed gymnosperm-like ancestor. The Mostly Male Theory (MMT; Frohlich, 2003, Frohlich and Parker, 2000) postulates the flower to be mostly derived from the male strobili of a gymnosperm-like ancestor. According to this hypothesis (Fig. 1A), the ancestor of the flowering plants would first have generated ectopic ovules on male sporophylls, which would thereby have become bisexual. The MMT postulates that ectopic ovules were concentrated on sporophylls near the apex of the strobilus, and that these sporophylls subsequently lost their ability to produce microsporangia, thus becoming functionally female. The MMT goes on to postulate these newly female sporophylls to have closed around the ovules to form the first carpels. In subsequent evolutionary steps, the residual female strobili of these proto-flowering



plants would have been lost, leaving only bisexual reproductive axes containing apical carpels and basal microsporophylls (later to become stamens).

The MMT is based on evidence from a number of sources, including data linked to *LEAFY* (*LFY*), which acts upstream of genes that specify the identities of floral organs in model angiosperms. In certain gymnosperms, a paralogue of *LFY* termed *NEEDLY* (*NLY*) has been shown to be expressed principally in female cones (Mouradov et al., 1998). No direct orthologue of *NLY* has been found in any angiosperm, suggesting this gene lineage to have been lost from a common ancestor of the living angiosperms, subsequent to the separation of the angiosperm and gymnosperm lineages. The MMT postulates the loss of *NLY* to have been accompanied by the extensive loss of female-specific developmental programmes, and this loss to have contributed to the origin of the flower. Hence, the MMT regards the carpel as derived from male reproductive organs. The MMT even accounts for the origin of the outer integument of the ovule, which is also specific to the flowering plants, by proposing this to have arisen from a cupule that surrounded the ovules in their presumed gymnosperm-like ancestor. Indeed, the MMT cites Jurassic fossil *Corystospermales* as having cupules of a type which could have evolved to generate the outer integument. Though the MMT has been a very widely discussed and conceptually useful hypothesis for flower origin, it should be noted that several more recent studies have questioned the sex-specific expression of *LFY* and *NLY* in gymnosperms on which the MMT is partly based (Carlsbecker et al., 2004, Dornelas and Rodriguez, 2005, Vazquez-Lobo et al., 2007).

Several further hypotheses of flower origin have been proposed, which differ from the MMT in that they postulate the bisexuality of the flower to have arisen by a spatial or temporal change in factors controlling the sex of reproductive organs. Hence, these hypotheses do not,

in contrast to the MMT, postulate the extensive loss of female developmental programmes during flowering plant evolution and consequently regard the carpel as homologous to female, rather than male, gymnosperm reproductive structures. The Out-of-Male (OOM) hypothesis (Theissen et al., 2002) proposes the bisexual flower to have evolved by the basipetal movement of the expression of a male-promoting, B-class MADS box gene in a previously male strobilus, resulting in the production of female structures at the apex (Fig. 1B). A sister hypothesis to the OOM hypothesis, termed the Out-of-Female (OOF) hypothesis (Theissen et al., 2002), postulates a sex-determining role for B-sister MADS box genes, whose expression is proposed to have moved acropetally in a female strobilus to leave male structures in basal positions (Fig. 1C). In general therefore, the OOM and OOF hypotheses focus below the level of *LFY* and *NLY* in the hierarchical control of gene expression, and postulate a spatial change in MADS box gene expression to form a boundary of B- or B-sister expression in a previously unisexual strobilus, thereby making this bisexual. It should be added that recent functional characterisation of B-sister genes in *Arabidopsis* has revealed roles in seed pigmentation (Nesi et al., 2002) and outer integument development (de Folter et al., 2006, Prasad et al., 2010), rather than in carpel development *per se*. However, it cannot be excluded that B-sister genes may have played a role in carpel development in early flowering plants.

Baum and Hileman (2006) have formulated a further hypothesis, which will be termed here the BHH, to account for the evolution of the first flowers (Fig. 1D). Like the MMT, the BHH proposes a central role for *LFY* in the origin of the flower, but postulates that the origin of floral bisexuality was caused, not by the loss of female-specific developmental programmes, but by a temporally generated switch in responses to *LFY*. According to this hypothesis, *LFY* protein levels increase with time in the meristems of developing reproductive axes and, at a certain threshold, cause these meristems to switch from the production of (male)

microsporophylls to (female) megasporophylls. This hypothesised switch may involve the action of *LFY* cofactors, such as *UNUSUAL FLORAL ORGANS (UFO)* and *WUSCHEL (WUS)*. Whatever the precise mechanism, the BHH proposes that a change occurred during early flower evolution in the relative response to *LFY* of B- and C-class MADS genes. Accordingly, C-class proteins are proposed to have predominated at the high *LFY* concentrations encountered at the apex of the strobilus at late developmental stages, resulting in MADS box complexes that were rich in C-class proteins. These proteins would have formed C-rich complexes which would consequentially have specified the development of megasporophylls at the apex of the strobilus.

The above hypotheses may, to some extent, be tested. Baum and Hileman (2006), for example, propose a list of predictions of their hypothesis that could be tested in basal angiosperms and gymnosperms. The MMT stands out from the other hypotheses described here in proposing the extensive loss of female developmental programmes during early flower evolution. This prediction might provide a means to eliminate either the MMT, or all other current contending hypotheses, from consideration. Thus, if the predictions of the MMT were correct, we would expect to find in gymnosperms numerous classes of genes with female-specific expression patterns, the orthologues of which had been lost from the angiosperms. Gymnosperm genes with male-specific expression patterns should not have been affected in this way. The testing of the MMT by this method has yet to be performed on a large scale. However, one question mark concerning such a test relates to the degree to which male and female developmental programmes in gymnosperms might be based on different sets of genes, rather than on subtle changes to the expression of a common set of genes. If the latter is predominately the case, such a relatively simple method of hypothesis testing may be unavailable.

### C. A possible role for the E-function in carpel evolution

The E-function MADS box genes may have played an important role in the origin of the flower and its carpel. In *Arabidopsis*, these genes encode the SEPALLATA1-4 (SEP1-4) transcription factors, which are hypothesised to act in quaternary complexes, together with combinations of A-, B- and C-function MADS box proteins, to specify organ identity in each whole of the flower (Honma and Goto, 2001, Pelaz et al., 2000, Theissen and Saedler, 2001). Theissen and Melzer (2007) discuss the possibility that, before the flower, dimers of C-function genes may have specified the development of female reproductive organs, and that the evolution of quaternary MADS box complexes, incorporating both C- and E-function proteins, may have built on this mechanism to generate the carpel. More precisely, the evolution of MADS box quaternary complexes is hypothesised to have caused transcription factor binding to two distinct MADS-box binding motifs, termed CArG boxes, in the cis-acting control regions of their target genes. According to this hypothesis, this newly evolved DNA binding behaviour would have generated the necessary multiplicity of interactions to specify at least three novel organ types in early flowers: carpels, stamens and tepals.

*SEP* genes, encoding the E-function, appear to be specific to the angiosperms (Becker and Theissen, 2003). However, recent evidence suggests the E-function may not be exclusively associated with the *SEP* clade: genes of the *AGAMOUS-LIKE6* (*AGL6*) clade have recently been demonstrated to contribute to the E-function in both *Petunia* (Rijkema et al., 2009) and Poaceae monocots (Li et al., 2010, Ohmori et al., 2009, Thompson et al., 2009). The *AGL6* clade is sister to the *SEP* clade in angiosperms, and is also present in gymnosperms (Becker et al., 2000), thus leaving open the possibility that quaternary MADS box complexes involving *AGL6* proteins might also form in gymnosperms. The presence of *AGL6* orthologues in

gymnosperms tends to suggest the loss of the *SEP* clade from this group, rather than its specific origin by duplication in the angiosperms, also casting some doubt on the idea that the genes responsible for the E-function arose specifically with the angiosperms. To resolve the question of the potential contribution of the E-function to the origin of the flower, and therefore of the carpel, attention is now being paid to the formation of higher order complexes of MADS box proteins in both angiosperms and gymnosperms (Melzer et al., 2010).

### **III. The Basic Developmental Plan of Lateral Organs**

#### **A. Leaves and carpels share basic regulatory pathways**

As mentioned in the previous section, leaves and floral organs most likely have a common evolutionary origin, or as Goethe memorably put it: *Alles ist Blatt* (All is Leaf; von Goethe, 1790). An increasing amount of evidence suggests that the pathways regulating the basic morphological outline of ancestral leaf-like organs have been recruited to the developmental programs of both leaves and floral organs, including the carpels, of present day plants, although in the latter case, these have been slightly modified and positioned downstream of genes determining floral organ identity. For instance, mutations in genes affecting carpel morphogenesis also result in defects in the basic morphogenesis of the leaves.

We will start by summarising what is known about the formation and patterning of leaves, in order to be able to use this information as a basis for a discussion of carpel and fruit morphogenesis in later sections. Impressive progress in our understanding of the genetic regulation of leaf initiation and morphogenesis has been made during the last decade, and this has been the subject recently of several in-depth reviews (Aida and Tasaka, 2006, Barkoulas et al., 2007, Bowman and Floyd, 2008, Byrne, 2006, Husbands et al., 2009, Kepinski, 2006, Pulido and Laufs, 2010, Shani et al., 2006), though in contrast to these, we aim only to give

an overview of the known regulatory networks.

### **B. Initiation of lateral organs at the flanks of the shoot apical meristem (SAM)**

In this section we will approach the following two basic questions: what regulatory events are required for lateral organ initiation at the peripheral zone (PZ) of the shoot apical meristem (SAM), and what are the events leading to the formation of lateral organ primordia? Before the initiation of lateral organs, the cells of the SAM are maintained in an undifferentiated state by the activity of a number of regulators, including members of the class I KNOTTED1-like homeobox (KNOX) and WUSHEL-like homeobox (WOX) transcription factor families (Fig. 2A; Hake et al., 2004, Laux et al., 1996). Cell expansion is strongly correlated to differentiation, and the above-mentioned factors act in part by promoting a high ratio of cell division to cell expansion through the modulation of hormonal balances in the SAM (Jasinski et al., 2005, Yanai et al., 2005). Cell division is stimulated by cytokinin-induced activation of Cyclin D, and high cytokinin levels in the *Arabidopsis* SAM result, at least in part, from KNOX-induced activation of the cytokinin biosynthesis gene *ISOPENTENYL TRANSFERASE7 (IPT7)*; Yanai et al., 2005). Not only cytokinin synthesis (Kurakawa et al., 2007), but also cytokinin responsiveness, is high in the central part in the meristem, and at least two members of the WOX family, WUS and STIMPY/WOX9, appear to stimulate cytokinin signalling in the SAM (Fig. 2A; Gordon et al., 2009, Skylar et al., 2010). Interestingly, recent data suggest that cytokinin stimulates *KNOX* and *WUS/WOX* activity, suggesting the presence of a positive feedback loop (Gordon et al., 2009, Kurakawa et al., 2007). At the same time, cell expansion is repressed by keeping the level of the hormone gibberellin low through the KNOX-mediated repression of the gibberellin biosynthesis gene *GA20-OXIDASE (ga20ox)* and by activation of the GA catabolism gene *ga2ox1* (Bolduc and Hake, 2009, Chen et al., 2004, Hay et al., 2002).

Lateral organ initiation in the PZ of the SAM requires the silencing of programmes that repress differentiation, such as the *KNOX* programme, and it has been suggested that the plant hormone auxin plays a major role in this process. Local auxin concentration maxima are formed at organ initiation sites by a directed auxin flux, which results from the action of auxin influx and efflux facilitators (Bainbridge et al., 2008, Heisler et al., 2005, Reinhardt et al., 2000, Reinhardt et al., 2003). In these “high auxin” organ initiation sites, *KNOX* gene activity is repressed by auxin action itself, as well as by the activity of a transcriptional repressor complex containing the *ASYMMETRIC LEAVES1* (*AS1*) and *AS2* proteins, resulting in the induction of leaf formation (Fig. 2A; Guo et al., 2008, Hay et al., 2006, Ikezaki et al., 2010). The *AS1/AS2* complex also maintains the silencing of *KNOX* throughout leaf development via the recruitment of the *HIRA* chromatin-remodelling factor to the *KNOX* gene (Guo et al., 2008, Phelps-Durr et al., 2005). Leaf initiation appears also to be regulated by a *SQUAMOSA PROMOTER BINDING PROTEIN-LIKE 9* (*SPL9*)-dependent, leaf-derived signal that may act via the modulation of auxin pathways (Wang et al., 2008).

### C. Establishment of boundaries

When a leaf primordium starts to grow, a boundary domain, in which cell expansion is reduced, becomes established to separate it from the neighbouring tissues. So the next question is: what regulatory pathways position and establish boundaries between the SAM and newly initiated lateral organs? It has been suggested that a signal from the CZ of the meristem participates in the positioning of organ boundaries, and that activities in the meristem and the each new organ together establish the molecular changes required for boundary formation (Rast and Simon, 2008). As soon as one organ primordium has been initiated, a redirection of auxin flux to a new position in the PZ establishes the initiation of the

next lateral organ (Heisler et al., 2005). As a consequence, auxin is depleted and changes to gene expression occur in the cell layers surrounding the newly formed primordium, which creates a morphological boundary of distinct cell types with reduced cell division activity that separates the primordium from the rest of the meristem (Fig. 2A; Heisler et al., 2005), as reviewed by Aida and Tasaka (2006). The NAC domain transcription factors CUP-SHAPED COTYLEDON (CUC1), CUC2 and CUC3 are expressed in the boundary domain, where their activity contributes to the repression of cell division and expansion (Hibara et al., 2006, Sieber et al., 2007, Vroemen et al., 2003). The expression of the growth inhibiting *CUC* genes is restricted by members of the *miRNA164* family, indicating *CUC* genes to be central regulators of boundary size (Laufs et al., 2004, Sieber et al., 2007). The JAGGED LATERAL ORGANS (JGL) LBD domain protein is also expressed at the SAM/organ boundary, where it promotes the boundary function by repressing *PINFORMED* (*PIN*) activity, potentially resulting in low auxin concentrations at the boundary, and by activating *KNOX* genes (Borghi et al., 2007, Husbands et al., 2007, Shuai et al., 2002). Recently, the BELL-type protein *ARABIDOPSIS THALIANA* HOMEODOMAIN BOX GENE1 (*ATH1*) was also suggested to participate in boundary formation in a pathway parallel to that of the *CUC* proteins (Gomez-Mena and Sablowski, 2008).

#### **D. Regulation of adaxial-abaxial polarity**

Concomitant with the lateral outgrowth of the leaf primordium, a distinct polarity along the adaxial-abaxial axis is established. Anatomical features that optimise the leaf for photosynthesis are formed on the upper or adaxial side, adjacent to the SAM, whereas the lower or abaxial side differentiates to carry out gas exchange (for recent reviews see Chitwood et al., 2007, Husbands et al., 2009 and Xu et al., 2007). Members of the class III homeodomain-leucine zipper (HD-ZIP III) gene family, such as *PHABULOSA* (*PHB*),



*PHAVOLUTA* (*PHV*) and *REVOLUTA* (*REV*), are expressed on the adaxial side of leaf primordia, where they play a major role in adaxial tissue specification (Fig. 2A; Emery et al., 2003, McConnell et al., 2001, McConnell and Barton, 1998, Prigge et al., 2005). HD-ZIPIII genes are also expressed in the CZ of the meristem, and have been suggested to coordinate communication between the SAM and the adaxial side of organ primordia (McConnell et al., 2001). By contrast, *miRNA166* accumulates on the abaxial side of leaf primordia, where it represses the activity of HD-ZIPIII genes via posttranscriptional cleavage and/or chromatin modifications (Alvarez et al., 2006, Bao et al., 2004, Emery et al., 2003, Kidner and Martienssen, 2004, Mallory et al., 2004, Tang et al., 2003, Williams et al., 2005, Zhou et al., 2007). The AS1 and AS2 proteins also support adaxial fate (Fu et al., 2007, Lin et al., 2003, Xu et al., 2003). The abaxial side of leaves is specified by another set of regulators, acting antagonistically to the adaxial determinants, suggesting the mutually exclusive and opposing nature of adaxial and abaxial cell fates (Fig. 2A). Transcription factors belonging to the KANADI (KAN) subgroup of the GARP family and the auxin response factors AUXIN RESPONSE FACTOR3/ETTIN (ARF3/ETT) and ARF4 together specify abaxial identity (Eshed et al., 2001, Kerstetter et al., 2001, Pekker et al., 2005). It has recently been shown that KAN1 directly represses AS2 transcription on the adaxial side of leaves, and other data suggest that KAN genes also act as negative regulators of HD-ZIPIII gene expression (Wu et al., 2008, Kerstetter et al., 2001, Eshed et al., 2004). The activity of *ARF3* and *ARF4* is restricted to the abaxial side of the leaf primordium by the action of *TAS3* encoded ta-siRNAs that accumulate on the adaxial side (Adenot et al., 2006, Fahlgren et al., 2006, Garcia et al., 2006, Hunter et al., 2006, Nogueira et al., 2007, Vazquez et al., 2004, Williams et al., 2005). Further data suggest the genes encoding YABBY (YAB) transcription factors to act downstream of other polarity determinants, including the KANs and the ARFs, to direct leaf lamina expansion at the adaxial-abaxial boundary (Sawa et al., 1999, Siegfried et al., 1999,

Eshed et al., 2001, Eshed et al., 2004). The antagonistic activity of the KAN and HD-ZIPIII genes has also been recruited to establish polarity during embryogenesis and vasculature formation, suggesting these to be important general regulators of polarity (Emery et al., 2003, Eshed et al., 2001, Ilegems et al., 2010, Izhaki and Bowman, 2007).

One important question that remains to be answered concerns how polarity information is generated to induce the expression of fate-specific regulators. It has been suggested that the establishment of the adaxial-abaxial axis of leaves is dependent on the conversion of positional signals provided by the SAM, and probably also by other surrounding areas, into the differential expression of the mutually antagonistic transcription factors mentioned above (Fig. 2A). Because the separation of the incipient leaf primordium from the CZ of the SAM through microsurgical sections results in radial abaxialised structures, it appears likely that adaxial identity is specified by a meristem-derived signal, continuously entering the primordium (Reinhardt et al., 2005, Sussex, 1954). However, the nature of this signal is still not known. Because the START domain of HD-ZIPIII proteins appears capable of lipid/sterol binding, this was suggested as the potential target of an unknown SAM-derived lipid/sterol signal (McConnell et al., 2001). Furthermore, as ta-siRNA-ARFs, encoded by *TAS3*, can move between cells, it has been suggested that these may play a role in SAM-to-primordia signalling (Garcia et al., 2006). However, although the ta-siRNA-ARFs can move from below the SAM into the meristem proper, as well as from the adaxial to the abaxial side of the leaf blade, there is no evidence that these molecules act as messengers from the SAM to leaf primordia (Chitwood et al., 2009). Instead, Chitwood et al. (2009) suggest that a gradient of these small RNAs is formed, which could define the expression boundary of their targets *ARF3* and *ARF4*. A signalling molecule often used for positional information is auxin, and Pekker et al. (2005) have suggested that auxin could act as an abaxially polarizing signal,

activating *ARF3* and *ARF4*. This is supported by the finding that the auxin influx facilitator *AUX1* localises specifically in the abaxial epidermal layer, suggesting that auxin may flow into the abaxial half of the incipient primordium and that an auxin gradient may be established across the primordium (Reinhardt et al., 2003).

#### **E. Leaf development**

Leaf development proceeds through various different steps during which cell proliferation, cell expansion and cell differentiation occur. One major question is: how is this process coordinated? Auxin gradients may contribute to leaf development by coordinating growth, and, for example, the differentiation and patterning of veins. An auxin maximum at the apical tip of the leaf primordium is established through auxin transport early in development (Reinhardt et al., 2003), and is maintained by the induction of auxin biosynthesis at the tip, and later on also in the hydathodes at the margins of the leaf primordium. It has been suggested that this process allows the formation of a distal-proximal auxin gradient (Benkova et al., 2003), which is important for controlled cell division and expansion, and gradients formed by auxin transport from the leaf tip have been suggested to be important for midvein development (Mattsson et al., 1999, Zgurski et al., 2005).

The mechanisms that establish the pattern of differentiation along the distal-proximal axis of the leaf have not yet been determined. It has been shown, however, that cell proliferation and differentiation/expansion occurs along a gradient from the (distal) leaf blade toward the (proximal) petiole, reflected by the gradual move of a front of cell cycle arrest from the tip to the base (Fig. 2B; Donnelly et al., 1999, Nath et al., 2003). Similarly, cell divisions in the mid-region decline slightly ahead of divisions at the leaf margins in a medio-lateral gradient (Byrne, 2005). Thus, it is quite clear that the control of cell division is an integral part of

pattern formation, and most likely contributes to the multitude of leaf shapes found in nature. Transcriptional regulators of cell division include: JAGGED (JAG), AINTEGUMENTA (ANT), GRF-INTERACTING FACTORS (GIFS), GROWTH-REGULATING-FACTORS (GRF), PEAPOD (PPD), LEAFY PETIOLE (LEP) and class I TCP (TEOSINTE BRANCHED1, CYCLOIDEA, PCF), all of which promote cell divisions in the leaf (Dinneny et al., 2006, Horiguchi et al., 2005, Hu et al., 2003, Lee et al., 2009, Li et al., 2005, Mizukami and Fischer, 2000, van der Graaff et al., 2000, White, 2006), as reviewed by Anastasiou and Lenhard (2007) and Ingram and Waites (2006). In addition, the final expression patterns of some of these regulators, such as the GRF and TCP genes, appear to be controlled by miRNAs (Palatnik et al., 2003, Palatnik et al., 2007, Rodriguez et al., 2010). By contrast, growth cessation is promoted by the bHLH protein SPATULA (SPT) and the class II TCP proteins, starting from the tip of the leaf and continuing to the base (Efroni et al., 2008, Ichihashi et al., 2010, Nath et al., 2003, Ori et al., 2007, Palatnik et al., 2003). The distal-proximal polarity of the leaf also results in an asymmetry in the size of the leaf blade. The leaf is wide in its distal region, but at the proximal end, a petiole is formed which has a very narrow blade (Fig. 2B). Two BTB/POZ transcriptional co-activators, BLADE-ON-PETIOLE1 (BOP1) and BOP2, repress cell proliferation and growth of the petiole. These regulators are expressed at the base of the developing leaf, where they directly activate *AS2* transcription, establishing the conditions for the repression of *KNOX* gene expression that is necessary to correctly pattern the petiole tissues that form at the proximal end of the leaf (Ha et al., 2003, Ha et al., 2004, Ha et al., 2007, Hepworth et al., 2005, Jun et al., 2010). Furthermore, the *BOP* genes also negatively affect cell proliferation in the proximal end of the leaf by repressing *JAG* and *NUBBIN* (*NUB*; Norberg et al., 2005).

## **F. Leaf margins**

Controlled cell proliferation and elongation are also required for shaping the leaf margins (Dinneny et al., 2004, Palatnik et al., 2003) and interestingly, auxin appears to play an important role in establishing a part of this growth pattern. Auxin responsiveness is evenly distributed in the leaf margin of simple leaves that lack serrations (ie. teeth; Aloni et al., 2003), whereas in simple leaf primordia, which will eventually form serrations at their margins, high auxin responsiveness can be detected in one or two marginal epidermal cells before teeth are recognizable, suggesting that auxin localization play an important role in patterning leaf margins (Kawamura et al., 2010). Furthermore, the depth of the indentation between the teeth of serrated leaves is regulated by the meristem and boundary gene *CUC2* (Nikovics et al., 2006), and *KNOX* meristem genes have also recently been implicated in leaf serration (Kawamura et al., 2010). The process of leaf serration shows many similarities to that of leaflet formation in dissected leaves. For example, leaf dissection in *Cardamine hirsuta* also relies on the activity of *CUC* boundary-specifying genes in the area delimiting leaflet primordia, and on the activity of *KNOX* genes at the leaflet initiation position, which cause a delay in cell differentiation relative to cell proliferation. This is followed by the production of auxin maxima at the leaflet tip (Canales et al., 2010, Hay et al., 2006). Interestingly, these modulations of leaf margins clearly provide evidence that processes related to meristem function are also used during lateral organ development.

#### **IV. The Basic Concepts of the *Arabidopsis* Gynoecium and Fruit**

##### **A. The *Arabidopsis* gynoecium**

As described in Sections I and II, above, the *Arabidopsis* gynoecium is a complex syncarpic structure, most commonly interpreted as being composed of two congenitally fused carpels. The *Arabidopsis* gynoecium is connected to the base of the flower by a short internode called the gynophore (Fig. 3). Above the gynophore is the ovary which contains between 50 and 80

ovules, and makes up most of the length of the gynoecium. The ovary is divided in two longitudinally by a septum which is formed postgenitally. The two ovary wall regions in the gynoecium are termed the valves, and the external part of the septum is termed the replum. At the apical end of the ovary are the style and the stigma. The stigma consists of a single layer of specialised epidermal cells bearing elongated processes termed stigmatic papillae. This surface receives pollen grains and permits their germination. Following this, pollen tubes are guided by the transmitting tract present in the style and septum towards the ovules. After fertilization, the ovules develop into seeds and the *Arabidopsis* gynoecium is transformed into a two-chambered fruit called a silique. This structure opens at maturity to release its seeds along four dehiscence zones which consist of thin regions present at the valve margins on either side of the replum. The lignification of specific cells in these zones contributes to the dehiscence process by providing mechanical tension which stimulates the detachment of the two valves (Balanza et al., 2006).

The gynoecial primordium arises in the centre of the floral meristem as a ring of cells enclosing a small depression, and then develops as an open-ended tube (Stages 6 to 8 of flower development, according to Smyth et al., 1990). Two opposing meristematic ridges form in the internal medial regions of this cylinder and fuse together to form the septum. Placental tissues, which will give rise to the ovules, develop in the zones where the vertical septum and the gynoecial walls meet. At Stage 9, valve, placenta, septum, style and stigmatic cells begin to differentiate. At Stages 11-12, the apical part of the gynoecium closes, the stigmatic papillae complete their development, and the style becomes distinct from the ovary. The gynoecium is mature at anthesis (Stage 13), when the flower opens and fertilization can take place. All the tissues required for fruit maturation and dehiscence are already present at

this stage, and will complete their development after the fertilisation of the ovules (Bowman et al., 1999, Roeder and Yanofsky, 2005).

### **B. From leaf to carpel: the identity genes**

Two decades ago, the ABC model was formulated to explain the genetic interactions which lead to floral organ identity (Coen and Meyerowitz, 1991). Since then, extensive research has been carried out that has broadly validated the ABC model and provided evidence for a biochemical model for the action of the A-, B- and C-functions, termed the “floral quartet model” (Krizek and Fletcher, 2005, Theissen and Melzer, 2007). According to this model, a leaf, which corresponds to the "ground state" for lateral organs, can be transformed into a carpel by expressing the C-function MADS-box gene *AG*, and at least one of three E-function MADS-box genes, *SEP1-3* (Honma and Goto, 2001). On the contrary, loss of function mutations in *AG* result in homeotic conversions of the carpel into a reiteration of the sequence sepals-petals-petals and the simultaneous loss of function of the redundant *SEP* genes results in a complete loss of carpel development programmes, transforming carpels into leaves (Ditta et al., 2004, Pelaz et al., 2000).

Despite its central role in specifying carpel identity, *AG* is not unique in providing carpelloid features. In fact, in the double mutant *apetala2 (ap2) ag*, organs with carpel characteristics still develop in the first whorl of the flower (Bowman, 1991). This observation led to the conclusion that other genes involved in carpel identity were present and were, like *AG*, negatively regulated by the A-class gene *AP2*. These factors have been identified as two other highly related and entirely redundant MADS-box genes *SHATTERPROOF1 (SHP1)* and *SHP2*, which are principally involved in the specification of valve margin identity (Liljegren et al., 2000; see Section IV.D, below). In the quadruple *ap2 ag shp1 shp2* mutant, all

carpelloid structures disappear. Other complementary studies have demonstrated that the AG and SHP proteins are extremely similar at a functional level, but play distinct roles during carpel development, mostly due to their different expression patterns (Pinyopich et al., 2001).

Two other putative transcription factors required for the development of carpel tissues are encoded by the bHLH gene *SPT* and the YAB gene *CRABS CLAW (CRC)*, which seem to act downstream of AG/SHP. As in the *ap2 ag shp1 shp2* quadruple mutant, the loss of *SPT* or *CRC* function in an *ap2 ag* background results in the loss of all carpelloid features (Alvarez and Smyth, 1999), showing that these two genes are also necessary for carpel development. *SPT* is widely expressed in both vegetative and reproductive structures throughout development (Heisler et al., 2001), although the main phenotypes associated with *spt* mutations are developmental defects in most of the marginal tissues of the carpel (Alvarez and Smyth, 1999). *CRC* is specifically expressed in nectaries and carpels; *crc* gynoecia are shorter and wider than wild type and partially unfused at the apex (Bowman and Smyth, 1999). In the *crc spt* double mutant, the gynoecium is completely unfused and possesses a considerably reduced number of ovules, in addition to much less stigmatic and stylar tissue (Alvarez and Smyth, 1999). Supporting the idea that *CRC* and *SPT* mediate the AG/SHP carpel identity function, Alvarez and Smyth (1999) showed that *CRC* and *SPT* were also involved in other aspects of AG activity, such as the termination of the floral meristem. Recently, it has been shown that *CRC* acts in combination with three other genes, *REBELOTE (RBL)*, *SQUINT (SQN)* and *ULTRAPETALA1 (ULT1)*, to control meristem determinacy (Prunet et al., 2008). Interestingly, *DROOPING LEAF (DL)*, the *CRC* ortholog from rice (Yamaguchi et al., 2004) and *EcCRC*, the *CRC* ortholog from poppy (Orashakova et al., 2009), have both been shown to play a role in the termination of the floral meristem, as



discussed further in Section V, below. Moreover *DL*, like *AG*, plays a prominent role in the C-function.

What do we know about the chronology and the hierarchy of the molecular events leading to carpel identity? *AG* and *SEP* genes are expressed in carpel primordia, even before any morphological sign of differentiation can be observed (Hempel et al., 1997, Savidge et al., 1995, Yanofsky et al., 1990). When the gynoecial primordium begins to form and develop as a cylinder, the expression of *AG* is uniform throughout this structure. Later however, *AG* is only expressed in specific cell types within the gynoecium, including the stigmatic papillae and ovules (Bowman et al., 1991a). In the very early stages of floral meristem development, the expression of *AG* is activated by the joint action of the floral identity regulator *LFY* and the meristem maintenance factor *WUS*. Once present, the *AG* protein acts to down-regulate *WUS*, leading to a loss of floral meristem activity (Lenhard et al., 2001, Lohmann et al., 2001). It has been shown that the A-function gene *AP2* inhibits *AG* expression in the perianth whorls of the flower (Drews et al., 1991). In *35S::SEP3* plants, *AG* is ectopically expressed, suggesting that *SEP3* could participate in the early activation of *AG* (Castillejo et al., 2005). Numerous studies have permitted the identification of many other factors involved in the regulation of *AG* activity, both at the transcriptional and post-transcriptional levels (Bao et al., 2004, Chen et al., 1999, Cheng et al., 2003, Das et al., 2009, Franks et al., 2002, Gregis et al., 2006, Krizek et al., 2000, Liu and Meyerowitz, 1995, Sieburth and Meyerowitz, 1997, Yu et al., 2009; see Section V for further details). In early stage of development, *SHP1* and *SHP2* are widely expressed in the *Arabidopsis* gynoecium. Later their expression is restricted to the valve margins, the top of the gynoecium, the placental tissue and the ovules (Flanagan et al., 1996, Savidge et al., 1995). *SHP* genes seem to act downstream of *AG* and could therefore represent direct *AG*-targets, though as demonstrated by *ap2 ag* mutants, these genes can also

be activated by other factors. Recently, it has been demonstrated that the transcription factors FILAMENTOUS FLOWER (FIL), YAB3 and JAG, also involved in leaf development, jointly activate *SHP* expression in valve margins (Dinneny et al., 2005; see Section IV.D for further details). Thus, both *SHP* and *AG* can be placed at the top of the carpel identity pathway, and could from this position directly or indirectly activate both *SPT* and *CRC*. *CRC* has been identified as a direct target of *AG* (Gomez-Mena et al., 2005), though less is known about *SPT* activation.

### C. Partitioning the carpel: adaxial-abaxial and medio-lateral patterning

Once organ identity has been specified, the gynoecial primordium is divided into different domains. In the first stages of development, abaxial-adaxial and medio-lateral patterning are specified, and later, as the primordium forms a cylinder, apical/basal polarity is defined. Abaxial-adaxial polarity refers to the differentiation between the outer (abaxial) and inner (adaxial) domains of the carpel (Fig. 3). These domains are respectively equivalent to the lower and upper sides of leaves, and the establishment of adaxial-abaxial polarity involves similar genetic mechanisms to those operating in leaves. Thus, the antagonistic interactions found in leaves between HD-ZIPIII genes, which direct adaxial fate, and KAN/YAB genes, which direct abaxial fate, also exist in the gynoecium (cf. Figs 4A and B). Indeed, HD-ZIPIII genes are expressed, as would be expected, in the adaxial domain of the carpel. However, inactivation of these genes seems to have a milder effect on carpel than on leaf development, indicating that additional factors involved in promoting adaxial polarity might operate in the gynoecium (Dinneny et al., 2006, McConnell and Barton, 1998). Recently, *JAG* and *NUB* have been identified as possible actors in carpel adaxialisation (Dinneny et al., 2006). *NUB*, unlike *JAG*, which is expressed in a non-polar manner in all lateral organs, is expressed only in the adaxial zones of leaves, stamens and carpels. Single

*nub* mutants do not present any obvious phenotype, but in the double *jag nub* mutant, floral organ growth is affected and carpels and stamens are abaxialised. In the abaxial domains of the carpel, as during leaf formation, the KAN genes, the auxin response factors *ETT* and *ARF4*, and the YAB genes are all expressed from very early stages of development (Kerstetter et al., 2001, Pekker et al., 2005, Siegfried et al., 1999). These genes are largely redundant in their abaxial activity, as single mutants of most of them only show a very weak adaxialisation phenotype. However, the combination of mutants can lead to severe polarity defects in which the gynoecium develops inside out. Thus, in *crc kan1*, *kan1 kan2* or *ett arf4* double mutants, the transmitting tract and ovules form on the exterior of the gynoecium (Eshed et al., 1999, Pekker et al., 2005).

Medio-lateral polarity is also specified very early during gynoecium development. The lateral domains of the gynoecium will give rise to the valves, while its medial domains correspond to the fused carpel margins and will develop internally into the placenta, septum, apical style and stigma, and externally into the replum (Fig. 3). Along the medio-lateral axis, two opposite types of tissue develop: in medial zones a new meristem called the medial ridge forms, while lateral domains develop as differentiated tissues. Accordingly, several genes involved in SAM maintenance, such as the class I KNOX factors and the “boundary genes” *CUC1* and *CUC2* are specifically expressed in the medial region, where they appear to be required for marginal tissue development. Ishida et al. (2000) analysed the gynoecium of *cuc1 cuc2* plants produced from calli, and therefore able to flower (grown from seed, these mutants would not progress beyond the seedling state). Interestingly, these plants showed defects in marginal tissue development and failed to develop a septum. Scofield et al. (2007) used inducible RNAi lines to study the effect of reduced activity of the KNOX I factor SHOOT MERISTEMLESS (STM) on gynoecium development. In some cases, the floral meristem aborted before forming

any carpels, whereas in others a gynoecium was present, but this lacked a septum and showed reduced marginal tissue development, thus forming a completely unfused structure. Conversely, the expression of genes that repress the undifferentiated state, and thus promote the development of lateral primordia, is restricted to the lateral domains of the gynoecium. Thus, YAB genes, *JAG/NUB* and *ASI/2* are specifically expressed in lateral regions of the young gynoecium (Alonso-Cantabrana et al., 2007, Bowman and Smyth, 1999, Dinneny et al., 2006, Siegfried et al., 1999). The genetic networks that maintain SAM and lateral primordia boundaries appear to perform similar functions in medio-lateral patterning of the gynoecium (Alonso-Cantabrana et al., 2007, Dinneny et al., 2005, Ragni et al., 2008). Dinneny et al. (2005) demonstrated *JAG*, *FIL* and *YAB3* to play a pivotal role in valve development in the gynoecium through the differential activation of valve and valve margin factors in the corresponding regions (Fig. 5). Alonso-Cantabrana et al. (2007) showed that later on in the development of the *as1* gynoecium, or in plants over-expressing *BP*, the replum is expanded and the valves are narrower. These authors concluded that *AS1* plays a specific role in promoting valve initiation, and this action is likely to involve the repression of *KNOX I* and other factors directing replum development. A model was proposed in which both a gradient of *FIL/YAB3/JAG* with a lateral maximum, and *AS1*, are present in the valve, while a *KNOX/RPL* gradient with a medial maximum is present in the replum. According to this model, the two gradients would have opposite actions and would thus define valve margin development in their region of overlap.

Another factor which plays an important role in the patterning of the gynoecium is the phytohormone auxin. It has been proposed that a gradient of auxin established during carpel development controls tissue patterning along the apical-basal axis (Nemhauser et al., 2000). From severe defects in apical-basal tissue distribution in *ett* mutants, the auxin response factor

*ETT* has been proposed to be the principal integrator of auxin gradient information during apical-basal development in the ovary (Nemhauser et al., 2000). In the same way, it is possible that auxin gradients also play important roles in the establishment of abaxial-adaxial and medio-lateral polarity. We mentioned above that *ETT* and *ARF4* were involved in specifying abaxial fate, as seen from the “inside-out” gynoecium phenotype of the *ett arf4* double mutant. In addition, the inactivation of *YUCCA* (*YUC*) genes, which encode enzymes of the auxin biosynthesis pathway, or *PINOID* (*PID*), which encodes a regulator of auxin transport, transforms the gynoecium into a completely radial structure (Bennett et al., 1995, Cheng et al., 2006). Moreover, in plants in which polar auxin transport is altered, the replum is expanded and the lateral valves are reduced (Bennett et al., 1995, Nemhauser et al., 2000). Very recently, it has also been demonstrated that a maximum of auxin is present in the replum, whereas a low level of this hormone is present in the valves (Sorefan et al., 2009). Therefore, it seems that auxin, beyond its central role in the establishment of apical-basal polarity in the gynoecium, is also involved in patterning its abaxial-adaxial and medio-lateral axes.

#### **D. Development of the carpel lateral domains**

The lateral domains of the gynoecial primordium give rise to the valves, which form the ovary wall, and the valve margins, which form at the valve/replum borders (Fig. 3). The valve margins later differentiate into the dehiscence zones, where the *Arabidopsis* fruit opens. *AG* has been proposed to specify valve identity, based on the phenotype of the *ap2 ag* double mutant, in which the carpeloid organs present in the first whorl of the flower do not show typical valve cell organization (Bowman, 1991). Valve margin specification and dehiscence zone formation have been extensively studied over the last few years, and the major factors involved in this process (Fig. 5) have now been identified, as reviewed by Dinneny and

Yanofsky (2005), Ferrandiz (2002) and Girin et al. (2009). Unlike *AG*, the closely related *SHP* genes direct valve margin identity. In the *shp1shp2* double mutant, the dehiscence zone fails to differentiate and the mature fruit does not open (Liljegren et al., 2000). Similar phenotypes are observed in plants in which the *INDEHISCENT (IND)* or *ALCATRAZ (ALC)* genes, encoding bHLH transcription factors, have been inactivated (Liljegren et al., 2004, Rajani and Sundaresan, 2001). *SHP* expression is restricted to the valve margins by the actions of the MADS-box gene *FRUITFULL (FUL)* in the valve, and of the homeodomain factor *RPL* in the replum (Ferrandiz et al., 2000b, Gu et al., 1998, Roeder et al., 2003). Then, in a narrow domain between these tissues, which is composed of 3-4 cell layers, *SHP* activates *IND* and *ALC*, which are necessary for the formation of the dehiscence zone (Liljegren et al., 2000, Liljegren et al., 2004, Rajani and Sundaresan, 2001). In *ful* mutants, *SHP*, *IND* and *ALC* are ectopically expressed in the valves and, as a result, small lignified cells, which are normally specific to the dehiscence zones, develop in these tissues. Consequently, *ful* fruits do not elongate, but instead break prematurely. In *rpl* mutants, the replum is reduced in width and the valve margins are expanded, as is expression of the corresponding valve margin genes. Based on genetic analyses, Dinneny et al. (2005) proposed a model in which the cooperative activity of *FIL*, *YAB3* and *JAG* would activate the transcription of *FUL* and *SHP* genes in the valves, and in the valve margins, respectively. According to this model, a high level of *FIL/YAB/JAG* activity would turn on *FUL* expression in the valves, while the activation of *SHP* in the valve margins would require a weaker activity of this same module. This model fits nicely with the observed phenotypes, but whether the activation of *FUL* and *SHP* lies in differences in *YAB/JAG* levels in different domains, or in some other type of molecular interaction, remains to be seen. In addition, it was shown in the replum that the homeodomain protein *RPL* represses the activity of the *FIL/YAB/JAG* module, therefore preventing *SHP* activation.

Auxin is also involved in the development of the lateral domains of the carpel. Sorefan et al. (2009) recently showed that a local auxin minimum is required for the differentiation of the dehiscence zones in the *Arabidopsis* fruit. These authors demonstrated that IND is involved in creating this auxin minimum by controlling the direction of auxin transport via PIN relocation. *SPT*, which has been proposed to mediate auxin signaling in apical-basal development, is also expressed in the dehiscence zones of the developing fruit (Heisler et al., 2001), where it appears to be regulated by IND (Groszmann et al., 2008). Though *SPT* expression suggests this factor to play a role in the dehiscence zones, no such function has yet been discovered, raising the possibility of redundancy with other factors.

#### **E. Development of carpel marginal tissues**

The carpel marginal tissues derive from the medial region of the gynoecial primordium, localised at the boundary between the two fused carpels (Fig. 3). A meristematic medial ridge of tissues develops along the adaxial side of the gynoecial tube and this gives rise to the placenta, septum, transmitting tract, style and stigma (Bowman et al., 1999). As discussed above, meristem-associated genes are expressed in this region and seem to be involved in early marginal tissue development.

Many genes have been shown to play a role in marginal tissue development (Fig. 6). Most of these share functional redundancy, and strong phenotypes in carpel development can thus frequently only be seen in multiple mutants. Among these genes, *ANT*, *LEUNIG (LUG)*, *SEUSS (SEU)* and *FIL* are major players in the development of carpel marginal tissues. Whereas the single mutants corresponding to these genes display relatively mild effects on the medial gynoecial domain, the gynoecium of double mutants combinations almost completely

lack marginal tissues. In *lug* mutants, the gynoecium is partially unfused at the apex, and also presents defects in septum and ovule development (Chen et al., 2000, Liu and Meyerowitz, 1995). The pistils of *ant* and *seu* single mutants present similar, though weaker, defects to those of *lug* mutants (Franks et al., 2002, Krizek et al., 2000). Strikingly, in the *ant lug* double mutant, the inner whorl of the flower consists of unfused valves-like structures which have style cells at their tips, but lack placenta, ovule, septum and stigma tissues (Liu et al., 2000). Similar carpel growth defects have been described in *lug seu*, *ant seu*, *fil ant* and *fil lug* double mutants (Azhakanandam et al., 2008, Chen et al., 1999, Franks et al., 2002, Nole-Wilson and Krizek, 2006). These four genes have been shown to repress *AG* in the first two whorls of the flower (Chen et al., 1999, Franks et al., 2002, Krizek et al., 2000, Liu and Meyerowitz, 1995). *LUG* and *SEU* encode transcriptional co-repressors that can interact to form a regulatory complex (Elliott et al., 1996, Siegfried et al., 1999, Sridhar et al., 2004). From all this information, it has been suggested that ANT, LUG, SEU and FIL form a multimeric complex involved in marginal tissue development in the gynoecium (Azhakanandam et al., 2008, Nole-Wilson and Krizek, 2006).

*SPT*, which has been mentioned above for its role in other aspects of gynoecium morphogenesis, is also a major factor in the formation of carpel marginal tissues. In *spt* mutants, the stigma and the style are reduced, the septum is distorted, and the transmitting tract is not properly formed (Alvarez and Smyth, 1999). As this phenotype can be partially rescued by the chemical inhibition of polar auxin transport (PAT), it has been proposed that *SPT* mediates auxin signalling (Heisler et al., 2001). Some other factors have also been specifically related to transmitting tissue development. The three closely related bHLH genes *HECATE1* (*HEC1*), *HEC2* and *HEC3* have been shown to redundantly specify stigma and transmitting tract development (Gremski et al., 2007). HEC proteins are able to physically



interact with SPT in yeast-two hybrid assays, so it has been suggested that HEC and SPT function together in this process. Crawford et al. (2007) described the first gene specifically required for *Arabidopsis* transmitting tract development: *NO TRANSMITTING TRACT (NTT)*. In *ntt* mutants, pollen tubes are unable to migrate efficiently due to the lack of transmitting tissues, and fertility is consequently much reduced.

Another set of factors that promote marginal tissue development in the apical domain of the *Arabidopsis* gynoecium are the members of the SHI/STY gene family, which encode zinc-finger transcriptional activators. *STYLISH1 (STY1)* is the only such factor showing a phenotype as a single mutant: *sty1* presents subtle defects in style development (Kuusk et al., 2002), though this phenotype is gradually enhanced in combination with mutations in further SHI/STY family members. Accordingly, multiple SHI/STY mutants show radically reduced style and stigma tissues, an abnormal septum and incomplete closure of the gynoecium apex. This phenotype is similar to that observed in *lug* mutants; moreover, it has been shown that the *lug* mutation is epistatic over *sty1*, and that *STY* expression is reduced in *lug* mutants. Thus, *STY* factors may act downstream of *LUG* to mediate marginal tissue formation (Kuusk et al., 2006). *STY1* activity has been linked to auxin as *STY1* is a direct activator of *YUC4*-an auxin biosynthesis gene (Eklund et al., 2010, Sohlberg et al., 2006; see Section IV.F for further details). Recently, the small NGATHA (*NGA*) gene family, encoding B3-domain transcription factors, has been described. The *NGA* genes are redundantly involved in style development (Alvarez et al., 2009, Trigueros et al., 2009). In the quadruple *nga* mutant, the style and stigma are completely absent, similarly to *shi/sty* multiple mutants. In addition to showing similar mutant phenotypes, *NGA* and *STY* genes share similar expression patterns and it has been shown that *YUC2* and *YUC4* expression is radically reduced in the *nga* quadruple mutant. Moreover, simultaneous over-expression of *NGA* and *STY* transforms the

ovary into style tissue. Accordingly, Trigueros *et al.* (2009) have suggested that *NGA* and *STY* may act cooperatively in style development, at least partially by promoting YUC-mediated auxin biosynthesis in the apical region of the gynoecium. While the precise regulatory hierarchy of *NGA* and *SHI/STY* factors has not yet been elucidated, several pieces of evidence point to a positive feedback loop acting between *STY* and *NGA* (Alvarez *et al.*, 2009, Trigueros *et al.*, 2009).

The *SHP* genes have been mentioned above for their important role in carpel identity and valve margin differentiation. In addition however, a recent study has indicated these genes to be also involved in style and stigma development (Colombo *et al.*, 2010). Gynoecia of the *ant crc shp1 shp2* quadruple mutant almost completely lack marginal tissues, resembling those of *ant lug* or *fil ant* double mutants. Surprisingly however, *STY* and *NGA* expression was not reduced in this quadruple mutant, while *SHP* expression was absent in the apical part of the carpel in *nga* quadruple mutants, and expanded when *NGA* genes were over-expressed (Alvarez *et al.*, 2009, Colombo *et al.*, 2010). These data suggest that *NGA* activity acts upstream of *SHP* expression in the style. Thus, *SHP* genes seem to act in a complicated and not fully understood regulatory network that controls most of the events which direct patterning and tissue specification in the gynoecium.

#### **F. Establishment of apical-basal polarity in the gynoecium**

Auxin is clearly one of the major morphogens involved in apical-basal patterning of the gynoecium: mutations in genes such as *PIN* and *PID*, which mediate polar auxin transport (PAT), result in defects in apical-basal patterning, producing enlarged apical and basal regions concomitantly with drastically reduced ovaries (Bennett *et al.*, 1995, Okada *et al.*, 1991). Similar defects are also found in mutants which lack functional auxin response factors

MONOPTEROS (MP/ARF5) or ETT/ARF3 (Przemeck et al., 1996, Sessions et al., 1997, Sessions and Zambryski, 1995). Nemhauser et al. (2000) have accordingly proposed that an auxin gradient spans the gynoecial primordium, and controls apical-basal patterning. Their model predicts maximum auxin levels at the apex, which induce the differentiation and proliferation of the stigma and style, and intermediate and low auxin levels lower down the gynoecium, which respectively specify the development of the ovary and gynophore (Fig. 7). In accordance with this model, the regulators of auxin biosynthesis *SHI/STY* and *NGA*, in addition to their downstream targets, the auxin biosynthesis genes *YUC4* and *YUC2*, have recently been shown to be expressed in the apex of the young gynoecium (Alvarez et al., 2009, Eklund et al., 2010, Kuusk et al., 2002, Kuusk et al., 2006, Sohlberg et al., 2006, Trigueros et al., 2009) as is also the auxin response reporter construct *ProDR5:GFP* (Aloni et al., 2003, Benkova et al., 2003). According to this model, auxin is trapped apically at the biosynthesis site under condition in which PAT is reduced, causing shifts in the boundaries between the different tissues which form along the apical-basal axis (Fig. 7). Interestingly, PAT inhibition, as well as over-expression of the auxin biosynthesis activator *STY1*, can restore style and stigma proliferation in mutant lines including: *lug*, *seu*, *ant*, *styl*, *spt*, *crc* and *jag*, which are affected in the development of apical tissues that are derived from marginal regions of the gynoecium. These observations suggest that auxin may act downstream of, or in parallel to, corresponding apical tissue-promoting factors during style and stigma development (Chen et al., 2000, Nemhauser et al., 2000, Sohlberg et al., 2006, Staldal et al., 2008, Staldal and Sundberg, 2009). An exception to this is the *nga* quadruple mutant, suggesting *NGA* genes to act not only upstream, but also downstream of auxin (Alvarez et al., 2009).

The repression of the apical program in the zone of the gynoecium corresponding to intermediate auxin levels may be required for the specification of the ovary. It has been suggested that ovary size may be controlled via a specific response of ETT/ARF3 to intermediate auxin levels (Heisler et al., 2001, Sessions et al., 1997, Sessions and Zambryski, 1995). As ETT/ARF3 represses the activity of *SPT* and the *HEC* genes, which promote stigma, style and transmitting tract differentiation (Alvarez and Smyth, 1999, Gremski et al., 2007, Heisler et al., 2001), ETT activity in the intermediate zone locally represses the apical developmental program. Gynophore development may be restricted to the basal end of the gynoecium by the activity of the zinc-finger protein KNUCKLES (KNU), via the establishment or maintenance of a tissue boundary at this location (Payne et al., 2004).

Though the auxin gradient model can be used to explain apical-basal patterning in the gynoecium, there is at present no direct evidence of the existence of such a gradient, other than the presence of an auxin maximum at the gynoecium apex. Auxin level measurements along the length of the developing gynoecial cylinder have not yet been made, and so the hypothesised intermediate and low auxin levels in the ovary and gynophore have yet to be demonstrated. In addition, it has recently been suggested that the basal end of the gynoecium may be distinguished from the ovary by an opposing cytokinin gradient (Ostergaard, 2009). We also need to know more about how a potential auxin gradient could be interpreted. Lines carrying mutations in some of the genes involved in the promotion of apical tissues, such as *LUG*, *SEU* and *STY*, are hyper-responsive to the chemical or genetic inhibition of PAT (Pfluger and Zambryski, 2004, Sohlberg et al., 2006, Staldal et al., 2008), as are auxin biosynthesis mutants (*yuc1*, *yuc4*) and auxin response mutants (*axr1-3*, *ett/arf3* and *tir1*) (Cheng et al., 2007, Nemhauser et al., 2000, Staldal et al., 2008), suggesting that this group of genes could promote either auxin gradients or response pathways. Conversely, *jag* and *ant*

mutants respond to PAT reductions to the same extent as wild type (Staldal and Sundberg, 2009), suggesting that JAG and ANT may not be involved in the establishment or responses to the hypothesised auxin gradient. Interestingly, *spt*, *nga* and *crc* mutants are less sensitive to PAT inhibition (Alvarez et al., 2009, Nemhauser et al., 2000, Staldal et al., 2008), which may suggest that these genes could participate in the modulation of PAT, or in the sensing of parts of the auxin gradient.

### G. Post-fertilisation carpel growth

Upon fertilisation, gynoecium and ovule developmental programs are switched to those of fruit and seed development (Fig. 7). In *Arabidopsis*, fruit development is characterised by a dramatic elongation of the ovary, concomitant with the differentiation of specific tissues along the carpel margins. Unfertilised gynoecia fail to elongate and develop seeds, and will eventually undergo senescence. Interestingly, the switch to fruit development after fertilisation appears to rely on a hormone-induced signal evoked in the fertilised ovules, and some data suggest that auxin signalling participates in this process (Dorcey et al., 2009, Goetz et al., 2006, Vivian-Smith et al., 2001). One piece of evidence supporting this suggestion is that a knock-out of the auxin response factor *ARF8* gene results in parthenocarpy, or fertilisation-independent fruit development (Goetz et al., 2006, Vivian-Smith et al., 2001).

The major external parts of the *Arabidopsis* ovary comprise the pod walls, the replum, which extends along the length of the fruit, and the carpel margins, which form at the carpel/replum border where fruit opening will occur. During fruit development, the carpel margins differentiate into narrow strips consisting of a separation layer and a lignified layer, both of which contribute to the process of fruit opening (see Section IV.E). The key regulators of valve margin specification have been identified (Ferrandiz et al., 2000a, Liljegren et al., 2004,

Roeder et al., 2003), and it was recently shown that a local auxin minimum, generated by the valve margin identity factor IND, is required for separation layer development (Figs 4 and 5; Sorefan et al., 2009). Thus, both auxin maxima and minima appear to contribute to the activation of specific developmental programmes during fruit development.

## **V. Floral Meristem Termination in the Central Zone of the *Arabidopsis* Gynoecium**

At the time when carpel primordia are initiated, another essential process takes place in the centre of the *Arabidopsis* flower, in a domain that will later correspond to the base of the gynoecium. Stem cells in this domain, which had previously been maintained within the flower bud, and whose divisions had generated the different floral organs, cease to be maintained as the carpel primordial form. This disruption of stem cell maintenance makes the flower determinate, and assures its future fertility by blocking the development of floral organs, in the place of ovules, within the gynoecium.

### **A. *AGAMOUS* (*AG*), the main gene responsible for carpel development, also plays a central role in floral meristem termination**

*AG* function is not restricted to the control of stamen and carpel identity. It also promotes floral meristem termination (Fig. 8): in strong *ag* mutants (*ag-1* to *ag-3*), stamens are transformed into petals, and carpels are replaced by a new flower bud, which turns into a new, abnormal flower (Bowman et al., 1991b, Yanofsky et al., 1990). Stem cells are thus indefinitely maintained within the FM, and allow for the endless production of floral organs. Interestingly, FM termination is the most sensitive role of *AG*: weaker *ag* alleles (eg. *ag-4* and *AG-Met-205*) trigger a strong loss of FM termination, but fewer organ identity defects (Sieburth et al., 1995), while indeterminacy is the first phenotypic flaw associated with reduced levels of *AG* (Chuang and Meyerowitz, 2000, Mizukami and Ma, 1995).

A delay or loss of FM termination has also been observed in various mutant backgrounds or transgenic plants (Alvarez et al., 1999, Bowman et al., 1992, Chen et al., 2004, Clark et al., 1993, Clark et al., 1995, Das et al., 2009, Fletcher, 2001, Jacobsen et al., 1999, Kayes et al., 1998, Liu et al., 2010, Maier et al., 2009, Payne et al., 2004, Prunet et al., 2008, Schultz et al., 1991, Sun et al., 2009, Zhao et al., 2007). However, unlike *AG*, none of the genes corresponding to these mutations, with the notable exception of *KNU*, are strictly required for the arrest of stem cell maintenance within the FM. Moreover, most of these genes appear to act upstream of *AG*, and thus control FM termination through this factor. For example, mutations to *ULT1*, *CLAVATA1 (CLV1)* or *PERIANTHIA (PAN)* can trigger a defect in *AG* expression (Clark et al., 1993, Das et al., 2009, Fletcher, 2001, Maier et al., 2009). *DICER-LIKE1/CARPEL FACTORY (DCL1/CAF)* and *HUA ENHANCER1 (HEN1)* encode proteins that are required for the proper accumulation of *miR172* (Chen et al., 2002, Jacobsen et al., 1999, Park et al., 2002), which promotes *AG* expression by down-regulating the *AG* repressor *AP2* (Aukerman and Sakai, 2003, Chen et al., 2002, Chen et al., 2004). Other proteins that participate in the control of FM termination control *AG* expression at the post-transcriptional level: *HUA1*, *HUA2*, *HEN2* and *HEN4* are required for proper splicing of *AG* pre-mRNA (Cheng et al., 2003), while the interaction of *AG* with *SEP* proteins is necessary for *AG* to perform its function (Ditta et al., 2004, Goto et al., 2001, Pelaz et al., 2001).

**B. The B-class genes *APETALA3 (AP3)* and *PISTILLATA (PI)* promote the male developmental programme and antagonise FM termination**

The various functions of *AG* rely on its interactions with different protein partners. In the third whorl, *AG* participates in a protein complex with *AP3* and *PISTILLATA (PI)* which promotes stamen identity (Fig. 8). *AP3* and *PI* are excluded from the fourth whorl, and cannot

therefore interact there with AG, and this results in carpel identity and FM termination (Goto et al., 2001, Krizek and Fletcher, 2005, Theissen, 2001, Theissen and Saedler, 2001). Indeed, the loss of function of *AP3* or *PI* results in over-determinate flowers with stamens transformed into carpels, and a strongly reduced number of floral organs within whorls three and four, compared to wild type (Fig. 8; Bowman et al., 1991b). Over-determinacy is even more striking in *Antirrhinum majus* plants which are mutant for *DEFICIENS* or *GLOBOSA*: the respective orthologs of *AP3* and *PI*, the flowers of which entirely lack a fourth whorl (Sommer et al., 1990, Trobner et al., 1992). Conversely, the over-expression of *AP3* alone (*p35S::AP3*), or together with *PI* (*p35S::AP3/PI*), as well as the over-expression of their activator *UFO* (*p35S::UFO*), delays FM termination: flowers of *p35S::AP3*, *p35S::AP3/PI* and *p35S::UFO* plants exhibit several extra whorls of stamens, while carpels in these plants are often staminoid or absent (Jack et al., 1994, Krizek and Meyerowitz, 1996, Lee et al., 1997). B-class genes thus appear to antagonise AG's fourth whorl functions, including the arrest of stem cell maintenance.

Plants which are mutant for the C2H2 zinc-finger transcription repressor *SUPERMAN* (*SUP*) exhibit a phenotype similar to that of plants over-expressing B-class genes: *sup* mutant flowers exhibit extra stamens which usually develop at the expense of carpels (Bowman et al., 1992, Schultz et al., 1991), although some alleles also trigger a moderate increase in carpel number (Jacobsen and Meyerowitz, 1997, Rohde et al., 1999). The extra stamens in *sup* mutants are indeed associated with an expansion of B-class gene expression within the fourth whorl (Bowman et al., 1992, Goto and Meyerowitz, 1994). Two models have been proposed to explain the origin of the extra organs in *sup* mutant flowers. One of these proposes the expansion of B-class gene expression observed in *sup* flowers to be responsible for prolonging the developmental state of the FM which normally precedes carpel initiation and



stem cell termination. According to this model, SUP thus promotes flower determinacy indirectly (Bowman et al., 1992, Schultz et al., 1991). Conversely, the other alternative model proposes SUP to directly repress cell division within the inner part of the third whorl, in which it is expressed together with B-class genes (Sakai et al., 1995; Sakai et al., 2000). According to this second model, extra stamens would then be generated by increased cell division in the inner part of the third whorl, and the expansion of B-class gene expression would thus be a consequence, rather than a cause, of this increased rate of division. More recently, ectopic expression, using various promoters, of *SUP* and its orthologs in several species, has shown these genes to be able to repress both the expression of B-class genes and cell proliferation (Bereterbide et al., 2001, Hiratsu et al., 2002, Nakagawa et al., 2004, Nandi et al., 2000, Yun et al., 2002). Neither model can thus be excluded so far, though ectopic expression of B-class genes is likely to at least participate in the formation of extra organs in *sup* mutant flowers, as reviewed in more detail by Prunet et al. (2009).

Several pieces of evidence suggest that stamen and carpel identity, as well as FM termination, are very sensitive to the ratio of AG to AP3/PI proteins. Firstly, increased ectopic expression of B-class genes correlates with an increased indeterminacy phenotype, as shown by the greater number of stamens in *p35S::AP3*, *sup-1* and *p35S::AP3 sup-1* flowers (Jack et al., 1994) and in *p35S::AP3*, *sup-1*, *p35S::AP3/PI* and *p35S::AP3/PI sup-1* flowers (Krizek and Meyerowitz, 1996). Similarly, a reduction in the dose or activity of AG strongly enhances the indeterminacy phenotype of *sup: ag-1/AG sup-1* and *ag-4 sup-1* flowers are fully indeterminate (Prunet et al., 2008, Schultz et al., 1991). Conversely, increased expression of AG within the third whorl, in which AG is normally expressed together with AP3 and PI, is sufficient to trigger a partial transformation of stamens into carpels (Lohmann et al., 2001). Also, stamen identity is less sensitive to a reduced dose of AG than is carpel identity or FM

termination (Chuang and Meyerowitz, 2000, Mizukami and Ma, 1995). Given that AG participates in several different transcription factors complexes, together with AP3 and PI in whorl 3 and without them in whorl 4, the importance of the relative dose of AG to AP3/PI hints that B-class genes may oppose AG's fourth whorl functions, including FM termination, by competing with AG in the formation of these complexes.

C. CRABS CLAW (CRC) and SPATULA (SPT), two genes involved in carpel development, also participate to FM termination

AG is not the only gene involved in the female developmental program to promote FM termination. *CRC* and, to a lesser extent, *SPT* encode two other transcription factors that control carpel growth, polarity and congenital fusion, and which also participate in floral determinacy (Fig. 8; Alvarez and Smyth, 1999, Bowman and Smyth, 1999, Heisler et al., 2001). Flowers of *crc* and *spt* single mutants are normally determinate, but those of the *crc spt* double mutants, and more often those of *crc AG/ag-1* plants, are indeterminate and possess extra whorls of stamens, and secondary carpels that developing within the gynoecium (Alvarez and Smyth, 1999). A strong loss of FM termination is also seen when *crc* is combined with mutations in genes such as *ULT1*, *RBL* or *SQN*, which do not show an extensive indeterminacy phenotype as single mutants (Berardini et al., 2001, Fletcher, 2001, Prunet et al., 2008). Interestingly, in several other angiosperm species, a loss of *CRC* function is sufficient to cause strong floral indeterminacy (Lee et al., 2005, Nagasawa et al., 2003, Orashakova et al., 2009, Yamaguchi et al., 2004), suggesting that *CRC*'s ancestral role in FM termination may be more important than its current role in that process in *Arabidopsis*.

AG promotes the expression of *CRC* and *SPT*, which occurs through direct transcriptional activation in the case of *CRC* (Bowman and Smyth, 1999, Gomez-Mena et al., 2005, Heisler

et al., 2001). *CRC* and *SPT* may thus mediate a proportion *AG*'s functions in carpel development and flower determinacy. Most genes known to control this latter process act upstream of *AG*, as described above, but no positive feedback loop between *CRC*, *SPT* and *AG* has yet been shown. Indeed, the function of both *CRC* and *SPT* is at least partly independent of *AG* (Bowman and Smyth, 1999, Heisler et al., 2001). How *CRC* and *SPT* influence stem cell termination is thus currently unclear. However, the fact that genes controlling female development other than *AG* share its role in stem cell termination confirms the close link between these two processes. It is particularly interesting to note that *CRC*'s role in floral determinacy is non cell-autonomous: *CRC* is expressed in the emerging carpel primordial, but not at the base of the gynoecium where indeterminacy phenotypes first become apparent (Bowman and Smyth, 1999). This observation clearly suggests primordia to signal back to the FM to oppose stem cell maintenance.

#### **D. *AG* is required in the centre of the FM to trigger FM termination**

*AG*'s three functions, in stamen and carpel specification and in FM termination, are dose-dependent (Chuang and Meyerowitz, 2000, Mizukami and Ma, 1995), but can also be separated on a spatial basis. *pAP3::AG ag-3* plants express functional *AG* in the third whorl, but not in the fourth, and produce completely indeterminate flowers which lack carpels, but have normal stamens (Jack et al., 1997). *AG* is therefore required specifically in the fourth whorl to promote carpel development and FM termination. While this phenotype tends to confirm the close association between female development and stem cell termination, these two programs also can be spatially separated. Indeed, numerous mutants suggest that FM termination requires the expression of *AG* in an even more restricted domain within the fourth whorl. In the flowers of *clv*, *ult1* and *pan* single mutants and of *crc sqn* and *crc ult1* double mutants, stamens and carpels are correctly specified, suggesting that *AG* is still active within

whorls 3 and 4 of their flowers. However, AG fails to terminate stem cell maintenance in these mutants, which all exhibit a delay in or loss of flower determinacy (Clark et al., 1993, Das et al., 2009, Fletcher, 2001, Maier et al., 2009, Prunet et al., 2008). Indeed, depending on allelic strengths, the indeterminacy phenotype of these mutants is associated with a transient or persistent defect in AG transcription in an inner, intercarpellary domain within the fourth whorl (Clark et al., 1993, Das et al., 2009, Fletcher, 2001, Maier et al., 2009, Prunet et al., 2008). Interestingly, a similar defect in AG expression is seen in the flowers of plants over-expressing a modified version of AG's repressor AP2 which has been made resistant to *miR172* (*35S::AP2m3*): these flowers are fully indeterminate, with numerous supernumerary stamens (Chen et al., 2004, Zhao et al., 2007). AG is thus specifically required at the base of the gynoecium to promote FM termination.

**E. To terminate stem cell maintenance, AG promotes a specific developmental programme at the base of the gynoecium**

The subdomain of whorl 4 at the base of the gynoecium corresponds to the centre of the FM and contains the floral stem cells. AG disrupts the maintenance of these cells by switching off the expression of the stem cell-promoting gene *WUS* at the moment of emergence of the carpel primordial. Accordingly, *WUS* mRNA becomes undetectable at Stage 6 (Smyth et al., 1990) of determinate, wild type flower development (Mayer et al., 1998), but persists until later stages in indeterminate, *ag* mutant flowers (Lenhard et al., 2001, Lohmann et al., 2001). Such a maintenance of *WUS* expression beyond Stage 6 has been described in most mutant or transgenic plants with indeterminate flowers (Carles et al., 2004, Das et al., 2009, Maier et al., 2009, Prunet et al., 2008, Schoof et al., 2000, Sun et al., 2009, Zhao et al., 2007) and appears both necessary and sufficient to maintain stem cells within the flower, unlike another pro-meristematic gene, *STM* (Lenhard et al., 2001, Lohmann et al., 2001).

AG is required in L2 cells, in which *WUS* is not expressed, to disrupt stem cell maintenance (Mayer et al., 1998, Sieburth et al., 1998), suggesting that, despite its role as a transcription factor, AG does not directly repress *WUS* transcription. Indeed, recent data suggest that *KNU*, a C2H2 zinc-finger protein (Payne et al., 2004), is the main intermediate between AG and *WUS* (Fig. 8; Sun et al., 2009). Accordingly, AG is required for the expression of *KNU*, and directly binds to its promoter, and in turn, *KNU* appears both necessary and sufficient to turn off the expression of *WUS* and therefore stem cell maintenance within the flower. Interestingly, *KNU* expression starts at Stage 6 and then becomes restricted to precisely the domain at the base of the gynoecium in which AG expression is specifically required for FM termination, as described above (Payne et al., 2004, Sun et al., 2009). Later on, *KNU* expression is also detected in anthers, though remains very strong at the basis of the gynoecium. The importance of *KNU* in FM termination, together with its expression pattern, confirms AG to activate a specific developmental program in the centre of the FM, resulting in the disruption of stem cell maintenance at Stage 6 of flower development.

## **F. Conclusions**

The data discussed above clearly demonstrate FM termination to be closely associated with the development of the gynoecium. FM termination relies on a genetic network centred on *AG*, which is responsible for both male and female developmental programs. However *AG* is able to promote carpel development and stem cell termination only if the male-promoting B-class genes are excluded from the centre of the flower. The *AG*-induced disruption of stem cell maintenance mainly depends on a cascade of transcription factors in a subdomain of whorl four, at the base of the gynoecium: *AG* activates *KNU* specifically at this location, and *KNU* in turn switches off the expression of *WUS*. However, FM termination also involves

signals from the developing carpels: another target of AG, *CRC*, is expressed only in carpel primordia, and not at the base of the gynoecium, but nonetheless participates by an as yet unknown mechanism in FM termination.

## **VI. Carpel Diversification in the Angiosperms**

### **A. A phylogenetic background**

To reconstruct the different paths that carpel evolution has taken in distinct angiosperm groups, it is first necessary to gain some insight into phylogenetic relationships within the angiosperms. Several independent studies published around ten year ago (reviewed by Kuzoff and Gasser, 2000) provided an early outline of angiosperm phylogeny, and this has been expanded and updated ever since by the Angiosperm Phylogeny Group (Stevens, 2001 onwards). These phylogenetic studies indicate the angiosperms to form a monophyletic clade in which three extant groups, Amborellales, Nymphaeales and Austrobaileyales, collectively known as the ANA grade, diverged in a series of basal bifurcations to leave a remaining lineage from which all other extant angiosperms are descended (Fig. 9). Amborellales contains the single species *Amborella trichopoda*, a shrub endemic to the South Pacific island of New Caledonia. Nymphaeales contains the three families of herbaceous aquatic plants Nymphaeaceae, Cabombaceae, and Hydatellaceae, which include a total of around 100 species, and Austrobaileyales contains Austrobaileyaceae, Schisandraceae (incorporating Illiciaceae) and Trimeniaceae, which are composed of a total of around 100 woody species. Following the divergence of the ANA lineages, the remaining angiosperm lineage later diversified to form the five further extant angiosperm groups of: eudicots, monocots, eumagnoliids, Chloranthaceae and *Ceratophyllum*. Of these, the eudicots and monocots together account for over 95% of the estimated 300 000 or more extant angiosperm species.

The resolution of angiosperm phylogenetic relationships provides an excellent framework to analyse the processes through which carpel development has diversified throughout the flowering plants. However, a further finding of molecular phylogenetic studies provides less encouraging news for the analysis of carpel evolution. To understand how the carpel first appeared, it would be very useful to have available for study one or more living groups whose lineages had diverged from that of the flowering plants shortly before the origin of the carpel. However, molecular phylogenetic analyses indicate the remaining seed plants, or gymnosperms, to form a sister clade to the angiosperms (Fig. 9), meaning that no individual group of living gymnosperms appears to be more closely related to the angiosperms than any other. Molecular clock estimates suggest a date for the divergence of the angiosperm and gymnosperm lineages of around 300 MYA (Goremykin et al., 1997, Savard et al., 1994), whereas the earliest known angiosperm macrofossils date from around 125 MYA (Sun et al., 1998). Thus, the extant flowering plants and gymnosperm lineages may have diverged up to 175 MY before the origin of the flower and carpel.

As will be apparent from Section II, above, lack of evidence concerning the origin of the angiosperms has not prevented the construction of numerous hypotheses for the evolution of the flower and carpel. However, given the absence of a living, non-flowering, close relative of the flowering plants, it seems likely that a full understanding of the evolutionary origin of these structures will require the inclusion of fossil data. In this respect, further evidence from mesofossils may prove extremely valuable. These are small fossils, of up to a few millimetres in diameter, many of which are “coalified” specimens that appear to have been generated by forest fires. Numerous early angiosperm mesofossils containing considerable anatomical detail have already been discovered (Friis et al., 2001, Friis et al., 2010) raising hopes that

further discoveries may provide important new insights into early flower evolution (Frohlich and Chase, 2007).

**B. Carpel morphology and function in ANA grade angiosperms**

Morphological comparisons of the three extant lineages of ANA grade angiosperms (Fig. 9) have enabled a number of conclusions to be made on the likely state of the gynoecium in the last common ancestor of the living flowering plants (Endress, 2001, Endress and Igersheim, 2000). The gynoecium in all extant ANA grade taxa is apocarpic, except Nymphaeaceae, which are syncarpic. Apocarpy is thus present throughout Amborellales and Austrobaileyales, and is also the basal condition in Nymphaeales, clearly implying this to be an ancestral trait of the living angiosperms. The carpels of apocarpic ANA grade angiosperms are ascidiate (bottle-shaped) rather than plicate (folded), as in many later diverging groups, implying the carpel to have been ascidiate in early angiosperms.

The carpels of many ANA grade species are not only separate from each other (apocarpic), but are also incompletely closed by cellular structures at maturity, leaving a secretion-filled aperture or canal through which pollen tubes grow to bring about fertilisation (Fig. 10A). The only exceptions to this are *Illicium* (Austrobaileyales, Schisandraceae) and Nymphaeales, in which closure of the carpel margins occurs, at least in part, though postgenital cell divisions (Endress and Igersheim, 2000). Thus, comparative analysis of ANA grade species clearly indicates the basal condition of the angiosperms to have been carpels that were closed at the apex by substances secreted from their margins, rather than by post-genital cell division.

In some ANA grade angiosperms, the stigmatic surface is covered by multicellular protrusions, rather than by the unicellular papillae present in most later diverging groups. This



is the case in *Amborella* (Amborellales) and *Trimenia* (Austrobaileyales), both of which have stigmatic surfaces containing multicellular ridge-like structures, and in Cabombaceae and Hydatellaceae (Nymphaeales), which possess bi- or multicellular stigmatic hairs (Endress, 2005, Endress and Igersheim, 2000, Rudall et al., 2007). However, all other Austrobaileyales and most Nymphaeaceae have stigmatic surfaces covered with unicellular papillae (Endress, 2001). Thus, multicellular protrusions on stigmatic surfaces are present in all three ANA grade lineages, leaving open the possibility that this may represent the ancestral condition in the angiosperms.

Self-incompatibility (SI) systems operating between carpel tissues and pollen grains are present in the Austrobaileyales species: *Illicium floridanum* (Thien et al., 1983), *Austrobaileya scandens* (Prakash and Alexander, 1984) and *Trimenia moorei* (Bernhardt et al., 2003). It is not yet clear whether these SI systems are homologous, which would make SI an ancestral feature of Austrobaileyales. *Amborella*, the only representative of the likely most basally diverging angiosperm lineage Amborellales, avoids in-breeding by dioecy, rather than through an SI mechanism. However, male *Amborella* flowers contain a structure which may be a relictual gynoecium, and female flowers of this species contain staminodes, or non-functional stamens (Endress, 2001), strongly suggesting *Amborella* to be descended from a bisexual ancestor. Hydatellaceae (Nymphaeales) are either dioecious, or possess bisexual reproductive units that may be derived from unisexual flowers (Rudall et al., 2007), suggesting their possible descent from a monocious ancestor. Most other out-breeding members of the ANA grade are protogynous, while *Barclaya* and *Euryale* (Nymphaeaceae) are partially cleistogamous and inbreeding (Endress, 2001). Protogyny, and possibly even SI, may thus have been present to promote out-breeding in the last common ancestor of the living

angiosperms. By contrast, it seems clear that outbreeding by dioecy, and inbreeding mechanisms, which are sporadically present in other ANA groups, have arisen secondarily.

Ovules in all ANA grade angiosperms are anatropous, except in *Amborella* and *Barclaya* (Nymphaeaceae) which have orthotropous ovules (Endress, 1986, Schneider, 1978). Of these, the *Amborella* ovule shows a residual curvature near its point of attachment, suggesting a previously anatropous structure (Endress and Igersheim, 2000). Thus, the ovule of the last common ancestor of the extant angiosperms was probably anatropous. Similar comparisons (Endress and Igersheim, 2000) indicate that this ancestral ovule was probably covered by two integuments, and crassinucellar (containing a large nucellus tissue).

Double fertilisation, leading to the production of an embryo and a bi-parental endosperm, also appears to be a plesiomorphic feature of the angiosperms. In the majority of flowering plants, the embryo sac arrangement is that of the *Polygonum* type, which contains seven cells of which the central cell is binucleate (Fahn, 1982). The two nuclei of this central cell combine with one sperm nucleus following fertilisation to generate a triploid endosperm. However, in the ANA grade taxa *Nuphar* and *Trithuria* (incorporating *Hydatella*) of Nymphaeales, and *Illicium* of Austobaileyales, the embryo sac contains only four cells, including a uninucleate central cell (Friedman, 2008, Williams and Friedman, 2002; Williams and Friedman, 2004). Double fertilisation in these taxa generates an embryo and endosperm which are both diploid. However, the likely most basal ANA grade lineage, Amborellales, does not fit this pattern: the *Amborella* embryo sac contains eight cells, including a binucleate central cell that produces a triploid endosperm after fertilisation (Friedman, 2006). The *Amborella* embryo sac arrangement is thus more similar to that of *Polygonum* and the majority of later-diverging angiosperms than to other members of the ANA grade, though it contains one extra cell in its

egg apparatus compared to the *Polygonum* type. The major difference in embryo sac arrangement between Amborellales and other ANA lineages renders the ancestral state of the embryo sac and endosperm ploidy ambiguous in the extant angiosperms. Interestingly, a perisperm is present in *Trithuria*, in addition to its diploid endosperm (Friedman, 2008). Like the endosperm, the perisperm is an embryo-nourishing tissue, though one which is derived exclusively from maternal cells. The presence of a perisperm is mainly associated with gymnosperms, and the occurrence of such a tissue in *Trithuria* has been suggested to form a link with the hypothesised gymnosperm-like ancestor of the flowering plants (Friedman, 2008).

### C. The molecular control of carpel development in the ANA grade

Phylogenetic analyses of MADS box genes in ANA grade angiosperms and gymnosperms suggest a duplication event to have taken place in the C-function lineage prior to the last common ancestor of the living flowering plants (Kim et al., 2005). As a result of this duplication, the ancestors of the clade-defining genes *AG* from *Arabidopsis thaliana*, and *FLORAL BINDING PROTEIN7 (FBP7)* from *Petunia hybrida* (reviewed by Kramer et al., 2004) were generated, and both of these lineages were thus present in the last common ancestor of the flowering plants. The *AG* clade contains angiosperm C-function genes, whereas the *FBP7* clade contains genes involved in ovule development in diverse later diverging angiosperm groups (Angenent et al., 1995, Colombo et al., 1995, Dreni et al., 2007). In addition to the C- and D-clades, two clades of *SEP* genes, encoding E-function MADS box proteins, have been found in basal angiosperms. The genes *SEP1*, *SEP2* and *SEP4* from *Arabidopsis* appear to be orthologous to one of these ANA grade *SEP* clades, while *SEP3* appears to be orthologous to the other (Zahn et al., 2005).

The expression of C-function genes in ANA grade angiosperms is mostly limited to the third and fourth floral whorls, while the *SEP*-like E-function genes of these species are expressed in all floral organs (Fig. 10A; Kim et al., 2005). These expression patterns closely resemble those of the corresponding genes in *Arabidopsis*, suggesting important elements of the control of carpel identity to have been conserved in both the *Arabidopsis* and ANA grade lineages throughout angiosperm evolution. Despite this broad conservation, Kim et al. (2005) noted some expression of C-function genes in the perianth organs of the ANA grade angiosperms *Amborella* (Amborellales) and *Illicium* (Austrobaileyales), in contrast to the expression patterns of C-function genes in model eudicots. However, as pointed out by these authors, such imprecise boundaries of gene expression may reflect the gradual transition of floral organ types that is apparent in ANA grade angiosperms, with intermediate organ types often present.

The expression patterns of several YAB transcription factors have also been analyzed in ANA grade angiosperms. As discussed in Sections III and IV above, studies of *Arabidopsis* indicate YAB factors to participate in the specification of abaxial cellular identity in lateral organs by defining the side of these organs that faces away from the developmental axis. The YAB gene *AmbCRC* (Fourquin et al., 2005), from the ANA grade angiosperm *Amborella*, shows a similar pattern of expression to that of its *Arabidopsis* orthologue *CRC* (Bowman and Smyth, 1999), suggesting these genes to have conserved a common developmental role in abaxial tissue specification in the carpel since the last common ancestor of the flowering plants. *INNER NO OUTER (INO)* represents a further YAB gene with a very specific role in female reproductive development. In *Arabidopsis*, *INO* is expressed specifically expressed in the outer ovule integument and is necessary for the development of this structure (Villanneva et al., 1999). A putative *INO* orthologue from the ANA grade angiosperm *Nymphaea alba* is

specifically expressed in both ovule integuments and in the suspensor (Yamada et al., 2003), suggesting this gene to have functioned in integument development since the last common ancestor of the flowering plants. Thus, the carpel and outer integument, which are both plesiomorphic features of the angiosperms, may have been associated with the expression of distinct YAB gene lineages throughout flowering plant evolution.

In general, the study of carpel development genes in ANA grade angiosperms has highlighted several instances of the broad conservation of gene function since the common ancestor of the last flowering plants. However, much work remains to be done in this field, as many families of transcription factors, and other regulatory proteins of known importance to carpel development, have yet to be analysed in ANA grade angiosperms. Furthermore, molecular studies of ANA grade angiosperms have, to date, relied principally on expression data to infer gene function. This approach may be tenable in the case of genes with highly characteristic and specific expression patterns, such as the MADS box and YAB genes discussed above. However, further substantial evolutionary-developmental advances in ANA grade angiosperms will surely require the development of methods for the direct study of gene function in these species.

#### **D. Carpel development in monocots**

The monocots form a monophyletic group of angiosperms numbering some 60 000 species. Phylogenetic studies have identified the small genus *Acorus* as sister to all other monocots, with the moderately large Alismatales as the next-earliest diverging group. *Acorus* contains a syncarpic gynoecium of three fused carpels (Fig. 10B), and comparison with Alismatales indicates a trimerous syncarpic arrangement to be the probable ancestral condition in the monocots (Igersheim et al., 2001). From similar comparisons, the presence of more than two

ovules per carpel can also be concluded as a probable ancestral feature of the monocots. Other characteristics of the gynoecium in the last common ancestor of the monocots are, however, more difficult to infer, partly due to differences between *Acorus* and other basal monocot lineages. For example, the carpels of *Acorus* are completely closed by postgenital cell divisions, whereas postgenital fusion is absent or partial in other early-diverging monocots (Igersheim et al., 2001), with the exception of Tofeldiaceae (Alismatales). Similarly, ovules are pendent in *Acorus*, but ascendant in other basal monocot groups, rendering the ancestral state of ovule orientation undetermined in monocots. Most molecular-developmental studies of monocots have, to date, been performed in two models from Poaceae (the grass family): *Oryza sativa* (rice) and *Zea mays* (maize). Poaceae are highly derived monocots in which the gynoecium, which contains a single ovule, appears also to have been reduced to a single carpel (discussed by Kellogg, 2001 and Rudall et al., 2005).

Phylogenetic analyses of carpel development genes in Poaceae suggest at least one major duplication event to have occurred in the MADS box C-clade, prior to the separation of the rice and maize lineages, with a further additional duplication in the maize lineage. Accordingly, the rice C-clade gene *OsMADS58* is orthologous to *ZAG1* in maize, while *OsMADS3* from rice is orthologous to both *ZMM2* and *ZMM23* (Yamaguchi et al., 2006). The inactivation of *OsMADS58* in rice leads to defects in carpel development, though does not completely eliminate carpels (Fig. 10B; Yamaguchi et al., 2006). In addition to abnormal carpels, *osmads58* mutants show a reduced level of determinacy in the spikelet (or Poaceae-type flower). Whereas *OSMADS58* appear to act mainly in the fourth whorl, the inactivation of its paralogue *OsMADS3* has little or no effect on either carpel development or floral determinacy. Instead, stamen development is eliminated in *osmads3* mutants (Kang et al., 1998, Yamaguchi et al., 2006). Rice plants in which both *OsMADS3* and *OsMADS58* have

been inactivated produce aberrant carpels, similar to those of *osmads58* mutants, suggesting *OsMADS3* to make no specific contribution to carpel development (Yamaguchi et al., 2006). In maize, *zag1* mutants show a defect in floral determinacy, indicating the functional conservation of *ZAG1* with its rice orthologue *OsMADS58*. It appears, therefore, that C-clade MADS box genes in the grass family have undergone significant sub-functionalisation following a monocot-specific gene duplication. The well known C-clade functions of carpel development, stamen development and floral determinacy are thus shared in rice and maize in a whorl specific manner between two and three C-clade genes, respectively. The functions of a paralogous pair of D-clade MADS box genes, *OsMADS13* and *OsMADS21*, have also been investigated in rice (Dreni et al., 2007). Of these, *OsMADS13* is ovule-specifically expressed, and its inactivation accordingly results in the ectopic conversion of ovules into internal carpeloid organs, similar to the D-function knockout phenotype observed in *Petunia* (Angenent et al., 1995, Colombo et al., 1995). *OsMADS21*, by contrast, appears to make no significant contribution to the D-function, and is expressed more widely than its paralogue in female reproductive tissues (Dreni et al., 2007).

Carpels are entirely replaced by stamens in rice mutants in which the YAB gene *DL* has been inactivated (Fig. 10B, Yamaguchi et al., 2004), which also causes the loss of the leaf mid-rib. In agreement with these functions, *DL* is expressed throughout the carpel anlagen (presumptive primordium), and in leaves, and its orthologues share similar expression patterns in other Poaceae (Ishikawa et al., 2009). It is not yet clear whether carpel development in rice depends on *DL* expression *per se*, or whether *DL* is responsible for preventing B-function gene expression, and thus ectopic stamen development, in the fourth floral whorl. *DL* is the rice orthologue of *CRC* in *Arabidopsis*, and its coding sequence, when expressed from the *CRC* promoter, is able to fully rescue *Arabidopsis crc* mutants

(Fourquin et al., 2007). It therefore seems that the functional differences between *DL* and *CRC* may reside principally outside their coding sequences. *DL* expression is maintained in the carpels of rice plants in which both *OsMADS3* and *OsMAD58* have been inactivated (Yamaguchi et al., 2006), demonstrating its action to be independent of these C-clade MADS box genes. This characteristic may represent a further difference between *DL* in rice and *CRC* in *Arabidopsis*, as the latter is known to be a direct target of the C-function transcription factor AG (Gomez-Mena et al., 2005).

*OsMADS1* from rice corresponds to the *LEAFY HULL STERILE1* locus, and groups within the *SEP1* Clade of E-function MADS-box genes which contains the genes *SEP1*, *SEP2* and *SEP4* from *Arabidopsis* (Zahn et al., 2005). Outer whorl floral organs in *osmads1* loss-of-function mutants take on a leaf-like appearance, whereas the inner whorl floral organs of these mutants are partially converted to paleas and lemmas, which are normally found in the outer two whorls of the Poaceae spikelet (Agrawal et al., 2005). These results suggest *OsMADS1* to contribute to the E-function in rice (Fig. 10B), while the functions of four remaining rice *SEP*-clade genes, *OsMADS5*, *OsMADS7*, *OsMADS8* and *RMADS217* (Zahn et al., 2005), remain to be determined. Interestingly, the inactivation of *AGL6*-clade MADS box genes in both rice and maize has also recently been shown to generate E-function-like mutant phenotypes (Li et al., 2010, Ohmori et al., 2009, Thompson et al., 2009). As discussed above, the *AGL6* and *SEP* clades occur in sister positions in MADS box phylogeny, and similar *SEP*-like phenotypes for an *AGL6* orthologue have also recently been demonstrated in *Petunia* (Rijpkema et al., 2009).

In general, carpel and ovule development in the highly derived Poaceae monocots seem to depend on the orthologues of regulatory genes that are known to play key roles in these



processes in *Arabidopsis* and other eudicots. However, specific duplications have taken place in several MADS box lineages in Poaceae, including those of the C-, D- and E-functions, in some cases leading to sub-functionalisation events between paralogous genes. The precise limits of this sub-functionalisation have not yet been defined, which might explain the currently unclear contribution made by MADS box genes to the specification of carpel identity in Poaceae models. Further work is also required in other monocot groups, and perhaps particularly in basal monocots, to establish the extent to which conclusions arising from molecular data of Poaceae models can be applied outside this highly derived family.

#### **E. Carpel development in basal eudicots**

The eudicots form the largest group of angiosperms, and are characterised by the single synapomorphic character of tricolpate pollen (containing three apertures in the pollen wall). Phylogenetic analyses clearly indicate Ranunculales to be sister to all other eudicots (reviewed by Judd and Olmstead, 2004), and the comparison of this and other basally diverging groups provides insight into the structure of the carpel in the last common ancestor of the eudicots (Endress and Doyle, 2009, Endress and Igersheim, 1999). These comparative studies suggest the gynoecium in the ancestral eudicot to have been apocarpic, and to have contained no extragynoecial compitum: a system which allows for the exchange of pollen tubes between the carpels of an apocarpous gynoecium. Its ovules were probably pendant, though no firm conclusions can be made of the number of ovules in each carpel, and on the mechanism of carpel closure (ie. secretion versus postgenital cell division) in the last common ancestor of the eudicots. The presumed ancestral features of the ancestral eudicot gynoecium are illustrated by the Ranunculaceae genus *Aquilegia* (Fig. 10C).

Numerous Ranunculales are being developed as molecular-genetic models, though most of the work on these species has so far been focussed on perianth structure, rather than on the gynoecium. An exception to this is *Eschscholzia californica* (California poppy) of Papaveraceae, though in contrast to the presumed state of the eudicot ancestor, this species possesses a gynoecium of two fused carpels (Fig. 10C). Placentation in *Eschscholzia* is parietal, with two rows of ovules forming in the single loculus of its syncarpic ovary. Molecular studies of *Eschscholzia*, with relevance to the gynoecium, have focussed on orthologues of *LFY/FLORICAULA* (*LFY/FLO*) and of *CRC*, which are respectively termed *EcFLO* and *EcCRC*. *EcFLO* expression was found to be absent from the centre of the *Eschscholzia* floral meristem (Becker et al., 2005), which represents a considerable difference from *Arabidopsis*, in which *LFY* expression is required for the expression of the C-function gene *AG* in the inner floral whorls (Hong et al., 2003). The functional analysis of *EcCRC* in *Eschscholzia* (Orashakova et al., 2009), performed using virus-induced gene silencing (VIGS; Wege et al., 2007), demonstrates a significant overlap of function of this gene with *CRC* in *Arabidopsis* (Alvarez and Smyth, 1999). Accordingly, both *CRC* and its *Eschscholzia* orthologue appear to contribute to abaxial tissue identity in the carpel, floral determinacy, and the growth of tissues that develop from the carpel margins. However, unlike its *Arabidopsis* orthologue, *EcCRC* appears to have acquired novel functions in placenta development and ovule initiation (Orashakova et al., 2009). Further molecular analyses in basal eudicots have focussed on the MADS box family, and comparison of this family in basal and core (later-diverging) eudicots has provided evidence of numerous duplications in MADS box lineages of relevance to carpel development, as discussed in the section below.

#### **F. The role of gene duplications in core eudicot carpel evolution**

The core eudicots form a monophyletic crown group of eudicots. This group includes the major clades of the rosids, asterids and Caryophyllales, which together account for the majority of extant angiosperm species. Several well known molecular-genetic models occur in the core eudicots (Fig. 10D), including *Arabidopsis thaliana* (rosids, Brassicales, Brassicaceae), *Antirrhinum majus* (asterids, Plantaginales, Plantaginaceae) and *Petunia hybrida* (asterids, Solanales, Solanaceae). Comparison of the major clades making up the core eudicots indicates the presence of numerous novel floral features, as compared to the earlier common ancestor of all eudicots. Perhaps the most notable change in carpel structure between these two successive stages in eudicot evolution was the origin of syncarpy, which appears to be a plesiomorphic feature of the core eudicots (Armbruster et al., 2002, Endress and Doyle, 2009).

Analysis of the *Arabidopsis* genome sequence has provided evidence of a large-scale duplication event that probably occurred not long before the last common ancestor of the core eudicots (De Bodt et al., 2005). Accordingly, single genes in basal eudicots are frequently found to be orthologous to pairs of genes in core eudicots, and this is the case for several classes of MADS box genes that control carpel development. For example, two C-clade lineages are present the core eudicots in place of a single *paleoAG* lineage in basally diverging eudicots. Thus in *Arabidopsis*, the *euAG* clade contains the *AG* gene itself, while the *PLENA (PLE)* clade, contains the paralogous genes *SHP1* and *SHP2*, which probably resulted from a further duplication within or near Brassicaceae (Fig. 11). In *Antirrhinum majus*, the probable orthologue of *AG* is termed *FARINELLI (FAR)*, and that of *SHP1/2* is the clade-defining gene *PLE*. Interestingly, the non-orthologous C-clade genes *AG* and *PLE* are responsible for specifying the C-function in *Arabidopsis* and *Antirrhinum*, respectively (Fig. 11; Davies et al., 1999, Kramer et al., 2004). *FAR*, by contrast, is redundantly involved in

stamen development and contributes to pollen fertility. In an example of neo-functionalisation, *SHP1* and *2* have acquired a novel role in *Arabidopsis* fruit development (Liljegren et al., 2000). In *Petunia hybrida*, which, as a member of the asterids, is more closely related to *Antirrhinum* than to *Arabidopsis* (Fig. 9), a further case of sub-functionalisation is to be found. The *Petunia* AG orthologue *PMADS3* is principally responsible for stamen development (Kapoor et al., 2002), though also plays a redundant role with the *PLE* orthologue *FBP6* in both carpel development and floral determinacy (Kramer et al., 2004).

Though sub-functionalisation between the paralogous AG and PLE clades in the *Arabidopsis* lineage has left AG playing the major C-function role, elegant experiments involving multiple mutants demonstrate the SHP genes also to have retained a capacity for C-function activity. Ectopic carpelloid organs are formed in the first floral whorl of *Arabidopsis ag* mutants, conditionally on the inactivation of AP2, which is known mainly for its contribution to the A-function (Bowman et al., 1991b). This effect is thought to occur because, in wild-type *Arabidopsis* plants, AP2 is responsible for down-regulating C-clade genes in the outer floral whorls. In the case of *ag/ap2* double mutants, the C-function activity responsible for specifying ectopic carpel development in the first whorl is provided by *SHP1* and *SHP2*, as demonstrated by the complete lack of carpelloid features to be observed in the first whorl organs of quadruple *ap2/ag/shp1/shp2* mutants (Pinyopich et al., 2003). These data indicate a subtle effect of functional overlap between paralogous gene clades, which does not equate to simple genetic redundancy: the SHP genes adopt a novel C-function role in *Arabidopsis*, conditionally on the inactivation of AP2 and AG.

The fluidity of functions among duplicated genes in the core eudicots is further illustrated by an exchange of function between MADS box genes of the C- and D-clades. Two paralogous D-function genes in *Petunia*, *FBP7* and *FBP11*, are redundantly required for ovule development (Angenent et al., 1995). The *Arabidopsis* orthologue of these genes, *SEEDSTICK* (*STK*), is also involved in ovule development, though *STK* shares this role redundantly with the C-clade genes *SHP1* and *SHP2* (Fig. 11). Accordingly, the *Arabidopsis* *stk/shp1/shp2* triple mutant (Pinyopich et al., 2003), like the *Petunia* *fpb7/fpb11* double mutant (Angenent et al., 1995), produces supernumerary carpels in the place of ovules within the gynoecium. In addition to its redundant role in ovule specification, *STK* plays non-redundant roles in the development of the funiculus and in seed abscission (Pinyopich et al., 2003). The combined C+D-clade in the eudicots, whose different lineages were separated by duplication events that occurred both before and after the radiation of the angiosperms, therefore represents a complex situation in which diverse evolutionary processes have taken place. Examples can be found in this clade of: the repartition of multiple pre-existing functions between paralogous genes (sub-functionalisation), the generation of novel functions associated with one or both genes of a pair of paralogues (neo-functionalisation), and exchanges in function, both between paralogues and non-paralogous genes (Fig. 11).

A further likely consequence of the genome duplication that appears to have occurred at the base of the core eudicots is the generation of a second sub-clade of MADS box genes within the A-clade (Litt and Irish, 2003). The *Arabidopsis* A-function MADS box gene *API* plays roles in floral meristem patterning and in the specification of perianth organ identity. However, gene duplications in the core eudicots have provided further A-clade sequences, of which *FUL* has acquired a role in the patterning of the gynoecium wall in *Arabidopsis* (Gu et al., 1998). *FUL* is known to act in a network involving a large number of genes (Roeder et al.,

2003, Liljegren et al., 2004), including the MADS box genes *SHP1* and *SHP2* (Ferrandiz et al., 2000b) that also function redundantly with *STK* in ovule development. Gene duplication in the MADS box A-clade, followed by neo-functionalisation, has thus resulted in the evolution of novel fruit shattering mechanisms in Brassicaceae.

An interesting feature of gene duplication in the A-clade is the evolution of a distinct C-terminal protein motif in AP1, which was apparently produced by a frame-shift that occurred near the 3'-extremity of the *API* coding sequence in a common ancestor of the core eudicots (Litt and Irish, 2003). This frame-shift created a farnesylation site in the encoded protein, which is known from studies of *Arabidopsis* to be post-translationally modified and required for wild type *API* protein activity (Yalovsky et al., 2000). Other frame-shift mutations in duplicated genes are also present in the B- and C-function MADS box clades in the core eudicots (Vandenbussche et al., 2003), though the motifs generated in these cases are distinct from that of the *API* lineage and do not contain farnesylation sites. The novel C-terminal motifs present in certain lineages within the eudicot A-, B- and C-clades of MADS box genes have been conserved over a long evolutionary timescale, clearly indicating their functional significance. However, it is not known whether the functions of these novel motifs are connected with biochemical processes in common, such as the higher-order assembly of MADS box complexes, or the sequestration of transcription factors through membrane attachment (Vandenbussche et al., 2003).

## **VII. General Conclusions**

In this chapter, our treatment of carpel and gynoecium development has been the most thorough in *Arabidopsis*, reflecting both our own research interests and the relative wealth of data available in this model species. Until recently, most molecular-genetic studies of

gynoecium development have focussed on individual genes and interactions, though the development of large-scale and modelling approaches means that more ambitious goals can now be set. Accordingly, a major challenge for the future will be to construct integrated models of gynoecium development using a system biology approach. A central factor in such an analysis will surely be the hormone auxin. As described in Section IV above, we already know that genes involved in auxin synthesis, transport and responsiveness have many important effects on *Arabidopsis* gynoecium development. A systems biology approach should permit the construction of integrated models of gynoecium development by using a dynamic map of auxin distribution and transport to link together auxin-related genetic elements. Further experimental and modelling approaches can be used to link the many known gynoecium development transcription factors to the proximal causes of development, such as cell division and differentiation.

Though much still remains to be done on *Arabidopsis* carpel development, even more work is required in other species if we are to better understand comparative aspects of carpel development and evolution. Such comparative studies will require in some cases the development of new angiosperm models to fill gaps that currently exist at key phylogenetic positions. The advent of a new generation of sequencing technology (eg. Eid et al., 2009) is expected to revolutionise many aspects of evolutionary-developmental biology, and should provide complete genome sequences from species occupying phylogenetic positions of relevance to flower and carpel evolution (Cyranoski, 2010, Soltis et al., 2008). However, attention must also be paid to the development of functional genetics approaches in new plant models, especially if the expected wealth of genome sequence data is to be exploited to its full potential.

One aspect of carpel development that has only briefly been touched on in this chapter concerns what becomes of the carpel after fertilisation. Indeed, the greater part of carpel-related biodiversity becomes apparent only at this later stage, as the gynoecium transforms into a fruit (Ostergaard, 2010). Fruits are, of course, plant parts of major economic interest, and the contribution of fruits to seed dispersal by air, wind, water, explosion, mammals, birds, reptiles, insects etc (Willson and Traveset, 2002), must also represent one of the major reasons why the flowering plants have been so successful. To accomplish their role in seed dispersal, fruits undergo considerable post-fertilisation changes, though many of the mechanisms that bring these about are established at earlier developmental stages. Thus, the basis for fruit development is laid down during the formation of the carpel (Roeder and Yanofsky, 2005, Sorefan et al., 2009), and involves many of the genetic programs described in Section IV of this chapter. An increasingly important goal of research in this area will be to understand how carpel genetic networks have been modified through evolution to generate the enormous diversity of fruit forms found in nature, and indeed how knowledge of these networks can help us to further modify fruit characteristics to meet our agricultural needs (Doebley et al., 2006, Ostergaard et al., 2006).

### **Acknowledgements**

Our work is supported by: grant BIO2009-09920 from the Spanish *Ministerio de Ciencia e Innovacion* and funding from *Generalitat Valenciana* to C. Ferrandiz; grants BLAN05-0280-01 and BLAN07-2/185566 from the French *Agence Nationale de la Recherche* to C. Scutt and C. Trehin, and funding from the Swedish Research Council Formas to E. Sundberg. We also acknowledge a post of *Attaché Temporaire d'Enseignement et de Recherche* of the *Ecole Normale Supérieure de Lyon* to N. Prunet, and doctoral thesis funding from the French *Ministère de l'Enseignement Supérieure et de la Recherche* to A. Vialette-Guiraud.



## **Figure Legends**

### **Figure 1. Hypotheses for the origin of the flower and its carpel.**

- A. According to the Mostly-Male Theory (Frohlich, 2003), ectopic ovules formed on previously male sporophylls and, in a second step, these sporophylls lost their microsporangia and closed around the ovule to form the carpel. The outer integument of the angiosperm ovule (thick line) was formed from a pre-existing female cupule structure.
- B. According to the Out-of-Male hypothesis (Theissen et al., 2002), the basipetal movement of male-determining B-function MADS box gene expression (shaded area) in a male strobilus left female structures at the apex, which later became carpels.
- C. According to the Out-of-Female hypothesis (Theissen et al., 2002), the acropetal movement of B-sister MADS box gene expression (shaded area) in a female strobilus left male structures at the base, which later became stamens. Female structures at the apex became carpels.
- D. According to the hypothesis of Baum and Hileman (2006), a temporal switch in the regulation by LFY of B- and C-function MADS box genes occurred in an ancestor of the flowering plants. This change generated high concentrations of C-function-rich MADS box complexes at late developmental stages, causing the patterning of the strobilus into apical female and basal male reproductive structures, and these later became carpels and stamens, respectively.

### **Figure 2. Genetic networks controlling plant lateral organ development.**

- A. Interactions between hormones, transcription factors, and RNAs for lateral organ initiation, boundary formation, and establishment of adaxial-abaxial polarity in the leaf.
- B. The establishment of distal-proximal polarity of the leaf.

Figure 3 Structure of the *Arabidopsis* gynoecium.

- A. Scanning electron micrograph of the mature gynoecium at anthesis, artificially coloured to mark different functional domains.
- B. The mature gynoecium at anthesis, cleared using chloral hydrate to reveal vascular patterning. Primary and secondary bifurcations of the medial veins are indicated with arrows.
- C. Cross section of the ovary at anthesis, artificially coloured to mark different functional domain.

Figure 4 Parallelism between genetic networks operating at the shoot apical meristem (SAM) and gynoecium in *Arabidopsis*.

- A. Scheme of the SAM in longitudinal section. Meristem maintenance genes such as *STM* and *RPL* are expressed in meristematic cells (blue), antagonistically with *AS*, *YAB*, *KAN* and *HD-ZIPIII* genes which are expressed in developing leaf primordia (green). Auxin maxima (yellow) precede the emergence of leaf primordia, and boundary genes *CUC* and *KNAT6* mark the outer boundary of the stem cell pool (pink).
- B. Scheme of the *Arabidopsis* gynoecium in transverse section, indicating genetic pathways that direct medio-lateral patterning. Meristem-associated genes *STM*, *RPL* and *BP* are expressed in the medial region at the replum (blue), antagonistically to the

valve-expressed AS, YAB, KAN and HD-ZIPIII genes (green). Boundary genes *KNAT2* and *KNAT6* are expressed at the valve margins (pink), where dehiscence zone genes *SHP* and *IND* also become activated. Auxin maxima occur in the replum, while auxin minima occur at the valve margins.

Figure 5. A genetic model for medio-lateral patterning in the *Arabidopsis* gynoecium and the differentiation of fruit dehiscence zones.

Figure 6. Genetic networks directing marginal tissue differentiation at the apex of the *Arabidopsis* gynoecium.

Black arrows indicate experimentally supported interactions, while grey arrows indicate possible interactions. Boxes indicate putative protein complexes.

Figure 7. Apical-basal patterning in the *Arabidopsis* gynoecium.

An apical-basal auxin gradient has been suggested to participate in the apical-basal patterning of the gynoecium by positioning the borders between the apical style and stigma, the centrally placed ovary, and the short stem or gynophores at the base. After fertilization, the ovary continues to elongate.

Figure 8: Genetic pathways controlling floral meristem (FM) termination.

AG is the main switch toward FM termination, but triggers different developmental programs depending on its protein partners: when AG interacts with B-class proteins AP3 and PI, it promotes stamen development, and without B-class proteins it promotes both carpel development and FM termination. SUP, which excludes B-class gene expression from the

fourth whorl, promotes these two latter functions. In whorl four, AG activates several targets, among which *KNU* plays a central role in switching off *WUS*, and thus stem cell termination. *CRC* and *SPT*, acting downstream of AG, also contribute stem cell termination, showing this process to involve signalling by the developing carpels to the FM.

#### Figure 9. Seed plant phylogeny.

The numbers of species in major clades are given in parentheses, while approximate dates of divergence within the angiosperms, based on a calibration of the molecular clock using fossil data, are estimated from Davies et al. (2004). The positions of some of the taxa referred to in the text are indicated as follows: *Am* = *Amborella trichopoda*, *An* = *Antirrhinum majus*, *Ar* = *Arabidopsis thaliana*, *Ca* = *Cabomba*, *Cr* = *Catharanthus roseus*, *Il* = *Illicium*, *Ze* = *Zea mays*, *Pe* = *Petunia hybrida*, *Or* = *Oryza sativa*.

#### Figure 10. Gynoecium structure and mechanisms specifying carpel development in the angiosperms.

- A. Typical gynoecium structures in ANA grade angiosperms are apocarpic, as in *Amborella trichopoda* and *Cabomba aquatica* (represented schematically in longitudinal section), with carpels that contain one or a few ovules. A secretion filled canal leading from the carpel apex to the ovary provides a route for pollen tube growth. C- and E-clade MADS box gene expression correlates with carpel identity in ANA grade angiosperms, as in core eudicot models. MADS box gene expression is, however, less tightly controlled in ANA grade angiosperms (graded colouring of gene expression boundaries), frequently producing intermediate floral organ types in boundary zones.

- B.** The basal monocot *Acorus calamus* (shown in transverse section) has a trimerous, syncarpic gynoecium, which probably represents the pleisomorphic condition in monocots. However, the gynoecium in the model monocot *Oryza sativa* (rice; shown in longitudinal section), as in other Poaceae, is reduced to a single carpel containing one ovule. Paralogous C-clade MADS box genes show partial sub-functionalisation between the third and fourth whorls of the rice flower: *OsMADS3* plays a major role in stamen development, while *OsMADS58* functions principally in floral determinacy (thick arrows = major roles, thin arrows = minor roles, “STOP” = floral determinacy function). The YAB gene *DL* plays a major role in carpel development either directly (solid arrow), or indirectly by limiting the inner boundary of B-function gene expression (dashed arrow).
- C.** Basal eudicots of Ranunculales include both apocarpic and syncarpic taxa, such as *Aquilegia* (shown in longitudinal section) and *Eschscholzia californica* (shown in transverse section), respectively, though apocarpy is believed to be the pleisomorphic condition in eudicots.
- D.** The last common ancestor of the core eudicots is believed to have possessed a dimerous, syncarpic gynoecium, as is the case in its present-day model plant descendents *Arabidopsis thaliana*, *Petunia hybrida* and *Antirrhinum majus* (all shown in transverse section). However, the dimerous gynoecium of *Arabidopsis* is divided into two loculi by the secondary development of a false septum, whereas those of *Petunia* and *Antirrhinum* are divided by the common wall of their congenitally fused carpels. Carpel and stamen identity, and floral determinism (“STOP”), are controlled by MADS box C-clade genes, though sub-functionalisation has occurred to differently partition these roles among C-clade paralogues in *Arabidopsis* (*AG*), *Petunia* (*PMADS3* and *FBP6*) and *Antirrhinum* (*FAR* and *PLE*).

(ca = carpels, io = intermediate organs, it = inner tepals, ot = outer tepals, pa/le = paleas/lemmas, pe = petals, se = sepals, st = stamens)

Figure 11. Fluidity in the functionalisation of C- and D-function MADS box genes in core eudicots.

**A.-C.** Venn diagrams representing the functions of genes from the *AG*, *PLE* and *FBP7* MADS box clades in three species of core eudicots. Overlapping regions represent functional redundancy between genes in wild-type genetic backgrounds

**D.** The sequence of duplications that generated of the eudicot *AG*, *PLE* and *FBP7* MADS box gene clades.

## References

- Adenot, X., Elmayan, T., Laussergues, D., Boutet, S., Bouche, N., Gascioli, V. & Vaucheret, H. (2006) DRB4-dependent TAS3 trans-acting siRNAs control leaf morphology through AGO7. *Current Biology*, 16, 927-932.
- Agrawal, G., Abe, K., Yamazaki, M., Miyao, A. & Hirochika, H. (2005) Conservation of the E-function for floral organ identity in rice revealed by the analysis of tissue culture-induced loss-of-function mutants of the OsMADS1 gene. *Plant Molecular Biology*, 59, 125-135.
- Aida, M. & Tasaka, M. (2006) Morphogenesis and patterning at the organ boundaries in the higher plant shoot apex u. *Plant Molecular Biology*, 60, 915-928.
- Aloni, R., Schwalm, K., Langhans, M. & Ullrich, C. I. (2003) Gradual shifts in sites of free-auxin production during leaf-primordium development and their role in vascular differentiation and leaf morphogenesis in Arabidopsis. *Planta*, 216, 841-853.
- Alonso-Cantabrana, H., Ripoll, J. J., Ochando, I., Vera, A., Ferrandiz, C. & Martinez-Laborda, A. (2007) Common regulatory networks in leaf and fruit patterning revealed by mutations in the Arabidopsis ASYMMETRIC LEAVES1 gene. *Development*, 134, 2663-2671.
- Alvarez, J. & Smyth, D. R. (1999) CRABS CLAW and SPATULA, two Arabidopsis genes that control carpel development in parallel with AGAMOUS. *Development*, 126, 2377-2386.
- Alvarez, J. P., Goldshmidt, A., Efroni, I., Bowman, J. L. & Eshed, Y. (2009) The NGATHA Distal Organ Development Genes Are Essential for Style Specification in Arabidopsis. *Plant Cell*, 21, 1373-1393.

- Alvarez, J. P., Pekker, I., Goldshmidt, A., Blum, E., Amsellem, Z. & Eshed, Y. (2006) Endogenous and synthetic microRNAs stimulate simultaneous, efficient, and localized regulation of multiple targets in diverse species. *Plant Cell*, 18, 1134-1151.
- Anastasiou, E. & Lenhard, M. (2007) Growing up to one's standard. *Current Opinion in Plant Biology*, 10, 63-69.
- Angenent, G. C., Franken, J., Busscher, M., Vandijken, A., Vanwent, J. L., Dons, H. J. M. & Vantunen, A. J. (1995) A Novel Class of Mads Box Genes Is Involved in Ovule Development in Petunia. *Plant Cell*, 7, 1569-1582.
- Armbruster, W. S., Debevec, E. M. & Willson, M. F. (2002) Evolution of syncarpy in angiosperms: theoretical and phylogenetic analyses of the effects of carpel fusion on offspring quantity and quality. *Journal of Evolutionary Biology*, 15, 657-672.
- Aukerman, M. J. & Sakai, H. (2003) Regulation of flowering time and floral organ identity by a microRNA and its APETALA2-like target genes. *Plant Cell*, 15, 2730-2741.
- Azhakanandam, S., Nole-Wilson, S., Bao, F. & Franks, R. G. (2008) SEUSS and AINTEGUMENTA mediate patterning and ovule initiation during gynoecium medial domain development. *Plant Physiology*, 146, 1165-1181.
- Bainbridge, K., Guyomarc'h, S., Bayer, E., Swarup, R., Bennett, M., Mandel, T. & Kuhlemeier, C. (2008) Auxin influx carriers stabilize phyllotactic patterning. *Genes & Development*, 22, 810-823.
- Balanza, V., Navarrete, M., Trigueros, M. & Ferrandiz, C. (2006) Patterning the female side of Arabidopsis: the importance of hormones. *Journal of Experimental Botany*, 57, 3457-3469.



- Bao, X. Z., Franks, R. G., Levin, J. Z. & Liu, Z. C. (2004) Repression of AGAMOUS by BELLRINGER in floral and inflorescence meristems. *Plant Cell*, 16, 1478-1489.
- Barkoulas, M., Galinha, C., Grigg, S. P. & Tsiantis, M. (2007) From genes to shape: regulatory interactions in leaf development. *Current Opinion in Plant Biology*, 10, 660-666.
- Baum, D. A. & Hileman, L. C. (2006) A developmental genetic model for the origin of the flower. IN AINSWORTH, C. (Ed.) *Flowering and its Manipulation*. Oxford, Blackwell.
- Becker, A., Gleissberg, S. & Smyth, D. R. (2005) Floral and vegetative morphogenesis in California poppy (*Eschscholzia californica* Cham.). *International Journal of Plant Sciences*, 166, 537-555.
- Becker, A. & Theissen, G. (2003) The major clades of MADS-box genes and their role in the development and evolution of flowering plants. *Molecular Phylogenetics and Evolution*, 29, 464-489.
- Becker, A., Winter, K. U., Meyer, B., Saedler, H. & Theissen, G. (2000) MADS-box gene diversity in seed plants 300 million years ago. *Molecular Biology and Evolution*, 17, 1425-1434.
- Benkova, E., Michniewicz, M., Sauer, M., Teichmann, T., Seifertova, D., Jurgens, G. & Friml, J. (2003) Local, efflux-dependent auxin gradients as a common module for plant organ formation. *Cell*, 115, 591-602.
- Bennett, S. R. M., Alvarez, J., Bossinger, G. & Smyth, D. R. (1995) MORPHOGENESIS IN PINOID MUTANTS OF ARABIDOPSIS-THALIANA. *Plant Journal*, 8, 505-520.
- Berardini, T. Z., Bollman, K., Sun, H. & Poethig, R. S. (2001) Regulation of vegetative phase change in *Arabidopsis thaliana* by cyclophilin 40. *Science*, 291, 2405-2407.

- Bereterbide, A., Hernould, M., Castera, S. & Mouras, A. (2001) Inhibition of cell proliferation, cell expansion and differentiation by the Arabidopsis SUPERMAN gene in transgenic tobacco plants. *Planta*, 214, 22-29.
- Bernhardt, P., Sage, T., Weston, P., Azuma, H., Lam, M., Thien, L. B. & Bruhl, J. (2003) The pollination of *Trimenia moorei* (Trimeniaceae): Floral volatiles, insect/wind pollen vectors and stigmatic self-incompatibility in a basal angiosperm. *Annals of Botany*, 92, 445-458.
- Bolduc, N. & Hake, S. (2009) The Maize Transcription Factor KNOTTED1 Directly Regulates the Gibberellin Catabolism Gene *ga2ox1*. *Plant Cell*, 21, 1647-1658.
- Borghi, L., Bureau, M. & Simon, R. (2007) Arabidopsis JAGGED LATERAL ORGANS is expressed in boundaries and coordinates KNOX and PIN activity. *Plant Cell*, 19, 1795-1808.
- Bowman, J. L. (1991). *Molecular Genetics of Flower Development in Arabidopsis thaliana*. Pasadena, California Institute of Technology Press.
- Bowman, J. L. (2000) The YABBY gene family and abaxial cell fate. *Current Opinion in Plant Biology*, 3, 17-22.
- Bowman, J. L., Baum, S. F., Eshed, Y., Putterill, J. & Alvarez, J. (1999) Molecular genetics of gynoecium development in Arabidopsis. *Current Topics in Developmental Biology*, Vol 45.
- Bowman, J. L., Drews, G. N. & Meyerowitz, E. M. (1991a) Expression of the Arabidopsis Floral Homeotic Gene *Agamous* Is Restricted to Specific Cell-Types Late in Flower Development. *Plant Cell*, 3, 749-758.
- Bowman, J. L. & Floyd, S. K. (2008) Patterning and polarity in seed plant shoots. *Annual Review of Plant Biology*, 59, 67-88.

- Bowman, J. L., Sakai, H., Jack, T., Weigel, D., Mayer, U. & Meyerowitz, E. M. (1992) Superman, a Regulator of Floral Homeotic Genes in Arabidopsis. *Development*, 114, 599-&.
- Bowman, J. L. & Smyth, D. R. (1999) CRABS CLAW, a gene that regulates carpel and nectary development in Arabidopsis, encodes a novel protein with zinc finger and helix-loop-helix domains. *Development*, 126, 2387-2396.
- Bowman, J. L., Smyth, D. R. & Meyerowitz, E. M. (1991b) Genetic Interactions among Floral Homeotic Genes of Arabidopsis. *Development*, 112, 1-20.
- Byrne, M. E. (2005) Networks in leaf development. *Current Opinion in Plant Biology*, 8, 59-66.
- Byrne, M. E. (2006) Shoot meristem function and leaf polarity: The role of class IIIHD-ZIP genes. *Plos Genetics*, 2, 785-790.
- Canales, C., Barkoulas, M., Galinha, C. & Tsiantis, M. (2010) Weeds of change: Cardamine hirsuta as a new model system for studying dissected leaf development. *Journal of Plant Research*, 123, 25-33.
- Carles, C. C., Lertpiriyapong, K., Reville, K. & Fletcher, J. C. (2004) The ULTRAPETALA1 gene functions early in Arabidopsis development to restrict shoot apical meristem activity and acts through WUSCHEL to regulate floral meirstem determinacy. *Genetics*, 167, 1893-1903.
- Carlsbecker, A., Tandre, K., Johanson, U., Englund, M. & Engstrom, P. (2004) The MADS-box gene DAL1 is a potential mediator of the juvenile-to-adult transition in Norway spruce (Picea abies). *Plant Journal*, 40, 546-557.
- Castillejo, C., Romera-Branchat, M. & Pelaz, S. (2005) A new role of the Arabidopsis SEPALLATA3 gene revealed by its constitutive expression. *Plant Journal*, 43, 586-596.

- Chen, C. B., Wang, S. P. & Huang, H. (2000) LEUNIG has multiple functions in gynoecium development in Arabidopsis. *Genesis*, 26, 42-54.
- Chen, H., Banerjee, A. K. & Hannapel, D. J. (2004) The tandem complex of BEL and KNOX partners is required for transcriptional repression of *ga20ox1*. *Plant Journal*, 38, 276-284.
- Chen, Q. Y., Atkinson, A., Otsuga, D., Christensen, T., Reynolds, L. & Drews, G. N. (1999) The Arabidopsis FILAMENTOUS FLOWER gene is required for flower formation. *Development*, 126, 2715-2726.
- Chen, X. M., Liu, J., Cheng, Y. L. & Jia, D. X. (2002) HEN1 functions pleiotropically in Arabidopsis development and acts in C function in the flower. *Development*, 129, 1085-1094.
- Cheng, Y. F., Dai, X. H. & Zhao, Y. D. (2006) Auxin biosynthesis by the YUCCA flavin monooxygenases controls the formation of floral organs and vascular tissues in Arabidopsis. *Genes & Development*, 20, 1790-1799.
- Cheng, Y. F., Dai, X. H. & Zhao, Y. D. (2007) Auxin synthesized by the YUCCA flavin Monooxygenases is essential for embryogenesis and leaf formation in Arabidopsis. *Plant Cell*, 19, 2430-2439.
- Cheng, Y. L., Kato, N., Wang, W. M., Li, J. J. & Chen, X. M. (2003) Two RNA binding proteins, HEN4 and HUM, act in the processing of AGAMOUS Pre-mRNA in Arabidopsis thaliana. *Developmental Cell*, 4, 53-66.
- Chitwood, D. H., Guo, M. J., Nogueira, F. T. S. & Timmermans, M. C. P. (2007) Establishing leaf polarity: the role of small RNAs and positional signals in the shoot apex. *Development*, 134, 813-823.

- Chitwood, D. H., Nogueira, F. T. S., Howell, M. D., Montgomery, T. A., Carrington, J. C. & Timmermans, M. C. P. (2009) Pattern formation via small RNA mobility. *Genes & Development*, 23, 549-554.
- Chuang, C. F. & Meyerowitz, E. M. (2000) Specific and heritable genetic interference by double-stranded RNA in *Arabidopsis thaliana*. *Proceedings of the National Academy of Sciences of the United States of America*, 97, 4985-4990.
- Clark, S. E., Running, M. P. & Meyerowitz, E. M. (1993) Clavata1, a Regulator of Meristem and Flower Development in *Arabidopsis*. *Development*, 119, 397-418.
- Clark, S. E., Running, M. P. & Meyerowitz, E. M. (1995) Clavata3 Is a Specific Regulator of Shoot and Floral Meristem Development Affecting the Same Processes as Clavata1. *Development*, 121, 2057-2067.
- Coen, E. S. & Meyerowitz, E. M. (1991) The War of the Whorls - Genetic Interactions Controlling Flower Development. *Nature*, 353, 31-37.
- Colombo, L., Franken, J., Koetje, E., Vanwent, J., Dons, H. J. M., Angenent, G. C. & Vantunen, A. J. (1995) The *Petunia* Mads Box Gene *Fbp11* Determines Ovule Identity. *Plant Cell*, 7, 1859-1868.
- Colombo, M., Brambilla, V., Marcheselli, R., Caporali, E., Kater, M. M. & Colombo, L. (2010) A new role for the SHATTERPROOF genes during *Arabidopsis* gynoecium development. *Developmental Biology*, 337, 294-302.
- Crawford, B. C. W., Ditta, G. & Yanofsky, M. F. (2007) The NTT gene is required for transmitting-tract development in carpels of *Arabidopsis thaliana*. *Current Biology*, 17, 1101-1108.
- Cyranoski, D. (2010) The Sequence Factory. *Nature*, 464, 22-24.

- Das, P., Ito, T., Wellmer, F., Vernoux, T., Dedieu, A., Traas, J. & Meyerowitz, E. M. (2009) Floral stem cell termination involves the direct regulation of AGAMOUS by PERIANTHIA. *Development*, 136, 1605-1611.
- Davies, B., Motte, P., Keck, E., Saedler, H., Sommer, H. & Schwarz-Sommer, Z. (1999) PLENA and FARINELLI: redundancy and regulatory interactions between two Antirrhinum MADS-box factors controlling flower development. *Embo Journal*, 18, 4023-4034.
- Davies, T. J., Barraclough, T. G., Chase, M. W., Soltis, P. S., Soltis, D. E. & Savolainen, V. (2004) Darwin's abominable mystery: Insights from a supertree of the angiosperms. *Proceedings of the National Academy of Sciences of the United States of America*, 101, 1904-1909.
- De Bodt, S., Maere, S. & Van de Peer, Y. (2005) Genome duplication and the origin of angiosperms. *Trends in Ecology & Evolution*, 20, 591-597.
- de Folter, S., Shchennikova, A. V., Franken, J., Busscher, M., Baskar, R., Grossniklaus, U., Angenent, G. C. & Immink, R. G. H. (2006) A B-sister MADS-box gene involved in ovule and seed development in petunia and Arabidopsis. *Plant Journal*, 47, 934-946.
- Dinneny, J. R., Weigel, D. & Yanofsky, M. F. (2005) A genetic framework for fruit patterning in Arabidopsis thaliana. *Development*, 132, 4687-4696.
- Dinneny, J. R., Weigel, D. & Yanofsky, M. F. (2006) NUBBIN and JAGGED define stamen and carpel shape in Arabidopsis. *Development*, 133, 1645-1655.
- Dinneny, J. R., Yadegari, R., Fischer, R. L., Yanofsky, M. F. & Weigel, D. (2004) The role of JAGGED in shaping lateral organs. *Development*, 131, 1101-1110.
- Dinneny, J. R. & Yanofsky, M. F. (2005) Drawing lines and borders: how the dehiscent fruit of Arabidopsis is patterned. *Bioessays*, 27, 42-49.

- Ditta, G., Pinyopich, A., Robles, P., Pelaz, S. & Yanofsky, M. F. (2004) The SEP4 gene of *Arabidopsis thaliana* functions in floral organ and meristem identity. *Current Biology*, 14, 1935-1940.
- Doebley, J. F., Gaut, B. S. & Smith, B. D. (2006) The molecular genetics of crop domestication. *Cell*, 127, 1309-1321.
- Donnelly, P. M., Bonetta, D., Tsukaya, H., Dengler, R. E. & Dengler, N. G. (1999) Cell cycling and cell enlargement in developing leaves of *Arabidopsis*. *Developmental Biology*, 215, 407-419.
- Dorcey, E., Urbez, C., Blazquez, M. A., Carbonell, J. & Perez-Amador, M. A. (2009) Fertilization-dependent auxin response in ovules triggers fruit development through the modulation of gibberellin metabolism in *Arabidopsis*. *Plant Journal*, 58, 318-332.
- Dornelas, M. C. & Rodriguez, A. P. M. (2005) A FLORICAULA/LEAFY gene homolog is preferentially expressed in developing female cones of the tropical pine *Pinus caribaea* var. *caribaea*. *Genetics and Molecular Biology*, 28, 299-307.
- Dreni, L., Jacchia, S., Fornara, F., Fornari, M., Ouwerkerk, P. B. F., An, G. H., Colombo, L. & Kater, M. M. (2007) The D-lineage MADS-box gene OsMADS13 controls ovule identity in rice. *Plant Journal*, 52, 690-699.
- Drews, G. N., Bowman, J. L. & Meyerowitz, E. M. (1991) Negative Regulation of the *Arabidopsis* Homeotic Gene *Agamous* by the *Apetala2* Product. *Cell*, 65, 991-1002.
- Efroni, I., Blum, E., Goldshmidt, A. & Eshed, Y. (2008) A Protracted and Dynamic Maturation Schedule Underlies *Arabidopsis* Leaf Development. *Plant Cell*, 20, 2293-2306.

- Eid, J., Fehr, A., Gray, J., Luong, K., Lyle, J., Otto, G., Peluso, P., Rank, D., Baybayan, P., Bettman, B., Bibillo, A., Bjornson, K., Chaudhuri, B., Christians, F., Cicero, R., Clark, S., Dalal, R., Dewinter, A., Dixon, J., Foquet, M., Gaertner, A., Hardenbol, P., Heiner, C., Hester, K., Holden, D., Kearns, G., Kong, X. X., Kuse, R., Lacroix, Y., Lin, S., Lundquist, P., Ma, C. C., Marks, P., Maxham, M., Murphy, D., Park, I., Pham, T., Phillips, M., Roy, J., Sebra, R., Shen, G., Sorenson, J., Tomaney, A., Travers, K., Trulson, M., Vieceli, J., Wegener, J., Wu, D., Yang, A., Zaccarin, D., Zhao, P., Zhong, F., Korlach, J. & Turner, S. (2009) Real-Time DNA Sequencing from Single Polymerase Molecules. *Science*, 323, 133-138.
- Eklund, D. M., Staldal, V., Valsecchi, I., Cierlik, I., Eriksson, C., Hiratsu, K., Ohme-Takagi, M., Sundstrom, J. F., Thelander, M., Ezcurra, I. & Sundberg, E. (2010) The Arabidopsis thaliana STYLISH1 Protein Acts as a Transcriptional Activator Regulating Auxin Biosynthesis. *Plant Cell*, 22, 349-363.
- Elliott, R. C., Betzner, A. S., Huttner, E., Oakes, M. P., Tucker, W. Q. J., Gerentes, D., Perez, P. & Smyth, D. R. (1996) AINTEGUMENTA, an APETALA2-like gene of arabidopsis with pleiotropic roles in ovule development and floral organ growth. *Plant Cell*, 8, 155-168.
- Emery, J. F., Floyd, S. K., Alvarez, J., Eshed, Y., Hawker, N. P., Izhaki, A., Baum, S. F. & Bowman, J. L. (2003) Radial patterning of Arabidopsis shoots by class III HD-ZIP and KANADI genes. *Current Biology*, 13, 1768-1774.
- Endress, P. K. (1982) Syncarpy and Alternative Modes of Escaping Disadvantages of Apocarpy in Primitive Angiosperms. *Taxon*, 31, 48-52.
- Endress, P. K. (1986) Reproductive Structures and Phylogenetic Significance of Extant Primitive Angiosperms. *Plant Systematics and Evolution*, 152, 1-28.



- Endress, P. K. (2001) The flowers in extant basal angiosperms and inferences on ancestral flowers. *International Journal of Plant Sciences*, 162, 1111-1140.
- Endress, P. K. (2005) Carpels in *Brasenia* (Cabombaceae) are completely ascidiate despite a long stigmatic crest. *Annals of Botany*, 96, 209-215.
- Endress, P. K. & Doyle, J. A. (2009) Reconstructing the Ancestral Angiosperm Flower and Its Initial Specializations. *American Journal of Botany*, 96, 22-66.
- Endress, P. K. & Igersheim, A. (1999) Gynoecium diversity and systematics of the basal eudicots. *Botanical Journal of the Linnean Society*, 130, 305-393.
- Endress, P. K. & Igersheim, A. (2000) Gynoecium structure and evolution in basal angiosperms. *International Journal of Plant Sciences*, 161, S211-S223.
- Eshed, Y., Baum, S. F. & Bowman, J. L. (1999) Distinct mechanisms promote polarity establishment in carpels of *Arabidopsis*. *Cell*, 99, 199-209.
- Eshed, Y., Baum, S. F., Perea, J. V. & Bowman, J. L. (2001) Establishment of polarity in lateral organs of plants. *Current Biology*, 11, 1251-1260.
- Eshed, Y., Izhaki, A., Baum, S. F., Floyd, S. K. & Bowman, J. L. (2004) Asymmetric leaf development and blade expansion in *Arabidopsis* are mediated by KANADI and YABBY activities. *Development*, 131, 2997-3006.
- Fahlgren, N., Montgomery, T. A., Howell, M. D., Allen, E., Dvorak, S. K., Alexander, A. L. & Carrington, J. C. (2006) Regulation of AUXIN RESPONSE FACTOR3 by TAS3 ta-siRNA affects developmental timing and patterning in *Arabidopsis*. *Current Biology*, 16, 939-944.
- Fahn, A. (1982) *Plant Anatomy*, Oxford, Pergamon.
- Ferrandiz, C. (2002) Regulation of fruit dehiscence in *Arabidopsis*. *Journal of Experimental Botany*, 53, 2031-2038.

- Ferrandiz, C., Gu, Q., Martienssen, R. & Yanofsky, M. F. (2000a) Redundant regulation of meristem identity and plant architecture by FRUITFULL, APETALA1 and CAULIFLOWER. *Development*, 127, 725-734.
- Ferrandiz, C., Liljegren, S. J. & Yanofsky, M. F. (2000b) Negative regulation of the SHATTERPROOF genes by FRUITFULL during Arabidopsis fruit development. *Science*, 289, 436-438.
- Flanagan, C. A., Hu, Y. & Ma, H. (1996) Specific expression of the AGL1 MADS-box gene suggests regulatory functions in Arabidopsis gynoecium and ovule development. *Plant Journal*, 10, 343-353.
- Fletcher, J. C. (2001) The ULTRAPETALA gene controls shoot and floral meristem size in Arabidopsis. *Development*, 128, 1323-1333.
- Fourquin, C., Vinauger-Douard, M., Chambrier, P., Berne-Dedieu, A. & Scutt, C. P. (2007) Functional conservation between CRABS CLAW orthologues from widely diverged Angiosperms. *Annals of Botany*, 100, 651-657.
- Fourquin, C., Vinauger-Douard, M., Fogliani, B., Dumas, C. & Scutt, C. P. (2005) Evidence that CRABS CLAW and TOUSLED have conserved their roles in carpel development since the ancestor of the extant angiosperms. *Proceedings of the National Academy of Sciences of the United States of America*, 102, 4649-4654.
- Franks, R. G., Wang, C. X., Levin, J. Z. & Liu, Z. C. (2002) SEUSS, a member of a novel family of plant regulatory proteins, represses floral homeotic gene expression with LEUNIG. *Development*, 129, 253-263.
- Friedman, W. E. (2006) Embryological evidence for developmental lability during early angiosperm evolution. *Nature*, 441, 337-340.
- Friedman, W. E. (2008) Hydatellaceae are water lilies with gymnospermous tendencies. *Nature*, 453, 94-97.

- Friedman, W. E. (2009) The Meaning of Darwin's "Abominable Mystery". *American Journal of Botany*, 96, 5-21.
- Friis, E. M., Pedersen, K. R. & Crane, P. R. (2001) Fossil evidence of water lilies (Nymphaeales) in the Early Cretaceous. *Nature*, 410, 357-360.
- Friis, E. M., Pedersen, K. R. & Crane, P. R. (2010) Diversity in obscurity: fossil flowers and the early history of angiosperms. *Philosophical Transactions of the Royal Society B-Biological Sciences*, 365, 369-382.
- Frohlich, M. W. (2003) An evolutionary scenario for the origin of flowers. *Nature Reviews Genetics*, 4, 559-566.
- Frohlich, M. W. & Chase, M. W. (2007) After a dozen years of progress the origin of angiosperms is still a great mystery. *Nature*, 450, 1184-1189.
- Frohlich, M. W. & Parker, D. S. (2000) The mostly male theory of flower evolutionary origins: from genes to fossils. *Systematic Botany*, 25, 155-170.
- Fu, Y. L., Xu, L., Xu, B., Yang, L., Ling, Q. H., Wang, H. & Huang, H. (2007) Genetic interactions between leaf polarity-controlling genes and ASYMMETRIC LEAVES1 and 2 in Arabidopsis leaf patterning. *Plant and Cell Physiology*, 48, 724-735.
- Garcia, D., Collier, S. A., Byrne, M. E. & Martienssen, R. A. (2006) Specification of leaf polarity in Arabidopsis via the trans-acting siRNA pathway. *Current Biology*, 16, 933-938.
- Girin, T., Sorefan, K. & Ostergaard, L. (2009) Meristematic sculpting in fruit development. *Journal of Experimental Botany*, 60, 1493-1502.
- Goetz, M., Vivian-Smith, A., Johnson, S. D. & Koltunow, A. M. (2006) AUXIN RESPONSE FACTOR8 is a negative regulator of fruit initiation in Arabidopsis. *Plant Cell*, 18, 1873-1886.

- Gomez-Mena, C., de Folter, S., Costa, M. M. R., Angenent, G. C. & Sablowski, R. (2005) Transcriptional program controlled by the floral homeotic gene AGAMOUS during early organogenesis. *Development*, 132, 429-438.
- Gomez-Mena, C. & Sablowski, R. (2008) ARABIDOPSIS THALIANA HOMEBOX GENE1 establishes the basal boundaries of shoot organs and controls stem growth. *Plant Cell*, 20, 2059-2072.
- Gordon, S. P., Chickarmane, V. S., Ohno, C. & Meyerowitz, E. M. (2009) Multiple feedback loops through cytokinin signaling control stem cell number within the Arabidopsis shoot meristem. *Proceedings of the National Academy of Sciences of the United States of America*, 106, 16529-16534.
- Goremykin, V. V., Hansmann, S. & Martin, W. F. (1997) Evolutionary analysis of 58 proteins encoded in six completely sequenced chloroplast genomes: Revised molecular estimates of two seed plant divergence times. *Plant Systematics and Evolution*, 206, 337-351.
- Goto, K., Kyojuka, J. & Bowman, J. L. (2001) Turning floral organs into leaves, leaves into floral organs. *Current Opinion in Genetics & Development*, 11, 449-456.
- Goto, K. & Meyerowitz, E. M. (1994) Function and Regulation of the Arabidopsis Floral Homeotic Gene Pistillata. *Genes & Development*, 8, 1548-1560.
- Gregis, V., Sessa, A., Colombo, L. & Kater, M. M. (2006) AGL24, SHORT VEGETATIVE PHASE, and APETALA1 redundantly control AGAMOUS during early stages of flower development in Arabidopsis. *Plant Cell*, 18, 1373-1382.
- Gremski, K., Ditta, G. & Yanofsky, M. F. (2007) The HECATE genes regulate female reproductive tract development in Arabidopsis thaliana. *Development*, 134, 3593-3601.

- Groszmann, M., Paicu, T. & Smyth, D. R. (2008) Functional domains of SPATULA, a bHLH transcription factor involved in carpel and fruit development in Arabidopsis. *Plant Journal*, 55, 40-52.
- Gu, Q., Ferrandiz, C., Yanofsky, M. F. & Martienssen, R. (1998) The FRUITFULL MADS-box gene mediates cell differentiation during Arabidopsis fruit development. *Development*, 125, 1509-1517.
- Guo, M. J., Thomas, J., Collins, G. & Timmermans, M. C. P. (2008) Direct repression of KNOX loci by the ASYMMETRIC LEAVES1 complex of Arabidopsis. *Plant Cell*, 20, 48-58.
- Ha, C. M., Jun, J. H., Nam, H. G. & Fletcher, J. C. (2004) BLADE-ON-PETIOLE1 encodes a BTB/POZ domain protein required for leaf morphogenesis in Arabidopsis thaliana. *Plant and Cell Physiology*, 45, 1361-1370.
- Ha, C. M., Jun, J. H., Nam, H. G. & Fletcher, J. C. (2007) BLADE-ON-PETIOLE1 and 2 control Arabidopsis lateral organ fate through regulation of LOB domain and adaxial-abaxial polarity genes. *Plant Cell*, 19, 1809-1825.
- Ha, C. M., Kim, G. T., Kim, B. C., Jun, J. H., Soh, M. S., Ueno, Y., Machida, Y., Tsukaya, H. & Nam, H. G. (2003) The BLADE-ON-PETIOLE 1 gene controls leaf pattern formation through the modulation of meristematic activity in Arabidopsis. *Development*, 130, 161-172.
- Hake, S., Smith, H. M. S., Holtan, H., Magnani, E., Mele, G. & Ramirez, J. (2004) The role of knox genes in plant development. *Annual Review of Cell and Developmental Biology*, 20, 125-151.
- Hay, A., Barkoulas, M. & Tsiantis, M. (2006) ASYMMETRIC LEAVES1 and auxin activities converge to repress BREVIPEDICELLUS expression and promote leaf development in Arabidopsis. *Development*, 133, 3955-3961.

- Hay, A., Kaur, H., Phillips, A., Hedden, P., Hake, S. & Tsiantis, M. (2002) The gibberellin pathway mediates KNOTTED1-type homeobox function in plants with different body plans. *Current Biology*, 12, 1557-1565.
- Heisler, M. G., Ohno, C., Das, P., Sieber, P., Reddy, G. V., Long, J. A. & Meyerowitz, E. M. (2005) Patterns of auxin transport and gene expression during primordium development revealed by live imaging of the Arabidopsis inflorescence meristem. *Current Biology*, 15, 1899-1911.
- Heisler, M. G. B., Atkinson, A., Bylstra, Y. H., Walsh, R. & Smyth, D. R. (2001) SPATULA, a gene that controls development of carpel margin tissues in Arabidopsis, encodes a bHLH protein. *Development*, 128, 1089-1098.
- Hempel, F. D., Weigel, D., Mandel, M. A., Ditta, G., Zambryski, P. C., Feldman, L. J. & Yanofsky, M. F. (1997) Floral determination and expression of floral regulatory genes in Arabidopsis. *Development*, 124, 3845-3853.
- Hepworth, S. R., Zhang, Y. L., McKim, S., Li, X. & Haughn, G. (2005) BLADE-ON-PETIOLE-dependent signaling controls leaf and floral patterning in Arabidopsis. *Plant Cell*, 17, 1434-1448.
- Hibara, K., Karim, M. R., Takada, S., Taoka, K. I., Furutani, M., Aida, M. & Tasaka, M. (2006) Arabidopsis CUP-SHAPED COTYLEDON3 regulates postembryonic shoot meristem and organ boundary formation. *Plant Cell*, 18, 2946-2957.
- Hiratsu, K., Ohta, M., Matsui, K. & Ohme-Takagi, M. (2002) The SUPERMAN protein is an active repressor whose carboxy-terminal repression domain is required for the development of normal flowers. *Febs Letters*, 514, 351-354.
- Hong, R. L., Hamaguchi, L., Busch, M. A. & Weigel, D. (2003) Regulatory elements of the floral homeotic gene AGAMOUS identified by phylogenetic footprinting and shadowing. *Plant Cell*, 15, 1296-1309.

- Honma, T. & Goto, K. (2001) Complexes of MADS-box proteins are sufficient to convert leaves into floral organs. *Nature*, 409, 525-529.
- Horiguchi, G., Kim, G. T. & Tsukaya, H. (2005) The transcription factor AtGRF5 and the transcription coactivator AN3 regulate cell proliferation in leaf primordia of *Arabidopsis thaliana*. *Plant Journal*, 43, 68-78.
- Hu, Y. X., Xie, O. & Chua, N. H. (2003) The *Arabidopsis* auxin-inducible gene ARGOS controls lateral organ size. *Plant Cell*, 15, 1951-1961.
- Hunter, C., Willmann, M. R., Wu, G., Yoshikawa, M., de la Luz Gutierrez-Nava, M. & Poethig, R. S. (2006) Trans-acting siRNA-mediated repression of ETTIN and ARF4 regulates heteroblasty in *Arabidopsis*. *Development*, 133, 2973-2981.
- Husbands, A., Bell, E. M., Shuai, B., Smith, H. M. S. & Springer, P. S. (2007) LATERAL ORGAN BOUNDARIES defines a new family of DNA-binding transcription factors and can interact with specific bHLH proteins. *Nucleic Acids Research*, 35, 6663-6671.
- Husbands, A. Y., Chitwood, D. H., Plavskin, Y. & Timmermans, M. C. P. (2009) Signals and prepatterns: new insights into organ polarity in plants. *Genes & Development*, 23, 1986-1997.
- Ichihashi, Y., Horiguchi, G., Gleissberg, S. & Tsukaya, H. (2010) The bHLH Transcription Factor SPATULA Controls Final Leaf Size in *Arabidopsis thaliana*. *Plant and Cell Physiology*, 51, 252-261.
- Igersheim, A., Buzgo, M. & Endress, P. K. (2001) Gynoecium diversity and systematics in basal monocots. *Botanical Journal of the Linnean Society*, 136, 1-65.
- Ikezaki, M., Kojima, M., Sakakibara, H., Kojima, S., Ueno, Y., Machida, C. & Machida, Y. (2010) Genetic networks regulated by ASYMMETRIC LEAVES1 (AS1) and

- AS2 in leaf development in *Arabidopsis thaliana*: KNOX genes control five morphological events. *Plant Journal*, 61, 70-82.**
- Ilegems, M., Douet, V., Meylan-Bettex, M., Uyttewaal, M., Brand, L., Bowman, J. L. & Stieger, P. A. (2010) Interplay of auxin, KANADI and Class III HD-ZIP transcription factors in vascular tissue formation. *Development*, 137, 975-984.**
- Ingram, G. C. & Waites, R. (2006) Keeping it together: co-ordinating plant growth. *Current Opinion in Plant Biology*, 9, 12-20.**
- Ishida, T., Aida, M., Takada, S. & Tasaka, M. (2000) Involvement of CUP-SHAPED COTYLEDON genes in gynoecium and ovule development in *Arabidopsis thaliana*. *Plant and Cell Physiology*, 41, 60-67.**
- Ishikawa, M., Ohmori, Y., Tanaka, W., Hirabayashi, C., Murai, K., Ogihara, Y., Yamaguchi, T. & Hirano, H. Y. (2009) The spatial expression patterns of DROOPING LEAF orthologs suggest a conserved function in grasses. *Genes & Genetic Systems*, 84, 137-146.**
- Izhaki, A. & Bowman, J. L. (2007) KANADI and class III HD-zip gene families regulate embryo patterning and modulate auxin flow during embryogenesis in *Arabidopsis*. *Plant Cell*, 19, 495-508.**
- Jack, T., Fox, G. L. & Meyerowitz, E. M. (1994) *Arabidopsis* Homeotic Gene *Apetala3* Ectopic Expression - Transcriptional and Posttranscriptional Regulation Determine Floral Organ Identity. *Cell*, 76, 703-716.**
- Jack, T., Sieburth, L. & Meyerowitz, E. (1997) Targeted misexpression of *AGAMOUS* in whorl 2 of *Arabidopsis* flowers. *Plant Journal*, 11, 825-839.**
- Jacobsen, S. E. & Meyerowitz, E. M. (1997) Hypermethylated *SUPERMAN* epigenetic alleles in *Arabidopsis*. *Science*, 277, 1100-1103.**



- Jacobsen, S. E., Running, M. P. & Meyerowitz, E. M. (1999) Disruption of an RNA helicase/RNase III gene in Arabidopsis causes unregulated cell division in floral meristems. *Development*, 126, 5231-5243.
- Jasinski, S., Piazza, P., Craft, J., Hay, A., Woolley, L., Rieu, I., Phillips, A., Hedden, P. & Tsiantis, M. (2005) KNOX action in Arabidopsis is mediated by coordinate regulation of cytokinin and gibberellin activities. *Current Biology*, 15, 1560-1565.
- Judd, W. S. & Olmstead, R. G. (2004) A survey of tricolpate (eudicot) phylogenetic relationships. *American Journal of Botany*, 91, 1627-1644.
- Jun, J. H., Ha, C. M. & Fletcher, J. C. (2010) BLADE-ON-PETIOLE1 Coordinates Organ Determinacy and Axial Polarity in Arabidopsis by Directly Activating ASYMMETRIC LEAVES2. *Plant Cell*, 22, 62-76.
- Kang, H. G., Jeon, J. S., Lee, S. & An, G. H. (1998) Identification of class B and class C floral organ identity genes from rice plants. *Plant Molecular Biology*, 38, 1021-1029.
- Kapoor, M., Tsuda, S., Tanaka, Y., Mayama, T., Okuyama, Y., Tsuchimoto, S. & Takatsuji, H. (2002) Role of petunia pMADS3 in determination of floral organ and meristem identity, as revealed by its loss of function. *Plant Journal*, 32, 115-127.
- Kayes, J. M. & Clark, S. E. (1998) CLAVATA2, a regulator of meristem and organ development in Arabidopsis. *Development*, 125, 3843-3851.
- Kawamura, E., Horiguchi, G. & Tsukaya, H. (2010) Mechanisms of leaf tooth formation in Arabidopsis. *Plant Journal*, 62, 429-441.
- Kellogg, E. A. (2001) Evolutionary history of the grasses. *Plant Physiology*, 125, 1198-1205.

- Kepinski, S. (2006) Integrating hormone signaling and patterning mechanisms in plant development. *Current Opinion in Plant Biology*, 9, 28-34.
- Kerstetter, R. A., Bollman, K., Taylor, R. A., Bomblied, K. & Poethig, R. S. (2001) KANADI regulates organ polarity in Arabidopsis. *Nature*, 411, 706-709.
- Kidner, C. A. & Martienssen, R. A. (2004) Spatially restricted microRNA directs leaf polarity through ARGONAUTE1. *Nature*, 428, 81-84.
- Kim, S., Koh, J., Ma, H., Hu, Y., Endress, P. K., Hauser, B. A., Buzgo, M., Soltis, P. S. & Soltis, D. E. (2005) Sequence and expression studies of A-, B-, and E-class MADS-box homologues in Eupomatia (Eupomatiaceae): Support for the bracteate origin of the calyptra. *International Journal of Plant Sciences*, 166, 185-198.
- Kramer, E. M., Jaramillo, M. A. & Di Stilio, V. S. (2004) Patterns of gene duplication and functional evolution during the diversification of the AGAMOUS subfamily of MADS box genes in angiosperms. *Genetics*, 166, 1011-1023.
- Krizek, B. A. & Fletcher, J. C. (2005) Molecular mechanisms of flower development: An armchair guide. *Nature Reviews Genetics*, 6, 688-698.
- Krizek, B. A. & Meyerowitz, E. M. (1996) The Arabidopsis homeotic genes APETALA3 and PISTILLATA are sufficient to provide the B class organ identity function. *Development*, 122, 11-22.
- Krizek, B. A., Prost, V. & Macias, A. (2000) AINTEGUMENTA promotes petal identity and acts as a negative regulator of AGAMOUS. *Plant Cell*, 12, 1357-1366.
- Kurakawa, T., Ueda, N., Maekawa, M., Kobayashi, K., Kojima, M., Nagato, Y., Sakakibara, H. & Kozuka, J. (2007) Direct control of shoot meristem activity by a cytokinin-activating enzyme. *Nature*, 445, 652-655.

- Kuusk, S., Sohlberg, J. J., Eklund, D. M. & Sundberg, E. (2006) Functionally redundant SHI family genes regulate *Arabidopsis* gynoecium development in a dose-dependent manner. *Plant Journal*, 47, 99-111.
- Kuusk, S., Sohlberg, J. J., Long, J. A., Fridborg, I. & Sundberg, E. (2002) STY1 and STY2 promote the formation of apical tissues during *Arabidopsis* gynoecium development. *Development*, 129, 4707-4717.
- Kuzoff, R. K. & Gasser, C. S. (2000) Recent progress in reconstructing angiosperm phylogeny. *Trends in Plant Science*, 5, 330-336.
- Laufs, P., Peaucelle, A., Morin, H. & Traas, J. (2004) MicroRNA regulation of the CUC genes is required for boundary size control in *Arabidopsis* meristems. *Development*, 131, 4311-4322.
- Laux, T., Mayer, K. F. X., Berger, J. & Jurgens, G. (1996) The WUSCHEL gene is required for shoot and floral meristem integrity in *Arabidopsis*. *Development*, 122, 87-96.
- Lee, B. H., Ko, J. H., Lee, S., Lee, Y., Pak, J. H. & Kim, J. H. (2009) The *Arabidopsis* GRF-INTERACTING FACTOR Gene Family Performs an Overlapping Function in Determining Organ Size as Well as Multiple Developmental Properties. *Plant Physiology*, 151, 655-668.
- Lee, I., Wolfe, D. S., Nilsson, O. & Weigel, D. (1997) A LEAFY co-regulator encoded by UNUSUAL FLORAL ORGANS. *Current Biology*, 7, 95-104.
- Lee, J. Y., Baum, S. F., Oh, S. H., Jiang, C. Z., Chen, J. C. & Bowman, J. L. (2005) Recruitment of CRABS CLAW to promote nectary development within the eudicot clade. *Development*, 132, 5021-5032.

- Lenhard, M., Bohnert, A., Jurgens, G. & Laux, T. (2001) Termination of stem cell maintenance in Arabidopsis floral meristems by interactions between WUSCHEL and AGAMOUS. *Cell*, 105, 805-814.
- Li, H. F., Liang, W. Q., Jia, R. D., Yin, C. S., Zong, J., Kong, H. Z. & Zhang, D. B. (2010) The AGL6-like gene OsMADS6 regulates floral organ and meristem identities in rice. *Cell Research*, 20, 299-313.
- Li, C. X., Potuschak, T., Colon-Carmona, A., Gutierrez, R. A. & Doerner, P. (2005) Arabidopsis TCP20 links regulation of growth and cell division control pathways. *Proceedings of the National Academy of Sciences of the United States of America*, 102, 12978-12983.
- Liljegren, S. J., Ditta, G. S., Eshed, H. Y., Savidge, B., Bowman, J. L. & Yanofsky, M. F. (2000) SHATTERPROOF MADS-box genes control seed dispersal in Arabidopsis. *Nature*, 404, 766-770.
- Liljegren, S. J., Roeder, A. H. K., Kempin, S. A., Gremski, K., Ostergaard, L., Guimil, S., Reyes, D. K. & Yanofsky, M. F. (2004) Control of fruit patterning in Arabidopsis by INDEHISCENT. *Cell*, 116, 843-853.
- Lin, W. C., Shuai, B. & Springer, P. S. (2003) The Arabidopsis LATERAL ORGAN BOUNDARIES-domain gene ASYMMETRIC LEAVES2 functions in the repression of KNOX gene expression and in adaxial-abaxial patterning. *Plant Cell*, 15, 2241-2252.
- Litt, A. & Irish, V. F. (2003) Duplication and diversification in the APETALA1/FRUITFULL floral homeotic gene lineage: Implications for the evolution of floral development. *Genetics*, 165, 821-833.

- Liu, Z. C., Franks, R. G. & Klink, V. P. (2000) Regulation of gynoecium marginal tissue formation by LEUNIG and AINTEGUMENTA. *Plant Cell*, 12, 1879-1891.
- Liu, X. D., Huang, J., Wang, Y., Khanna, K., Xie, Z. X., Owen, H. A. & Zhao, D. Z. (2010) The role of floral organs in carpels, an Arabidopsis loss-of-function mutation in MicroRNA160a, in organogenesis and the mechanism regulating its expression. *Plant Journal*, 62, 416-428.
- Liu, Z. C. & Meyerowitz, E. M. (1995) Leunig Regulates Agamous Expression in Arabidopsis Flowers. *Development*, 121, 975-991.
- Lohmann, J. U., Hong, R. L., Hobe, M., Busch, M. A., Parcy, F., Simon, R. & Weigel, D. (2001) A molecular link between stem cell regulation and floral patterning in Arabidopsis. *Cell*, 105, 793-803.
- Lolle, S. J. & Pruitt, R. E. (1999) Epidermal cell interactions: a case for local talk. *Trends in Plant Science*, 4, 14-20.
- Maier, A. T., Stehling-Sun, S., Wollmann, H., Demar, M., Hong, R. L., Haubeiss, S., Weigel, D. & Lohmann, J. U. (2009) Dual roles of the bZIP transcription factor PERIANTHIA in the control of floral architecture and homeotic gene expression. *Development*, 136, 1613-1620.
- Mallory, A. C., Reinhart, B. J., Jones-Rhoades, M. W., Tang, G. L., Zamore, P. D., Barton, M. K. & Bartel, D. P. (2004) MicroRNA control of PHABULOSA in leaf development: importance of pairing to the microRNA 5' region. *Embo Journal*, 23, 3356-3364.
- Mattsson, J., Sung, Z. R. & Berleth, T. (1999) Responses of plant vascular systems to auxin transport inhibition. *Development*, 126, 2979-2991.

- Mayer, K. F. X., Schoof, H., Haecker, A., Lenhard, M., Jurgens, G. & Laux, T. (1998) Role of WUSCHEL in regulating stem cell fate in the Arabidopsis shoot meristem. *Cell*, 95, 805-815.
- McConnell, J. R. & Barton, M. K. (1998) Leaf polarity and meristem formation in Arabidopsis. *Development*, 125, 2935-2942.
- McConnell, J. R., Emery, J., Eshed, Y., Bao, N., Bowman, J. & Barton, M. K. (2001) Role of PHABULOSA and PHAVOLUTA in determining radial patterning in shoots. *Nature*, 411, 709-713.
- Melzer, R., Wang, Y. Q. & Theissen, G. (2010) The naked and the dead: The ABCs of gymnosperm reproduction and the origin of the angiosperm flower. *Seminars in Cell & Developmental Biology*, 21, 118-128.
- Mizukami, Y. & Fischer, R. L. (2000) Plant organ size control: AINTEGUMENTA regulates growth and cell numbers during organogenesis. *Proceedings of the National Academy of Sciences of the United States of America*, 97, 942-947.
- Mizukami, Y. & Ma, H. (1995) Separation of Ag Function in Floral Meristem Determinacy from That in Reproductive Organ Identity by Expressing Antisense Ag Rna. *Plant Molecular Biology*, 28, 767-784.
- Mouradov, A., Glassick, T., Hamdorf, B., Murphy, L., Fowler, B., Maria, S. & Teasdale, R. D. (1998) NEEDLY, a Pinus radiata ortholog of FLORICAULA/LEAFY genes, expressed in both reproductive and vegetative meristems. *Proceedings of the National Academy of Sciences of the United States of America*, 95, 6537-6542.
- Nagasawa, N., Miyoshi, M., Sano, Y., Satoh, H., Hirano, H., Sakai, H. & Nagato, Y. (2003) SUPERWOMAN1 and DROOPING LEAF genes control floral organ identity in rice. *Development*, 130, 705-718.

- Nakagawa, H., Ferrario, S., Angenent, G. C., Kobayashi, A. & Takatsuji, H. (2004) The petunia ortholog of Arabidopsis SUPERMAN plays a distinct role in floral organ morphogenesis. *Plant Cell*, 16, 920-932.
- Nandi, A. K., Kushalappa, K., Prasad, K. & Vijayraghavan, U. (2000) A conserved function for Arabidopsis SUPERMAN in regulating floral-whorl cell proliferation in rice, a monocotyledonous plant. *Current Biology*, 10, 215-218.
- Nath, U., Crawford, B. C. W., Carpenter, R. & Coen, E. (2003) Genetic control of surface curvature. *Science*, 299, 1404-1407.
- Nemhauser, J. L., Feldman, L. J. & Zambryski, P. C. (2000) Auxin and ETTIN in Arabidopsis gynoecium morphogenesis. *Development*, 127, 3877-3888.
- Nesi, N., Debeaujon, I., Jond, C., Stewart, A. J., Jenkins, G. I., Caboche, M. & Lepiniec, L. (2002) The TRANSPARENT TESTA16 locus encodes the ARABIDOPSIS BSISTER MADS domain protein and is required for proper development and pigmentation of the seed coat. *Plant Cell*, 14, 2463-2479.
- Nikovics, K., Blein, T., Peaucelle, A., Ishida, T., Morin, H., Aida, M. & Laufs, P. (2006) The balance between the MIR164A and CUC2 genes controls leaf margin serration in Arabidopsis. *Plant Cell*, 18, 2929-2945.
- Nogueira, F. T. S., Madi, S., Chitwood, D. H., Juarez, M. T. & Timmermans, M. C. P. (2007) Two small regulatory RNAs establish opposing fates of a developmental axis. *Genes & Development*, 21, 750-755.
- Nole-Wilson, S. & Krizek, B. A. (2006) AINTEGUMENTA contributes to organ polarity and regulates growth of lateral organs in combination with YABBY genes. *Plant Physiology*, 141, 977-987.

- Norberg, M., Holmlund, M. & Nilsson, O. (2005) The **BLADE ONPETIOLE** genes act redundantly to control the growth and development of lateral organs. *Development*, 132, 2203-2213.
- Ohmori, S., Kimizu, M., Sugita, M., Miyao, A., Hirochika, H., Uchida, E., Nagato, Y. & Yoshida, H. (2009) **MOSAIC FLORAL ORGANS1**, an **AGL6-Like MADS Box Gene**, Regulates Floral Organ Identity and Meristem Fate in Rice. *Plant Cell*, 21, 3008-3025.
- Okada, K., Ueda, J., Komaki, M. K., Bell, C. J. & Shimura, Y. (1991) **REQUIREMENT OF THE AUXIN POLAR TRANSPORT-SYSTEM IN EARLY STAGES OF ARABIDOPSIS FLORAL BUD FORMATION**. *Plant Cell*, 3, 677-684.
- Orashakova, S., Lange, M., Lange, S., Wege, S. & Becker, A. (2009) The **CRABS CLAW** ortholog from California poppy (*Eschscholzia californica*, Papaveraceae), **EcCRC**, is involved in floral meristem termination, gynoecium differentiation and ovule initiation. *Plant Journal*, 58, 682-693.
- Ori, N., Cohen, A. R., Etzioni, A., Brand, A., Yanai, O., Shleizer, S., Menda, N., Amsellem, Z., Efroni, I., Pekker, I., Alvarez, J. P., Blum, E., Zamir, D. & Eshed, Y. (2007) Regulation of **LANCEOLATE** by **miR319** is required for compound-leaf development in tomato. *Nature Genetics*, 39, 787-791.
- Ostergaard, L. (2009) Don't 'leaf' now. The making of a fruit. *Current Opinion in Plant Biology*, 12, 36-41.
- Ostergaard, L. (2010) *Fruit Development and seed Dispersal*, Oxford, Wiley-Blackwell.
- Ostergaard, L., Kempin, S. A., Bies, D., Klee, H. J. & Yanofsky, M. F. (2006) Pod shatter-resistant Brassica fruit produced by ectopic expression of the **FRUITFULL** gene. *Plant Biotechnology Journal*, 4, 45-51.



- Palatnik, J. F., Allen, E., Wu, X. L., Schommer, C., Schwab, R., Carrington, J. C. & Weigel, D. (2003) Control of leaf morphogenesis by microRNAs. *Nature*, 425, 257-263.
- Palatnik, J. F., Wollmann, H., Schommer, C., Schwab, R., Boisbouvier, J., Rodriguez, R., Warthmann, N., Allen, E., Dezulian, T., Huson, D., Carrington, J. C. & Weigel, D. (2007) Sequence and expression differences underlie functional specialization of Arabidopsis MicroRNAs miR159 and miR319. *Developmental Cell*, 13, 115-125.
- Park, W., Li, J. J., Song, R. T., Messing, J. & Chen, X. M. (2002) CARPEL FACTORY, a Dicer homolog, and HEN1, a novel protein, act in microRNA metabolism in Arabidopsis thaliana. *Current Biology*, 12, 1484-1495.
- Payne, T., Johnson, S. D. & Koltunow, A. M. (2004) KNUCKLES (KNU) encodes a C2H2 zinc-finger protein that regulates development of basal pattern elements of the Arabidopsis gynoecium. *Development*, 131, 3737-3749.
- Pekker, I., Alvarez, J. P. & Eshed, Y. (2005) Auxin response factors mediate Arabidopsis organ asymmetry via modulation of KANADI activity. *Plant Cell*, 17, 2899-2910.
- Pelaz, S., Ditta, G. S., Baumann, E., Wisman, E. & Yanofsky, M. F. (2000) B and C floral organ identity functions require SEPALLATA MADS-box genes. *Nature*, 405, 200-203.
- Pelaz, S., Tapia-Lopez, R., Alvarez-Buylla, E. R. & Yanofsky, M. F. (2001) Conversion of leaves into petals in Arabidopsis. *Current Biology*, 11, 182-184.
- Pfluger, J. & Zambryski, P. (2004) The role of SEUSS in auxin response and floral organ patterning. *Development*, 131, 4697-4707.

- Phelps-Durr, T. L., Thomas, J., Vahab, P. & Timmermans, M. C. P. (2005) Maize rough sheath2 and its Arabidopsis orthologue ASYMMETRIC LEAVES1 interact with HIRA, a predicted histone chaperone, to maintain knox gene silencing and determinacy during organogenesis. *Plant Cell*, 17, 2886-2898.
- Pinyopich, A., Ditta, G. & Yanofsky, M. F. (2001) Roles of SEEDSTICK MADS box gene during ovule and seed development. *12th International Conference on Arabidopsis Research*. Madison, Wisconsin, University of Wisconsin Press.
- Pinyopich, A., Ditta, G. S., Savidge, B., Liljegren, S. J., Baumann, E., Wisman, E. & Yanofsky, M. F. (2003) Assessing the redundancy of MADS-box genes during carpel and ovule development. *Nature*, 424, 85-88.
- Prakash, N. & Alexander, J. H. (1984) Self-incompatibility in *Austrobaileya scandens*. IN WILLIAMS, E. G. & KNOX, R. B. (Eds.) *Pollination '84*. Melbourne, University of Melbourne.
- Prasad, K., Zhang, X. W., Tobon, E. & Ambrose, B. A. (2010) The Arabidopsis B-sister MADS-box protein, GORDITA, represses fruit growth and contributes to integument development. *Plant Journal*, 62, 203-214.
- Prigge, M. J., Otsuga, D., Alonso, J. M., Ecker, J. R., Drews, G. N. & Clark, S. E. (2005) Class III homeodomain-leucine zipper gene family members have overlapping, antagonistic, and distinct roles in Arabidopsis development. *Plant Cell*, 17, 61-76.
- Prunet, N., Morel, P., Negrutiu, I. & Trehin, C. (2009) Time to Stop: Flower Meristem Termination. *Plant Physiology*, 150, 1764-1772.
- Prunet, N., Morel, P., Thierry, A. M., Eshed, Y., Bowman, J. L., Negrutiu, I. & Trehin, C. (2008) REBELOTE, SQUINT, and ULTRAPETALA1 function redundantly in the temporal regulation of floral meristem termination in Arabidopsis thaliana. *Plant Cell*, 20, 901-919.

- Przemeck, G. K. H., Mattsson, J., Hardtke, C. S., Sung, Z. R. & Berleth, T. (1996) Studies on the role of the Arabidopsis gene MONOPTEROS in vascular development and plant cell axialization. *Planta*, 200, 229-237.
- Pulido, A. & Laufs, P. (2010) Co-ordination of developmental processes by small RNAs during leaf development. *Journal of Experimental Botany*, 61, 1277-1291.
- Ragni, L., Belles-Boix, E., Gunl, M. & Pautot, V. (2008) Interaction of KNAT6 and KNAT2 with BREVIPEDICELLUS and PENNYWISE in Arabidopsis inflorescences. *Plant Cell*, 20, 888-900.
- Rajani, S. & Sundaresan, V. (2001) The Arabidopsis myc/bHLH gene ALCATRAZ enables cell separation in fruit dehiscence. *Current Biology*, 11, 1914-1922.
- Rast, M. I. & Simon, R. (2008) The meristem-to-organ boundary: more than an extremity of anything. *Current Opinion in Genetics & Development*, 18, 287-294.
- Reinhardt, D., Frenz, M., Mandel, T. & Kuhlemeier, C. (2005) Microsurgical and laser ablation analysis of leaf positioning and dorsoventral patterning in tomato. *Development*, 132, 15-26.
- Reinhardt, D., Mandel, T. & Kuhlemeier, C. (2000) Auxin regulates the initiation and radial position of plant lateral organs. *Plant Cell*, 12, 507-518.
- Reinhardt, D., Pesce, E. R., Stieger, P., Mandel, T., Baltensperger, K., Bennett, M., Traas, J., Friml, J. & Kuhlemeier, C. (2003) Regulation of phyllotaxis by polar auxin transport. *Nature*, 426, 255-260.
- Rijkema, A. S., Zethof, J., Gerats, T. & Vandenbussche, M. (2009) The petunia AGL6 gene has a SEPALLATA-like function in floral patterning. *Plant Journal*, 60, 1-9.
- Rodriguez, R. E., Mecchia, M. A., Debernardi, J. M., Schommer, C., Weigel, D. & Palatnik, J. F. (2010) Control of cell proliferation in Arabidopsis thaliana by microRNA miR396. *Development*, 137, 103-112.

- Roeder, A. H. K., Ferrandiz, C. & Yanofsky, M. F. (2003) The role of the REPLUMLESS homeodomain protein in Patterning the Arabidopsis fruit. *Current Biology*, 13, 1630-1635.
- Roeder, A. H. K. & Yanofsky, M. F. (2005) Fruit development in Arabidopsis. *The Arabidopsis Book*.
- Rohde, A., Grunau, C., De Beck, L., Van Montagu, M., Rosenthal, A. & Boerjan, W. (1999) carpel, a new Arabidopsis epi-mutant of the SUPERMAN gene: Phenotypic analysis and DNA methylation status. *Plant and Cell Physiology*, 40, 961-972.
- Rudall, P. J., Sokoloff, D. D., Remizowa, M. V., Conran, J. G., Davis, J. I., Macfarlane, T. D. & Stevenson, D. W. (2007) Morphology of Hydatellaceae, an anomalous aquatic family recently recognized as an early-divergent angiosperm lineage. *American Journal of Botany*, 94, 1073-1092.
- Rudall, P. J., Stuppy, W., Cunniff, J., Kellogg, E. A. & Briggs, B. G. (2005) Evolution of reproductive structures in grasses (Poaceae) inferred by sister-group comparison with their putative closest living relatives, Ecteiocoleaceae. *American Journal of Botany*, 92, 1432-1443.
- Sakai, H., Krizek, B. A., Jacobsen, S. E. & Meyerowitz, E. N. (2000) Regulation of SUP expression identifies multiple regulators involved in arabidopsis floral meristem development. *Plant Cell*, 12, 1607-1618.
- Sakai, H., Medrano, L. J. & Meyerowitz, E. M. (1995) Role of Superman in Maintaining Arabidopsis Floral Whorl Boundaries. *Nature*, 378, 199-203.
- Savard, L., Li, P., Strauss, S. H., Chase, M. W., Michaud, M. & Bousquet, J. (1994) Chloroplast and Nuclear Gene-Sequences Indicate Late Pennsylvanian Time for

- the Last Common Ancestor of Extant Seed Plants. *Proceedings of the National Academy of Sciences of the United States of America*, 91, 5163-5167.
- Savidge, B., Rounsley, S. D. & Yanofsky, M. F. (1995) Temporal Relationship between the Transcription of 2 Arabidopsis MADS Box Genes and the Floral Organ Identity Genes. *Plant Cell*, 7, 721-733.
- Sawa, S., Watanabe, K., Goto, K., Kanaya, E., Morita, E. H. & Okada, K. (1999) FILAMENTOUS FLOWER, a meristem and organ identity gene of Arabidopsis, encodes a protein with a zinc finger and HMG-related domains. *Genes & Development*, 13, 1079-1088.
- Schneider, E. L. (1978) Morphological-Studies of Nymphaeaceae .9. Seed of *Barclaya-Longifolia* Wall. *Botanical Gazette*, 139, 223-230.
- Schoof, H., Lenhard, M., Haecker, A., Mayer, K. F. X., Jurgens, G. & Laux, T. (2000) The stem cell population of Arabidopsis shoot meristems is maintained by a regulatory loop between the CLAVATA and WUSCHEL genes. *Cell*, 100, 635-644.
- Schultz, E. A., Pickett, F. B. & Haughn, G. W. (1991) The Flo10 Gene-Product Regulates the Expression Domain of Homeotic Genes Ap3 and Pl in Arabidopsis Flowers. *Plant Cell*, 3, 1221-1237.
- Scofield, S., Dewitte, W. & Murray, J. A. H. (2007) The KNOX gene SHOOT MERISTEMLESS is required for the development of reproductive meristematic tissues in Arabidopsis. *Plant Journal*, 50, 767-781.
- Scutt, C. P., Vinauger-Douard, M., Fourquin, C., Ailhas, J., Kuno, N., Uchida, K., Gaude, T., Furuya, M. & Dumas, C. (2003) The identification of candidate genes for a reverse genetic analysis of development and function in the Arabidopsis gynoecium. *Plant Physiology*, 132, 653-665.

- Sessions, A., Nemhauser, J. L., McColl, A., Roe, J. L., Feldmann, K. A. & Zambryski, P. C. (1997) ETTIN patterns the Arabidopsis floral meristem and reproductive organs. *Development*, 124, 4481-4491.
- Sessions, R. A. & Zambryski, P. C. (1995) Arabidopsis Gynoecium Structure in the Wild-Type and in Effin Mutants. *Development*, 121, 1519-1532.
- Shani, E., Yanai, O. & Ori, N. (2006) The role of hormones in shoot apical meristem function. *Current Opinion in Plant Biology*, 9, 484-489.
- Shuai, B., Reynaga-Pena, C. G. & Springer, P. S. (2002) The LATERAL ORGAN BOUNDARIES gene defines a novel, plant-specific gene family. *Plant Physiology*, 129, 747-761.
- Sieber, P., Wellmer, F., Gheyselinck, J., Riechmann, J. L. & Meyerowitz, E. M. (2007) Redundancy and specialization among plant microRNAs: role of the MIR164 family in developmental robustness. *Development*, 134, 1051-1060.
- Sieburth, L. E., Drews, G. N. & Meyerowitz, E. M. (1998) Non-autonomy of AGAMOUS function in flower development: use of a Cre/loxP method for mosaic analysis in Arabidopsis. *Development*, 125, 4303-4312.
- Sieburth, L. E. & Meyerowitz, E. M. (1997) Molecular dissection of the AGAMOUS control region shows that cis elements for spatial regulation are located intragenically. *Plant Cell*, 9, 355-365.
- Sieburth, L. E., Running, M. P. & Meyerowitz, E. M. (1995) Genetic Separation of 3rd and 4th Whorl Functions of Agamous. *Plant Cell*, 7, 1249-1258.
- Siegel, B. A. & Verbeke, J. A. (1989) Diffusible Factors Essential for Epidermal-Cell Redifferentiation in Catharanthus-Roseus. *Science*, 244, 580-582.

- Siegfried, K. R., Eshed, Y., Baum, S. F., Otsuga, D., Drews, G. N. & Bowman, J. L. (1999) Members of the YABBY gene family specify abaxial cell fate in *Arabidopsis*. *Development*, 126, 4117-4128.
- Skyllar, A., Hong, F. X., Chory, J., Weigel, D. & Wu, X. L. (2010) STIMPY mediates cytokinin signaling during shoot meristem establishment in *Arabidopsis* seedlings. *Development*, 137, 541-549.
- Smyth, D. R., Bowman, J. L. & Meyerowitz, E. M. (1990) Early Flower Development in *Arabidopsis*. *Plant Cell*, 2, 755-767.
- Sohlberg, J. J., Myrenas, M., Kuusk, S., Lagercrantz, U., Kowalczyk, M., Sandberg, G. & Sundberg, E. (2006) STY1 regulates auxin homeostasis and affects apical-basal patterning of the *Arabidopsis* gynoecium. *Plant Journal*, 47, 112-123.
- Soltis, D. E., Albert, V. A., Leebens-Mack, J., Palmer, J. D., Wing, R. A., Depamphilis, C. W., Ma, H., Carlson, J. E., Altman, N., Kim, S., Wall, P. K., Zuccolo, A. & Soltis, P. S. (2008) The Amborella genome: an evolutionary reference for plant biology. *Genome Biology*, 9.
- Sommer, H., Beltran, J. P., Huijser, P., Pape, H., Lonnig, W. E., Saedler, H. & Schwarzsommer, Z. (1990) Deficiens, a Homeotic Gene Involved in the Control of Flower Morphogenesis in *Antirrhinum-Majus* - the Protein Shows Homology to Transcription Factors. *Embo Journal*, 9, 605-613.
- Sorefan, K., Girin, T., Liljegren, S. J., Ljung, K., Robles, P., Galvan-Ampudia, C. S., Offringa, R., Friml, J., Yanofsky, M. F. & Ostergaard, L. (2009) A regulated auxin minimum is required for seed dispersal in *Arabidopsis*. *Nature*, 459, 583-588.
- Sridhar, V. V., Surendrarao, A., Gonzalez, D., Conlan, R. S. & Liu, Z. C. (2004) Transcriptional repression of target genes by LEUNIG and SEUSS, two

- interacting regulatory proteins for Arabidopsis flower development. *Proceedings of the National Academy of Sciences of the United States of America*, 101, 11494-11499.
- Staldal, V., Sohlberg, J. J., Eklund, D. M., Ljung, K. & Sundberg, E. (2008) Auxin can act independently of CRC, LUG, SEU, SPT and STY1 in style development but not apical-basal patterning of the Arabidopsis gynoecium. *New Phytologist*, 180, 798-808.
- Staldal, V. & Sundberg, E. (2009) The role of auxin in style development and apical-basal patterning of the Arabidopsis thaliana gynoecium. *Plant Signal Behav*, 4, 83-5.
- Stevens, P. F. (2001 onwards) Angiosperm Phylogeny Website.
- Sun, B., Xu, Y. F., Ng, K. H. & Ito, T. (2009) A timing mechanism for stem cell maintenance and differentiation in the Arabidopsis floral meristem. *Genes & Development*, 23, 1791-1804.
- Sun, G., Dilcher, D. L., Zheng, S. L. & Zhou, Z. K. (1998) In search of the first flower: A Jurassic angiosperm, *Archaeofructus*, from northeast China. *Science*, 282, 1692-1695.
- Sussex, I. M. (1954) Morphogenesis in *Solanum tuberosum* L: experimental investigation of leaf dorsiventrality and orientation in the juvenile shoot. . *Phytomorphology* 5, 286-300.
- Tang, G. L., Reinhart, B. J., Bartel, D. P. & Zamore, P. D. (2003) A biochemical framework for RNA silencing in plants. *Genes & Development*, 17, 49-63.
- Taylor, E. L. & Taylor, T. N. (2009) Seed Ferns from the Late Paleozoic and Mesozoic: Any Angiosperm Ancestors Lurking There? *American Journal of Botany*, 96, 237-251.



- Theissen, G. (2001) Genetics of identity. *Nature*, 414, 491-491.
- Theissen, G., Becker, A., Winter, K.-U., Munster, T., Kirchner, C. & Saedler, H. (2002) How the land plants learned their floral ABCs: the role of MADS-box genes in the evolutionary origin of flowers. IN CRONK, Q. B. C., BATEMAN, R. M. & HAWKINS, J. A. (Eds.) *Developmental Genetics and Plant Evolution*. London, Taylor & Francis.
- Theissen, G. & Melzer, R. (2007) Molecular mechanisms underlying origin and diversification of the angiosperm flower. *Annals of Botany*, 100, 603-619.
- Theissen, G. & Saedler, H. (2001) Plant biology - Floral quartets. *Nature*, 409, 469-471.
- Thien, L. B., White, D. A. & Yatsu, L. Y. (1983) The Reproductive-Biology of a Relict *Illicium-Floridanum* Ellis. *American Journal of Botany*, 70, 719-727.
- Thompson, B. E., Bartling, L., Whipple, C., Hall, D. H., Sakai, H., Schmidt, R. & Hake, S. (2009) bearded-ear Encodes a MADS Box Transcription Factor Critical for Maize Floral Development. *Plant Cell*, 21, 2578-2590.
- Trigueros, M., Navarrete-Gomez, M., Sato, S., Christensen, S. K., Pelaz, S., Weigel, D., Yanofsky, M. F. & Ferrandiz, C. (2009) The NGATHA Genes Direct Style Development in the Arabidopsis Gynoecium. *Plant Cell*, 21, 1394-1409.
- Trobner, W., Ramirez, L., Motte, P., Hue, I., Huijser, P., Lonig, W. E., Saedler, H., Sommer, H. & Schwarzsommer, Z. (1992) Globosa - a Homeotic Gene Which Interacts with Deficiens in the Control of Antirrhinum Floral Organogenesis. *Embo Journal*, 11, 4693-4704.
- van der Graaff, E., Den Dulk-Ras, A., Hooykaas, P. J. J. & Keller, B. (2000) Activation tagging of the LEAFY PETIOLE gene affects leaf petiole development in *Arabidopsis thaliana*. *Development*, 127, 4971-4980.

- Vandenbussche, M., Theissen, G., Van de Peer, Y. & Gerats, T. (2003) Structural diversification and neo-functionalization during floral MADS-box gene evolution by C-terminal frameshift mutations. *Nucleic Acids Research*, 31, 4401-4409.
- Vazquez-Lobo, A., Carlsbecker, A., Vergara-Silva, F., Alvarez-Buylla, E. R., Pinero, D. & Engstrom, P. (2007) Characterization of the expression patterns of LEAFY/FLORICAULA and NEEDLY orthologs in female and male cones of the conifer genera *Picea*, *Podocarpus*, and *Taxus*: implications for current evo-devo hypotheses for gymnosperms. *Evolution & Development*, 9, 446-459.
- Vazquez, F., Vaucheret, H., Rajagopalan, R., Lepers, C., Gascioli, V., Mallory, A. C., Hilbert, J. L., Bartel, D. P. & Crete, P. (2004) Endogenous trans-acting siRNAs regulate the accumulation of *Arabidopsis* mRNAs. *Molecular Cell*, 16, 69-79.
- Verbeke, J. A. (1992) Fusion Events During Floral Morphogenesis. *Annual Review of Plant Physiology and Plant Molecular Biology*, 43, 583-598.
- Vialette-Guiraud, A. C. & Scutt, C. P. (2010) Carpel evolution. IN OSTERGAARD, L. (Ed.) *Fruit Development and Seed Dispersal*. Oxford, Wiley-Blackwell.
- Villanneva, J. M., Broadhvest, J., Hauser, B. A., Meister, R. J., Schneitz, K. & Gasser, C. S. (1999) INNER NO OUTER regulates abaxial-adaxial patterning in *Arabidopsis* ovules. *Genes & Development*, 13, 3160-3169.
- Vivian-Smith, A., Luo, M., Chaudhury, A. & Koltunow, A. (2001) Fruit development is actively restricted in the absence of fertilization in *Arabidopsis*. *Development*, 128, 2321-2331.
- von Goethe, J. W. (1790) *Versuch die Metamorphose der Pflanzen zu erklären*.
- Vroemen, C. W., Mordhorst, A. P., Albrecht, C., Kwaaitaal, M. & de Vries, S. C. (2003) The CUP-SHAPED COTYLEDON3 gene is required for boundary and shoot meristem formation in *Arabidopsis*. *Plant Cell*, 15, 1563-1577.

- Walker, D. B. (1978) Post-Genital Carpel Fusion in *Catharanthus-Roseus* (Apocynaceae) .4. Significance of Fusion. *American Journal of Botany*, 65, 119-121.
- Wang, J. W., Schwab, R., Czech, B., Mica, E. & Weigel, D. (2008) Dual effects of miR156-targeted SPL genes and CYP78A5/KLUH on plastochron length and organ size in *Arabidopsis thaliana*. *Plant Cell*, 20, 1231-1243.
- Wang, X., Duan, S. Y., Geng, B. Y., Cui, J. Z. & Yang, Y. (2007) *Schmeissneria*: A missing link to angiosperms? *Bmc Evolutionary Biology*, 7.
- Wege, S., Scholz, A., Gleissberg, S. & Becker, A. (2007) Highly efficient virus-induced gene silencing (VIGS) in california poppy (*Eschscholzia californica*): An evaluation of VIGS as a strategy to obtain functional data from non-model plants. *Annals of Botany*, 100, 641-649.
- White, D. W. R. (2006) PEAPOD regulates lamina size and curvature in *Arabidopsis*. *Proceedings of the National Academy of Sciences of the United States of America*, 103, 13238-13243.
- Williams, J. H. & Friedman, W. E. (2002) Identification of diploid endosperm in an early angiosperm lineage. *Nature*, 415, 522-526.
- Williams, J. H. & Friedman, W. E. (2004) The four-celled female gametophyte of *Illicium* (Illiciaceae; Austrobaileyales): Implications for understanding the origin and early evolution of monocots, eumagnoliids, and eudicots. *American Journal of Botany*, 91, 332-351.
- Williams, L., Carles, C. C., Osmont, K. S. & Fletcher, J. C. (2005) A database analysis method identifies an endogenous trans-acting short-interfering RNA that targets the *Arabidopsis* ARF2, ARF3, and ARF4 genes. *Proceedings of the National Academy of Sciences of the United States of America*, 102, 9703-9708.

- Willson, M. & Traveset, A. (2002) The ecology of seed dispersal. IN FENNER, M. (Ed.) *Seeds: The Ecology of Regeneration in Plant Communities*. Wallingford (UK), CAB International.
- Wu, G., Lin, W. C., Huang, T. B., Poethig, R. S., Springer, P. S. & Kerstetter, R. A. (2008) KANADI1 regulates adaxial-abaxial polarity in Arabidopsis by directly repressing the transcription of ASYMMETRIC LEAVES2. *Proceedings of the National Academy of Sciences of the United States of America*, 105, 16392-16397.
- Xu, L., Xu, Y., Dong, A. W., Sun, Y., Pi, L. M., Xu, Y. Q. & Huang, H. (2003) Novel as1 and as2 defects in leaf adaxial-abaxial polarity reveal the requirement for ASYMMETRIC LEAVES1 and 2 and ERECTA functions in specifying leaf adaxial identity. *Development*, 130, 4097-4107.
- Xu, L., Yang, L. & Huang, H. (2007) Transcriptional, post-transcriptional and post-translational regulations of gene expression during leaf polarity formation. *Cell Res*, 17, 512-9.
- Yalovsky, S., Rodriguez-Concepcion, M., Bracha, K., Toledo-Ortiz, G. & Gruissem, W. (2000) Prenylation of the floral transcription factor APETALA1 modulates its function. *Plant Cell*, 12, 1257-1266.
- Yamada, T., Ito, M. & Kato, M. (2003) Expression pattern of INNER NO OUTER homologue in Nymphaea (water lily family, Nymphaeaceae). *Development Genes and Evolution*, 213, 510-513.
- Yamaguchi, T., Lee, D. Y., Miyao, A., Hirochika, H., An, G. H. & Hirano, H. Y. (2006) Functional diversification of the two C-class MADS box genes OSMADS3 and OSMADS58 in *Oryza sativa*. *Plant Cell*, 18, 15-28.

- Yamaguchi, T., Nagasawa, N., Kawasaki, S., Matsuoka, M., Nagato, Y. & Hirano, H. Y. (2004) The YABBY gene DROOPING LEAF regulates carpel specification and midrib development in *Oryza sativa*. *Plant Cell*, 16, 500-509.**
- Yanai, O., Shani, E., Dolezal, K., Tarkowski, P., Sablowski, R., Sandberg, G., Samach, A. & Ori, N. (2005) Arabidopsis KNOXI proteins activate cytokinin biosynthesis. *Current Biology*, 15, 1566-1571.**
- Yanofsky, M. F., Ma, H., Bowman, J. L., Drews, G. N., Feldmann, K. A. & Meyerowitz, E. M. (1990) The Protein Encoded by the Arabidopsis Homeotic Gene *Agamous* Resembles Transcription Factors. *Nature*, 346, 35-39.**
- Yu, L. F., Patibanda, V. & Smith, H. M. S. (2009) A novel role of BELL1-like homeobox genes, PENNYWISE and POUND-FOOLISH, in floral patterning. *Planta*, 229, 693-707.**
- Yun, J. Y., Weigel, D. & Lee, I. (2002) Ectopic expression of SUPERMAN suppresses development of petals and stamens. *Plant and Cell Physiology*, 43, 52-57.**
- Zahn, L. M., Kong, H., Leebens-Mack, J. H., Kim, S., Soltis, P. S., Landherr, L. L., Soltis, D. E., de Pamphilis, C. W. & Ma, H. (2005) The Evolution of the SEPALLATA Subfamily of MADS-Box Genes: A Preangiosperm Origin With Multiple Duplications Throughout Angiosperm History. *Genetics*, 169, 2209-2223.**
- Zgurski, J. M., Sharma, R., Bolokoski, D. A. & Schultz, E. A. (2005) Asymmetric auxin response precedes asymmetric growth and differentiation of asymmetric leaf1 and asymmetric leaf2 Arabidopsis leaves. *Plant Cell*, 17, 77-91.**
- Zhao, L., Kim, Y. J., Dinh, T. T. & Chen, X. M. (2007) miR172 regulates stem cell fate and defines the inner boundary of APETALA3 and PISTILLATA expression domain in Arabidopsis floral meristems. *Plant Journal*, 51, 840-849.**

**Zhou, G. K., Kubo, M., Zhong, R. Q., Demura, T. & Ye, Z. H. (2007) Overexpression of miR165 affects apical meristem formation, organ polarity establishment and vascular development in Arabidopsis. *Plant and Cell Physiology*, 48, 391-404.**

Figure 1  
[Click here to download high resolution image](#)





Figure 2A  
[Click here to download high resolution image](#)





**B**




Figure 3  
[Click here to download high resolution image](#)




Figure 4  
[Click here to download high resolution image](#)




Figure 5  
[Click here to download high resolution image](#)




Figure 6  
[Click here to download high resolution image](#)




Figure 7  
[Click here to download high resolution image](#)




Figure 8

[Click here to download high resolution image](#)




Figure 9  
[Click here to download high resolution image](#)






Figure 10

[Click here to download high resolution image](#)




Figure 11

[Click here to download high resolution image](#)

