



HAL
open science

The XcpV/GspI pseudopilin has a central role in the assembly of a quaternary complex within the T2SS pseudopilus

Badreddine Douzi, Sébastien Alphonse, Herve Darbon, Alain Filloux, Christian Cambillau, Cedric Bernard, Mariella Tegoni, Romé Voulhoux

► To cite this version:

Badreddine Douzi, Sébastien Alphonse, Herve Darbon, Alain Filloux, Christian Cambillau, et al.. The XcpV/GspI pseudopilin has a central role in the assembly of a quaternary complex within the T2SS pseudopilus. Colloque de l'école doctorale Aix-Marseille, Jun 2009, Marseille, France. 1p, 2009. hal-02824819

HAL Id: hal-02824819

<https://hal.inrae.fr/hal-02824819>

Submitted on 6 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Copyright


The XcpV/GspI Pseudopilin has a central role in the assembly of a quaternary complex at the tip of the T2SS Pseudopilus

Badreddine Douzi¹, Eric Durand², Sébastien Alphonse¹, Hervé Darbon¹, Alain Filloux³, Christian Cambillau¹, Cédric Bernard¹, Mariella Tegoni¹ and Romé Voulhoux²


1. AFMB, UMR6098, CNRS, University of Aix-Marseille I & II case 932, 163 avenue de Luminy, 13288 Marseille cedex 09, France
 2. LISM UPR9027, CNRS, University of Aix-Marseille II, 31 chemin Joseph Aiguier, 13402 Marseille cedex 20, France
 3. Center of Molecular Microbiology and Infection (CMMI), Division of Cell and Molecular Biology, Imperial College London, London SW7 2AZ, United Kingdom
 §. Authors to whom correspondence should be addressed: Mariella.Tegoni@afmb.univ-mrs.fr/voulhoux@ibsm.cnrs-mrs.fr

Gram-negative bacteria use the sophisticated type II secretion system (T2SS) to transport a large number of exoproteins from the periplasmic space into the extracellular environment. However, if individual T2SS components are well characterized, very little is known about their multimeric organization during the secretion process. Five proteins of the T2SS, the pseudopilins GspG-H-I-J-K, are thought to assemble into a pseudopilus involved in the extrusion of the substrate through the outer membrane pore. Recent structural data have revealed that three of them, GspI-J-K, are organized in a trimeric complex proposed to be located at the tip of the GspG constituted pseudopilus. In the present work we combined two different biochemical techniques to investigate protein-protein interactions between the five *Pseudomonas aeruginosa* Xcp pseudopilins. The soluble domains of XcpT-U-V-W-X (respectively homologous to GspG-H-I-J-K) were purified and tested by surface plasmon resonance (SPR) and affinity purification for all possible interactions combinations. We found that XcpX_K interacts with XcpV_I which itself interacts with XcpW_J confirming the crystallized trimeric complex. Interestingly, our systematic approach also revealed a new interaction between XcpU_H and one member of the trimeric complex, XcpW_J, thus suggesting the existence of a quaternary complex involving XcpU_H-V_I-W_J-X_K at the tip of the pseudopilus. This quaternary complex was then successfully identified by affinity chromatography. Moreover, by testing various combinations of purified pseudopilins by SPR and affinity chromatography we were able to dissect the different successive steps involved in the development of the quaternary complex. We then propose a model for the quaternary pseudopilin complex formation centered around XcpV_I which first recruits XcpX_K on one side and XcpW_J on another side on which XcpU_H then binds. This central role of XcpV_I in pseudopilus formation fully agrees with our previous physiological data on its proposed involvement in pseudopilus initiation.


Pseudomonas aeruginosa



T2SS




The T2SS Piston Model




1 The secretion is performed by a Pseudopilus assembly

Expression of Xcp pseudopilin soluble domains Use of BIAcore 1000 for interaction study




Pseudopilin interaction network using Surface Plasmon Resonance (SPR)




(A) Interactions tested: each ligand with the 5 analytes. The K_{diss} values (μM) for the interaction detected. In bracket: the chip used for each ligand and the molecular weight of each analyte.
 (B) Schemes for all positive interactions, with the K_{diss} values indicated.

Epitope mapping of binary mixes using Surface Plasmon Resonance (SPR)



(A) W_J and binary mixes (W_J-U_H and W_J-X_K, 5 μM) were passed on V_I.
 (B, C) Schemes of the ternary complexes proposed, with K_{diss} of the interaction and K_{on} , k_{off} values when calculable.

Co-migration of pseudopilin soluble domains, batch-co-purified on affinity column




Each of the his6-tagged protein was mixed with different untagged protein partners. After affinity co-purification of proteins bound to the Ni²⁺-NTA-magnetic-beads, the different fractions were analyzed on a 15% SDS-PAGE. L1, containing the his6-tagged protein; L2, containing the untagged protein partners; L, containing tagged and untagged proteins. FT, flow-through; W, final wash; and E, elution.

Team How does the pseudopilus form?




Affinity determination of W_J-X_K and W_J-X_K-U_H mixes on V_I immobilized



(A) Binding pattern of W_J-X_K mixture at concentrations between 10 and 0.62 μM .
 (C) Binding pattern of W_J-X_K-U_H mixture at concentrations between 2.5 and 0.31 μM .
 (B, D) Schemes of the corresponding ternary complexes with K_{diss} of the interaction, and K_{on} , k_{off} values when calculable.

Proposed sequential assembly of the pseudopilins quaternary complex



- 1 Is time zero of the pseudopilus formation.
- 2 The presence of the substrate to be translocated activates (white stars) the formation of the complex.
- 3 The pseudopilus formed by the assembled quaternary complex and the polymerization of XcpT_G, pushes the substrate through the secretion pore.
- 4 After secretion, the interaction between XcpX_K and secretin activates (yellow star), the pseudopilus disassembly, and the system goes back to the resting state of time zero.