



**HAL**  
open science

## Responses of photosynthesis to high temperatures in trees: acclimation of the thermal break point of photochemistry to heat and drought

Daniel D. Epron, Hana Ghouil, Chiraz Daas, Fabienne Froux, Charlie Warren, Erwin Dreyer

### ► To cite this version:

Daniel D. Epron, Hana Ghouil, Chiraz Daas, Fabienne Froux, Charlie Warren, et al.. Responses of photosynthesis to high temperatures in trees: acclimation of the thermal break point of photochemistry to heat and drought. Impacts of the Drought and Heat in 2003 on Forests, Nov 2004, Freiburg, Germany. hal-02826899

**HAL Id: hal-02826899**

**<https://hal.inrae.fr/hal-02826899v1>**

Submitted on 7 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

#### **2.4.4 Responses of photosynthesis to high temperatures in trees: acclimation of the thermal break point of photochemistry to heat and drought**

Daniel Epron<sup>1</sup>, Hana Ghouil<sup>2</sup>, Chiraz Daas<sup>2</sup>, Fabienne Froux<sup>1</sup>, Charlie Warren<sup>3</sup>,  
Erwin Dreyer<sup>1</sup>

<sup>1</sup> UMR INRA-UHP « Ecologie et Ecophysiologie Forestières », F 54280  
CHAMPENOUX, France; dreyer@nancy.inra.fr

<sup>2</sup> Laboratoire d'Ecophysiologie et de Reboisement, Institut National de Recherches  
Forestières, Tunis, Tunisie

<sup>3</sup> Forest Science Center, University of Melbourne and Department of Environment of  
Victoria, Creswick, Vic. 3363, Australia

The physiological processes contributing to photosynthesis (i.e., photochemistry, carbon reduction cycle, CO<sub>2</sub> diffusion through stomata and to the chloroplasts) display different responses to high temperatures that still need to be better documented, despite the knowledge acquired during the past decades. The thermal optima of photochemistry and rubisco activity have been found to be around 35 and 38 C; but neither the inter and intra-specific diversity of this parameter, nor the ability of these optima to acclimate to rapidly changing environmental conditions have yet been clearly established. The thermal breakpoint of chlorophyll fluorescence is frequently used as one of the indexes for the thermostability of PSII and thylakoid membranes. This breakpoint is usually close to 45 C in many species. Recent experiments have shown that it is highly plastic, increases by up to 10 C being induced either by drought or by gradually increasing temperatures. In this presentation, we will summarise the information available in this field, as well as show the latest evidence documenting the plasticity of this parameter and the consequences it might have for the photosynthetic functions as a whole, basing on experiments with oak trees.