



**HAL**  
open science

## Specialist seed insects can affect the conservation of endangered conifer species

Alain Roques, Marie-Anne Auger-Rozenberg

► **To cite this version:**

Alain Roques, Marie-Anne Auger-Rozenberg. Specialist seed insects can affect the conservation of endangered conifer species. 22. International Congress of Entomology, Aug 2004, Brisbane, Australia. 1 p. hal-02827405

**HAL Id: hal-02827405**

**<https://hal.inrae.fr/hal-02827405>**

Submitted on 7 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## **Specialist seed insects can affect the conservation of endangered conifer species**

A Roques and MA Auger-Rozenberg

INRA, Unité de Zoologie Forestière, BP 20619 Ardon 45166 Olivet cedex, France,

[Alain.Roques@orleans.inra.fr](mailto:Alain.Roques@orleans.inra.fr)

Most insects exploiting conifer seed cones are highly host-specific. Several ecological adaptations, especially prolonged diapause, allow these insects to cope partially with the annual variations usually observed in host abundance. Long-term, standardized surveys were carried out in Europe and China in order to assess subsequent insect impact on the potential of natural regeneration of conifers. In widespread species (e.g., larches, *Larix* spp.), insect impact dramatically decreased the potential seed yield but a sufficient natural regeneration occurred when unforeseeable climatic events (e.g. late frosts) perturbed the natural synchrony between insect populations and cone abundance. Contrasted results were obtained for endangered conifers. In incense-juniper (*Juniperus thurifera*), a relict species with a very scattered range, low numbers of initial seeds per cone in Morocco and France resulted in a quasi-absence of viable seeds following insect attack whereas higher seed numbers allowed an important natural regeneration in Spain. In China, intensive forest exploitation limited the range of *Keteleeria xerophylla* and *Pseudotsuga sinensis* to very small, isolated plots where up to 100% of the seeds were annually destroyed by insects. Insect control seems the only way to prevent species disappearance.