



Spatial modelling and ecology of *Echinococcus multilocularis*

Francis M. Danson, Patrick Giraudoux, P.S. Craig

► To cite this version:

Francis M. Danson, Patrick Giraudoux, P.S. Craig. Spatial modelling and ecology of *Echinococcus multilocularis*. 3. Congress of the Federation of Asian Parasitologists, Jul 2005, Asahikawa, China. 1 p. hal-02828519

HAL Id: hal-02828519

<https://hal.inrae.fr/hal-02828519>

Submitted on 7 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

SPATIAL MODELLING AND ECOLOGY OF *ECHINOCOCCUS MULTILOCULARIS* TRANSMISSION IN CENTRAL ASIA

Francis M. Danson¹, P. Giraudoux² and P. S. Craig¹

¹Centre for Environmental Systems Research and Biosciences Research Institute, School of Environment and Life Sciences, University of Salford, Manchester M5 4WT, UK

²ISTE (EA 3184 MRT – UC INRA; EA 2276) – OMS, Université de Franche-Comté, 25030, Besançon Cedex, France

Transmission of *Echinococcus multilocularis* to humans occurs by accidental ingestion of eggs either indirectly through contaminated vegetation, or directly through contact with an infected canid. Our work in China, and more recently in Kazakhstan, has suggested that the most likely transmission mechanism in these regions is *via* domestic dogs which are allowed to roam freely and hunt (infected) small mammals within areas close to villages or tented pastures. This assertion has led to the hypothesis that there is a landscape control on transmission risk since the proximity of suitable habitat for susceptible small mammals appears to be key. We have tested this hypothesis in a number of endemic areas in China, notably south Gansu and Ningxia Provinces and most recently in the Tibetan region of western Sichuan Province. The fundamental landscape control is its effect at a regional scale on small mammal species assemblages (susceptible species are not ubiquitous) and, at a local scale, the spatial distributions of small mammal populations. To date we have examined relationships between landscape composition and patterns of human infection, and landscape and small mammal distributions. In current work we are examining the relationships between landscape and dog infection rates. The key tool to characterize landscape is satellite remote sensing and these data are used as inputs to drive spatial models of transmission risk. This paper reviews the progress that has been made so far in spatial modeling of the ecology of *E. multilocularis*, outlines current research issues, and describes a framework for building a spatial-temporal model of transmission risk.