



**HAL**  
open science

## **Transmission ecology of Echinococcus in China: advances and hot questions**

Patrick Giraudoux, Francis Raoul, Jean-Pierre Quere, David Pleydell, Qian Wang, Jiamen Qiu, Wen Yang, Philip S. Craig

► **To cite this version:**

Patrick Giraudoux, Francis Raoul, Jean-Pierre Quere, David Pleydell, Qian Wang, et al.. Transmission ecology of Echinococcus in China: advances and hot questions. 3. Congress of the Federation of Asian Parasitologists, Jul 2005, Asahikawa, China. pp.53. hal-02828792

**HAL Id: hal-02828792**

**<https://hal.inrae.fr/hal-02828792v1>**

Submitted on 7 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## TRANSMISSION ECOLOGY OF *ECHINOCOCCUS* IN CHINA: ADVANCES AND HOT QUESTIONS

Patrick Giraudoux<sup>1</sup>, Francis Raoul<sup>1</sup>, Jean-Pierre Quéré<sup>2</sup>, David Pleydell<sup>1</sup>, Qian Wang<sup>1, 3</sup>, Jiamen Qiu<sup>3</sup>, Wen Yang<sup>3</sup> and Philip S. Craig<sup>4</sup>

<sup>1</sup> Department of Environmental Biology – EA3184 - usc INRA, Université de Franche Comté, Place Leclerc, 25030 Besançon cedex, France

<sup>2</sup> Campus International du Baillarguet, CBGP, CS 30016 Montferrier sur Lez, 34988 Saint Gély du Fesc cedex, France

<sup>3</sup> Sichuan Institute of Parasitic Diseases, 10, University Road, Chengdu 610041 Sichuan, P.R. China

<sup>4</sup> School of Environmental and Life Sciences, University of Salford, M54WT, Salford, UK

In continental China *Echinococcus multilocularis* (Em) transmission intensity is spatially heterogeneous and unstable. This instability makes transmission responsive to fox/dog dosing control campaigns and variations in host population densities. Parasite extinction may occur as a consequence of landscape/land use change (e.g. Zhang-Puma area of southern Gansu, Northern Liu Pan Shan, Ningxia). By contrast, intensive and more stable transmission occurs in some areas of the Tibetan plateau due to sustainable high densities of small mammal intermediate host populations, the existence of two species of fox definitive hosts, and the extremely high number of dogs in and around villages with easy access to small mammal colonies. Yak herd management and subsequent grazing patterns may be an important factor impacting both the composition of small mammal communities and the population dynamics of intermediate host-species. Using data collected from comprehensive surveys in China and Europe as evidence, this paper sets out four concepts of transmission addressed by the following questions:

- 1) Do short-term studies within localized (sub-county) areas provide sufficient information to understand the processes leading to stable transmission of Em?
- 2) If no, how many small mammal communities and landscape habitats may be significantly involved in transmission?
- 3) Thus, what is the spatial extent of areas within which meta-stability operates and which areas can subsequently be considered stable foci ?
- 4) To which geographical range can the parasite be spread by foxes and the dog trade, thus can the likely meta-stable foci of the Tibetan plateau be a source of infection to transitory favorable remote areas of the spurs of the plateau (e.g. Southern Gansu) or even of more remote ranges (e.g. Liu Pan Shan)?