



**HAL**  
open science

# **A Statistical Application in Biological Phenomena: The Use of Mixed Models in the Analysis of the Growth Curve of Beef Cattle and Feed Intake of Lactating Sows**

Jean-Luc Gourdine, David Renaudeau, Michel Naves, Alberto Menendez Buxadera

## ► To cite this version:

Jean-Luc Gourdine, David Renaudeau, Michel Naves, Alberto Menendez Buxadera. A Statistical Application in Biological Phenomena: The Use of Mixed Models in the Analysis of the Growth Curve of Beef Cattle and Feed Intake of Lactating Sows. 6e édition de l'université de juillet 2004, Jul 2004, Pointe-à-Pitre, Guadeloupe, France. 2004. hal-02829356

**HAL Id: hal-02829356**

**<https://hal.inrae.fr/hal-02829356v1>**

Submitted on 7 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.


# A Statistical Application in Biological Phenomena : The Use of Mixed Models in the Analysis of the Growth Curve of Beef Cattle and Feed Intake of Lactating Sows.



J.L. Gourdine, D. Renaudeau, M. Naves and A. Menendez-Buxadera.  
INRA, Unité de Recherches Zootechniques, Domaine Duclos, 97170 Petit-Bourg, Guadeloupe.



## A biostatistical process



## An introduction to Linear Mixed Model

Let  $y_i$  be the vector of  $n$  measurements on the same  $i^{\text{th}}$  of  $m$  individuals.

The linear mixed model is written as

$$Y = \text{fixed regressions } (X\beta) + \text{random part } (\epsilon = Zu + W(t) + e)$$

With  $\epsilon \sim N_{nm}(0, V(t,u))$  and  $V(t,u) = ZGZ' + R$  where  $R = \sigma^2H + \tau^2I$   
where :

- $X$  is the design matrix of fixed effects
- $\beta$  is the unknown vector of fixed effects
- $Z$  is the known design matrix of random effects
- $u$  is the vector of unknown random-effects parameters
- $W(t)$  is the unknown vector of serial correlation
- and  $e$  is measurement error.

Estimation :

- Likelihood-based methods provide estimation of  $G$ ,  $R$ ,  $\beta$  and  $u$
- Estimation of  $\beta$  and  $u$  are the empirical best linear unbiased estimator (EBLUE) and best linear unbiased predictor (EBLUP)

## Application : two zootechnical examples

### Effect of breed (Creole vs. Large White) on lactating sow feed intake

#### Knowledge evaluation :

Genotype difference :

Prolificity : + LW sow (12 vs. 9 piglets/farrowing); Adaptation to heat stress: + CR sows (assumed).

Factors influencing feed intake (FI) during lactation :

Animal : breed, litter size, etc ...; Environmental : temperature, humidity, etc...; Dietary : energy and protein contents, etc ...

#### Purpose of the study :

Is sow feed intake pattern a good criteria to study heat stress tolerance in lactating sows?

Scientific objective : a descriptive purpose

Quantify the effect of genotype on feed intake during lactation

#### Model and data

**Data** : Daily feed intake (FI) performance from farrowing to 21 d post-partum in 35 Creole (CR) and 44 Large White (LW) lactating sows (225 lactations).

**Model** : Let  $y_{ijkl}$  be the observed value of FI of the sow  $j(i)$  of genotype  $i$  of the contemporary group  $l$ , taken in day  $k$  of lactation. Feed intake was expressed per metabolic weight and per one weaned piglet to take into account different maintenance and production requirements between CR and LW sows.

The mixed model is written as :  $Y_{ijkl} = \text{Fixed part } (\sum_m [\beta_m + \delta_m] t_k^m) + \text{Random part } (g_{j(i)} + r_i(t_k) + e_{ijkl})$

$\beta$  is the fixed regression coefficients to fit mean FI pattern

$\delta$  is the fixed parameters to fit the effect of breed on the mean curve


$g$  is the random-effect associated with the animal (because a sow has several lactations);  $r$  is the random-effect associated with time  $t_k$  (because of correlation between measurements on the same sow at different times);  $e$  is the measurement error

#### Results and discussion

Genotype effect	Pvalue
$\delta_{0_1}$ deviation from the intercept	0.42
$\delta_{1_1}$ deviation from the linear parameter	< 0.001
$\delta_{2_1}$ deviation from the quadratic parameter	< 0.001
$\delta_{3_1}$ deviation from the cubic parameter	< 0.01

The intercept was the same in both breeds. Nevertheless, the differences were significant when the linear, quadratic and cubic coefficients were considered; so LW and CR sows have different feed intake pattern (Figure 1a). FI of LW sow was greater during the first two weeks post-partum, but it was lower at week 3 of lactation.

CR sows were able to increase their FI (and their heat production) as their requirements for milk production increased. At the opposite, it seems that LW sows needs more time for adaptation because their consumption decreased during the third week and increased after.



### Estimation of genetic variability of Creole cow growth curve

#### Knowledge evaluation :

Factors influencing body growth of Creole cows :

• genetic : pedigree, non genetic effect : Management, environment, etc ...

Selection criteria : Liveweight at 18 months of age (LW18).

#### Purpose of the study :

Is possible to make the selection at an earlier age than 18 months of age?

What is the level of relationships between selection criteria and liveweight at different ages?

Scientific objective : predictive purpose

Estimate genetic variability of growth parameters

#### Model and data

**Data** : Growth performance from birth to 10 years in 227 Creole cows (117 dams).

**Model** : Let  $y_{ij}$  be the  $j^{\text{th}}$  liveweight record for cow  $i$  taken at age  $t_{ij}$ .

The mixed model is written as :  $Y_{ijkl} = \text{Fixed part } (F_i + \sum_m (\beta_m t_{ij}^m)) + \text{Random part } (g(t_{ij}) + r(t_{ij}) + e_{ij})$

$\beta$  is the fixed regression coefficients to fit mean growth curve

$F_i$  is the fixed effect of mother calving number on growth on body weight at birth


$g$  is the random-effect associated with the genetic additive effect

$r$  is the random-effect associated with the permanent environment effect (non genetic variation due to the cow)

$e$  is the measurement error


#### Results and discussion

Figure 1b. Heritability of liveweight of Creole cow.



Genetic correlation between liveweight at 10, 12 and 16 months of age and the selection criteria at LW18 are high (>0.84). We can estimate heritability  $h^2$  (figure 1b) from the estimation of the additive genetic covariance function  $g$ .  $h^2$  is the part of variance due to additive genetic component related to the total phenotypic variance.  $h^2$  at 10, 12, 16 months of age (• in figure 1b) are to the same extent as heritability at 18 months of age (> 30%). That means, selection for growth performance at 10 months of age is approximately equivalent as selection at LW18. This is illustrated in Figure 2b which presented the best genetic animal (in red) of the studied population and the worst cow (in blue) for growth performance criteria. The deviation at LW10 and at LW18 is 180 and 200 kg, respectively, and genetic part both represent 40 % in the deviation.

Figure 2b. Best and worst genetic Creole cow for growth performance.



## Conclusion

The use of linear mixed models, with random regression, is an answer to the difficulty in the choice of many statistical models in zootechnic area:

- Mixed models take into account random and fixed parts to provide good estimations (Best linear unbiased)
- They tend to be generally used in zootechnical research because characters of economical interest are often longitudinal data, that means repeated measurements on the same animal.