

Pinus yunnanensis resistance to a bark beetle associated fungus is increased during a natural water stress

François Lieutier, Ye Hui, Annie Yart

► **To cite this version:**

François Lieutier, Ye Hui, Annie Yart. Pinus yunnanensis resistance to a bark beetle associated fungus is increased during a natural water stress. 15. International Plant Protection Congress, May 2004, Beijing, China. pp.Number 10.2. hal-02829719

HAL Id: hal-02829719

<https://hal.inrae.fr/hal-02829719>

Submitted on 7 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

15th International Plant Protection Congress
Beijing China, May11-16, 2004

ABSTRACT SUBMISSION FORM

Abstracts must be received before January 31, 2004

PINUS YUNNANENSIS RESISTANCE TO A BARK BEETLE ASSOCIATED FUNGUS IS INCREASED BY A NATURAL WATER STRESS.

François LIEUTIER¹, YE Hui² and Annie YART³

1 Laboratoire de Biologie des Ligneux et des Grandes Cultures, Université d'Orléans, rue de Chartres, B.P. 6759, F-45067 Orléans Cedex 2, France.

2 Yunnan University, Institute of Ecology and Geobotany, 650091 Kunming, Yunnan, P.R. China.

3 INRA, Unité de Zoologie Forestière, Ardon, F-45160 Olivet, France.

Email: francois.lieutier@univ-orleans.fr

Two neighboring sites of an area forested with Yunnan pine (*Pinus yunnanensis*), located in Yunnan province (China) and clearly differing by their water availability, were compared for tree resistance to isolated and mass artificial inoculations with *Leptographium yunnanense*, a fungus associated with the bark beetle *Tomicus* sp. During 18 months alternating dry and humid seasons in each site, trees' physiological parameters (soil water content at different depths, predawn needle water potential, needle water content) were followed in the two sites. Simultaneously, the performances of the fungus after isolated inoculation (growth in the phloem, ability to induce the development of a phloem induced reaction) and its virulence after mass inoculations were periodically measured to appreciate tree resistance. The phloem induced reactions were more extended during the wet than during the dry season, and fungus grew further into the phloem during the humid season and in the wet site than during the dry season and in the dry site. Fungus virulence after mass inoculations was higher during the humid period and in the wet site than during the dry period and in the dry site. These results demonstrate that tree resistance to fungus inoculation and consequently to bark beetle attacks, is increased by a mild water stress. This is in agreement with the growth-differentiation balance hypothesis and with the idea that the fungus is favored by a large availability of water. In conclusion, drought is not a general explanation for *Tomicus* damage in Yunnan.

Abstract Title PINUS YUNNANENSIS NATURAL RESISTANCE TO A BARK BEETLE ASSOCIATED FUNGUS IS INCREASED BY A MILD WATER STRESS.

Choice of Presentation: Oral Poster

The equipment I need: Data projector (PowerPoint) 35 mm slide Overhead

Symposium Number 10.2 (first Choice) _____ (Second Choice)

See the Symposium Numbers in the Scientific Program of this circular.

The Name of Presenting Author and the Address:

First Name Francois Middle Name _____ Last Name LIEUTIER

Institution University of Orleans

Mailing Address Laboratoire de Biologie des Ligneux et des Grandes Cultures,
rue de Chartres, B.P. 6759, 45067 Orleans Cedex 2 FRANCE.

Tel 33 / 2 38 41 72 30 Fax 33 / 2 38 41 72 02 E-mail francois.lieutier@univ-orleans.fr