



HAL
open science

Fate of ivermectin in soil and impact on micro-organisms

Christian Mougin, J.C. Fournier, M. Alvinerie, D. Berry, Dominique Kerboeuf, J.P. Lumaret, Pierre Galtier

► To cite this version:

Christian Mougin, J.C. Fournier, M. Alvinerie, D. Berry, Dominique Kerboeuf, et al.. Fate of ivermectin in soil and impact on micro-organisms. *Envirpharma European Conference on Human and Veterinary Pharmaceuticals in the Environment*, Apr 2003, Lyon, France. 1 p., 2003. hal-02830951

HAL Id: hal-02830951

<https://hal.inrae.fr/hal-02830951>

Submitted on 13 Mar 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

INTRODUCTION

-Avermectins are a class of macrocyclic lactone drugs with insecticidal and anthelmintic properties, which have been developed for the protection of animals, humans and crops
-The intraruminal-sustained release bolus of ivermectin is considered an important advance in the control of endoparasitic nematodes in ruminants because of the long-term protection of animals
-High levels of faecal excretion of the drug could induce toxic effects on key dung-colonizing families of insects, and on soil or aquatic ecosystems inhabiting (micro)-organisms

→ We present here our results concerning the fate of ivermectin in soils, as a basis for exposure assessment, and its impact on bacteria and fungi.

EXCRETION OF IVERMECTIN

Faecal concentrations of ivermectin are obtained after the administration of a sustained-release bolus to calves (initial amount: 1.72 g per animal)

Time after the administration of the bolus (days)	Time of presence on pasture (days)	Faecal concentrations (mg/kg)
4	15	0,688
	30	1,617
	60	0,456
	90	0,181
14	15	1,123
	30	1,473
42	15	0,839
	30	1,184
70	15	1,056

→ 1 mg/kg untransformed drug is released in faeces during a 4-month period

BIOAVAILABILITY OF IVERMECTIN

Amount of ivermectin measured in the soil solution was 24 ± 14 ng
→ 0.2 % of initial amount in soil → $6 \mu\text{g/L} \rightarrow 10^{-5}$ M


GLOBAL EFFECTS OF IVERMECTIN ON BACTERIA

No effect of ivermectin (10, 50 and 250 mg/kg) on the number of colony-forming organisms after 15 days on solid media
No effect of ivermectin (10, 50 and 250 mg/kg) on the microbial biomass after 15 days in the soil (carbon extracted after fumigation)

No effect of ivermectin (10 and 50 mg/kg) on the nitrification of NH_4^+ in the soil
No effect of ivermectin (10 mg/kg) on the mineralization of labeled carbofuran (insecticide)
No effect of ivermectin (10, 50 and 250 mg/kg) on the turn-over of microbial carbon (using labeled glucose) in the soil

No effect of ivermectin (10 mg/l) on the metabolic activities of soil isolates using Biolog plates


→ No effect of the drug was detected either on the size of the microbial biomass or functions, even at high concentrations



SHORT-TERM EFFECTS OF IVERMECTIN ON FUNGI

• Fungal strains, namely *Trametes versicolor* (●), *Fusarium solani* (○), *Mucor racemosus* (▼), and *Fusarium solani* (△) have been cultured in liquid media in the presence of ivermectin at various concentrations

• Fungal cultures, inoculated with spore suspensions, were allowed to grow in the presence of ivermectin (10^{-4} M). The spores produced by the new hyphae have been harvested and counted. (*T. versicolor*, unable to produce spores, was replaced by *P. chrysosporium*, another white-rot strain)



→ No negative impact of the drug has been evidenced on the fungal biomass production after 4, 8 and 12 days of growth

→ Ivermectin was also unable to inhibit the germination of fungal spores, even at high concentrations (10^{-3} M, data not shown)

→ The drug diversely affected fungal sporulation, according to the strains

INDUCTION OF ENZYMATIC ACTIVITIES BY IVERMECTIN

Fungal cultures have been grown in liquid media in the presence of 10^{-5} M ivermectin. Enzymatic activities were then measured using colorimetric assays

Laccases of *T. versicolor* after 3 days of treatment

Controls	
Water	0.16 ± 0.02 U/mL
Xylydine	5.69 ± 0.21
Ivermectin	0.32 ± 0.06
8,9-Z isomer	0.33 ± 0.06
10,11-Z isomer	0.49 ± 0.09


Lignin-peroxidases of *P. chrysosporium* after 4, 8 and 12 days

→ No effect

→ Ivermectin is not an inducer of fungal enzymatic activities

LONG-TERM EFFECTS OF IVERMECTIN ON FUNGI

Fungal cultures have been inoculated with spores, without ivermectin (●) or in the presence of the drug (◊) at 10^{-5} M. The culture were then allowed to grow and sporulate for 10 days. Spores were harvested and the experiments were still repeated 3-times. The spore multiplication factor reflected the ratio between the amounts of spores produced by a culture versus the amounts of spores used for inoculation.



→ Ivermectin diversely affected fungal sporulation, according to the strains
→ It was without effect on fungal growth (data not shown)

CONCLUSIONS

- 1 - Ivermectin can reach the soil in high amounts after the treatment of cattle with the intraruminal sustained release bolus
- 2 - In the soil, the drug undergoes minor transformation reactions, only due to photo-isomerization
- 3 - The bioavailability of ivermectin for soil micro-organisms is very reduced
- 4 - Ivermectin induces no negative effect on bacteria
- 5 - It only impacts fungal reproduction, with positive or negative effects according to the strains