

Invasive patterns of an exotic seed chalcid, Megastigmus schimitscheki Novitzky, in the cedar plantations of southern France

Marie-Anne Auger-Rozenberg, Lionel Roques, Solen Boivin, Alain Chalon, Jean-Pierre Fabre, Alain Roques

▶ To cite this version:

Marie-Anne Auger-Rozenberg, Lionel Roques, Solen Boivin, Alain Chalon, Jean-Pierre Fabre, et al.. Invasive patterns of an exotic seed chalcid, Megastigmus schimitscheki Novitzky, in the cedar plantations of southern France. 15. International Plant Protection Congress, Jul 2003, Beijing, China. 1 p. hal-02831115

HAL Id: hal-02831115 https://hal.inrae.fr/hal-02831115

Submitted on 7 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

15th International Plant Protection Congress Beijing, China, July 6-11, 2003

INVASIVE PATTERNS OF AN EXOTIC SEED CHALCID, MEGASTIGMUS SCHIMITSCHEKI NOVITZKY, IN THE CEDAR PLANTATIONS OF SOUTHERN FRANCE

M.A. AUGER-ROZENBERG 1 , L. ROQUES 2 , S. BOIVIN 1 , A. CHALON 3 , J.P. FABRE 3 , AND <u>A. ROQUES $^1^*$ </u>

¹Institut National de la Recherche Agronomique (INRA), Station de Zoologie Forestière, 45166 Olivet Cedex, France

Worldwide exchange and trade of tree reproductive materials is rapidly increasing with the development of plantations and ornaments using exotic tree species. Such exchanges are highly favorable to the invasion of new pests because no or few phytosanitary regulations exist for importation of such material in a number of countries, and seed-borne organisms are not easily surveyed because of their cryptic way of life. An exotic seed chalcid, Megastigmus schimitscheki (Hymenoptera: Torymidae), specialized in the exploitation of seeds of cedars (Cedrus spp.), has thus been introduced from Turkey to southern France during the early 1990s. From 1994 to 2002, seeds were screened for insect damage in the main cedar plantations of Southern France, as well as in the stands where cedar is mixed with firs (Abies spp.). X-rays were used to identify chalcid-infested seeds from which the insects were then reared. The surveys revealed that M. schimitscheki progressively invaded all the stands of Atlas cedar, C. atlantica, planted in south-eastern France. However, it has not yet reached the plantations in south-western France, where the seeds are colonised by a related exotic insect, M. pinsapinis, originating from North Africa. In the invasion zone, M. schimitscheki seems to be competitively superior to M. pinsapinis possibly because of an earlier emergence of adults. From the survey data, a mathematical model was tentatively developed to forecast the annual expansion of M. schimitscheki. In parallel, we intended to compare the genetic structure of the invasive populations with these of the native, Turkish populations using microsatellite markers. A native chalcid species, M. suspectus, was also shown to have shifted to a slight extent from a native fir, Abies alba, onto cedar. The presence of three chalcid species competing for cedar seed resources may result in a substantial decline of the regeneration potential of that tree species. At Mt Ventoux, up to 92.6% of the cedar seeds were attacked, with 86.8% due to M. schimitscheki.

²University Paris 6, Laboratory J.L.Lions, 175 rue du Chevaleret, 75013 Paris, France ³Institut National de la Recherche Agronomique (INRA), Unité de Recherches Forestière Méditerranéennes, Av. Vivaldi, 84000 Avignon, France