



**HAL**  
open science

## Spatial variation in soil CO<sub>2</sub> efflux in french and belgian temperate forests

Bernard B. Longdoz, Jérôme J. Ngao, D. Perrin, G. Vincent, Marc Aubinet, André A. Granier, Daniel Epron

### ► To cite this version:

Bernard B. Longdoz, Jérôme J. Ngao, D. Perrin, G. Vincent, Marc Aubinet, et al.. Spatial variation in soil CO<sub>2</sub> efflux in french and belgian temperate forests. Conference on Processes underlying soil carbon fluxes, Oct 2003, Kiel, Germany. 13 p. hal-02833208

**HAL Id: hal-02833208**

**<https://hal.inrae.fr/hal-02833208v1>**

Submitted on 7 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# SPATIAL VARIATION IN SOIL CO<sub>2</sub> EFFLUX IN FRENCH AND BELGIAN TEMPERATE FORESTS

LONGDOZ B.<sup>1</sup>, NGAO J.<sup>1</sup>, PERRIN D.<sup>2</sup>, VINCENT G.<sup>3</sup>,  
AUBINET M.<sup>2</sup>, GRANIER A.<sup>1</sup>, EPRON D.<sup>4</sup>

<sup>1</sup> National Institute of the Agronomic Research, Nancy, France

<sup>2</sup> University of Agricultural Sciences, Gembloux, Belgium

<sup>3</sup> University of Franche-Comté, Besançon, France


<sup>4</sup> University H. Poincaré, Nancy, France

## MAIN OBJECTIVE

Quantification of the relative importance of climatic factors and ecosystem properties on the spatial variability of soil CO<sub>2</sub> efflux (SR) from different ecosystems (plots).

1. System comparison and set up of a common protocol.
2. Quantification of climatic influences on SR for each plots.
3. Identification of the main ecosystem properties influencing SR<sub>plot</sub> variability

## SITES PRESENTATION



- Vielsalm, University of agricultural science, Gembloux, Belgium
- Hesse, National Institute of the Agronomic Research, Nancy, France
- Chaux, University of Franche-Comté, Besançon, France

Site	Position	Precipitation	Air temperature	Elevation
Chaux	47°07'N 05°42'E	950 mm	10,3°C	250 m
Hesse	48°40'N 7°05'E	1150 mm	10.5°C	300 m
Vielsalm	50°18'N 6°00'E	800 mm	8,5°C	450 m

Sites plots	Hesse			Chaux			Vielsalm	
	H1	H2	H3	C1	C2	C3	V1	V2
Tree species	Beech	Beech/Spruce		Mixed broadleaves			Beech	Douglas fir
LAI	6.3	5.5 - 8	4 - 8	7.2	7.9	7.4	3.3	5.0
Age (years)	35	35	35	Stratified forest			95	68
Ground area (m <sup>2</sup> .ha)	20.7	19.7	22.7				129	112
Soil class	Luvisol/Stagnic luvisol			Gleyic Luvisol			Dystric Cambisol	
Texture	7%(S <sub>a</sub> ); 64%(S <sub>i</sub> ); 29%(C)			20%(S <sub>a</sub> ); 50%(S <sub>i</sub> ); 30%(C)			27%(S <sub>a</sub> ); 55%(S <sub>i</sub> ); 18%(C)	
Soil density	1.078	1.103	1.108	1.024	0.978	0.997	1.038	1.044
Litter height (cm)				0.78	1.09	1.12	1.07	1.75
[C] g.kg <sup>-1</sup> (0-8 cm)	29.61	27.44	29.67				43.7	57.3
[N] g.kg <sup>-1</sup> (0-8 cm)	1.84	1.41	1.90				2.9	4.8
pH H <sub>2</sub> O	4.43	4.49	4.53	4.5	4.5	5	4.46	4.26
pH KCl	3.68	3.61	3.59				3.79	3.46

# MATERIAL AND METHODOLOGY


## 1. Four SR measurement systems

## 2. Soil Temperature ( $T_s$ )

- Thermocouple
- 10 cm depth
- Fixed (climatic station) or mobile (inserted near the measurement point)

## 3. Soil Water Content (SWC)

- Thetaprobe
- 0-20 cm depth
- Fixed or mobile



## COMPARISON

Two campaigns of measurements with the four systems :


- ✓ 05/03 Vielsam (12 collars)
- ✓ 09/03 Hesse (20 collars)

One campaign in each plots with the reference and the « local » system



1. Protocol
2. Fluxes
3. SWC
4. Ts

# 1. Protocol






➤ Direct insertion of chambers generates higher SR (confirmation)

- $P_{ch} - P_{air} \leq 0.1 \text{ Pa}$  (Fang & Moncrieff, 98; Longdoz & al. 2000)
- 60 s are sufficient between two measurements on the same point
- No impact of the duration of measurement (up to 75 s)



## 2. Fluxes




« Systems » equations :

**a**  $SR_{\text{Licor INRA}} = 0.9733 \cdot SR_{\text{Licor Ref}} + 0.0291$

**b**  $SR_{\text{PP System}} = 1.1472 \cdot SR_{\text{Licor Ref}} + 0.1392$

**c**  $SR_{\text{Licor Gx}} = 1.044 \cdot SR_{\text{Licor Ref}} + 0.3496$

### 3. SWC



### 4. Ts


- This year : measurements with fixed probes
- Future : new probes (0 - 5 cm, 10 cm)



# CLIMATIC IMPACTS

For each plots :

1. measurements on minimum 12 collars, at several dates
2. conversion to  $SR_{ref}$  using the "systems" equations
3. one average value of  $SR_{ref}$  ,  $T_s$  ,  $SWC$  , by measurement dates




## Model


Multiplicative model : separation of SWC and Ts impacts

$$SR = SR_{10} * Q_{10} \left( \frac{T-10}{10} \right) * fct(RSWC)$$

where  
 $RSWC = SWC / SWC_{sat}$



# INTERPLOTS VARIABILITY



- Ts range to small during measurement periods for Hesse and Vielsalm
- Tendency : small  $SR_{10}$  and  $Q_{10}$  for young and coniferous plots (to be confirmed)

## CONCLUSIONS AND PERSPECTIVES

- Relatively good accuracy between systems → inter-plots variability consistent
- Work still in progress (more data to improve climatic impact)
- Small  $SR_{10}$  and  $Q_{10}$  for young and coniferous plots ?
- More plots with contrasted environmental conditions and properties

OTHERS TEAMS ARE WELCOME