



**HAL**  
open science

## Molecular phylogeny of *Megastigmus* seed chalcids (Hymenoptera: Torymidae) related to conifers

Marie-Anne Auger-Rozenberg, Carole Kerdelhue, Emmanuelle Magnoux,  
Alain Roques

► **To cite this version:**

Marie-Anne Auger-Rozenberg, Carole Kerdelhue, Emmanuelle Magnoux, Alain Roques. Molecular phylogeny of *Megastigmus* seed chalcids (Hymenoptera: Torymidae) related to conifers. 22. International Congress of Entomology, Aug 2004, Brisbane, Australia. 1 p. hal-02833794

**HAL Id: hal-02833794**

**<https://hal.inrae.fr/hal-02833794>**

Submitted on 7 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## **Molecular phylogeny of *Megastigmus* seed chalcids (Hymenoptera: Torymidae) related to conifers**

MA Auger-Rozenberg, C Kerdelhué, E Magnoux and [A Roques](#)

INRA, Unité de Zoologie Forestière, BP 20619 Ardon 45166 Olivet cedex, France,

[Alain.Roques@orleans.inra.fr](mailto:Alain.Roques@orleans.inra.fr)

Most chalcid species in the genus *Megastigmus* (Hymenoptera: Torymidae) develop entirely within tree seeds, especially of conifers on which 48 species are currently recognized. Phylogenetic relationships within this group were inferred from the molecular analysis of 22 species related to host families Pinaceae (15 spp.) and Cupressaceae (7 spp.). We used both mitochondrial (cytochrome *b*) and ribosomal sequences (D2 variable region of the 28S subunit) individually and in combined analyses. Although the cyt. *b* gene seemed to be more suited than the 28S to resolve inter-specific relationships, maximum likelihood and maximum parsimony analyses produced congruent results. Whatever the method of analysis or the dataset analysed, the species grouped per conifer family, and then per conifer genus for most of them. This grouping pattern probably proceeds from the strong specific association existing with host seeds. A phylogeographic radiation at host-species level was also suggested for the species groups related to Pinaceae. In addition, the analyses clearly confirmed that a number of species are invasive in regions other than the native ones.