

Regulation of cyclin D1 expression in ES cells: the role of the Ras/MAPK pathway

Ludmila Jirmanova, Marielle Afanassieff, Ludovic Vallier, Suzy S. Markossian, Jacques Samarut, Pierre Savatier

► To cite this version:

Ludmila Jirmanova, Marielle Afanassieff, Ludovic Vallier, Suzy S. Markossian, Jacques Samarut, et al.. Regulation of cyclin D1 expression in ES cells: the role of the Ras/MAPK pathway. Cold Spring Harbor Meetings: Mouse Molecular Genetics, Aug 2000, New York, United States. 2000. hal-02840306

HAL Id: hal-02840306 https://hal.inrae.fr/hal-02840306

Submitted on 7 Jun2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Mouse Molecular Genetics

August 31 – September 03, 2000 Abstract deadline : May 22, 2000

Organizers :

Andras Nagy, Samuel Lunenfeld Research Institute, Toronto, Canada
Patrick Tam, Children's Medical Research Institute, Sydney, Australia
Siew-Lan Ang, National Institute for Medical research, London, UK
Achim Gossler, Hannover Medical School, Hannover Germany



Cold Spring Harbor Meetings : Mouse Molecular Genetics New-York, 2000 Poster Abstract :

Regulation of cylin D1 expression in ES cells : the role of the Ras/MAPK pathway.

<u>Jirmanova Ludmila</u>, Afanassieff Marielle, Vallier Ludovic, Markosian Suzy, Samarut Jacques, and Savatier Pierre. Ecole Normale Supérieure de Lyon CNRS UMR5665, INRA LA913, Lyon, France

The cellular machinery that is organized to collect extracellular signals and transduce them via tyrosine kinase receptors and the SOS-RAS-MEKK-MAPK pathay seems to be dedicated largely to driving RB phosphorylation. Cyclin D1 is likely to play a key role in this process as : (i) cyclin D1 expression is regulated by MAPK-dependent transcription factors (jun, fos, ets) and (ii) cyclin D1/CDK4 complexes are involved in RB phosphorylation during G1 phase of the cell cycle. The circuitry appears to be operative in virtually all cell types and its revelance is illustrated by its disruption in many types of human tumors. Using ES cells which express the *neo^r* gene driven off the *oct4* promoter (IOUD2 cells) so as to kill all spontaneously differentiating cells, we show that ES cells express low, although constant, evel of cyclin D1. This results from low activity of the cyl1 promoter. Promoter deletion analysis shows that this background activity originates from a proximal region of the promoter, and taht distal regions known to be involved in cyclin D1 expression in response to serum stimulation are not required. Treatment of IOUD2 cells with wortmannin (an inhibitor of RAS activation) or PD98059 an inhibitor of MEK) does not inhibit cyclin D1 expression. None of these inhibitors induce growth retardation. Induction of differntiation with retinoic acid (RA) up-regulates cyclin D1 expression, and distal - serum-responsive elements of the cyl1 promoter are involved. Differentiating cells become sensitive to inhibitors of the Ras \rightarrow ERK cascade. Both wortmannin and PD98059 inhibit cell growth and down-regulate expression of cyclin D1. Hence, cyclin D1 expression seems not to be regulated by the RAS \rightarrow MERK \rightarrow ERK phosphorylation in ES cells. This regulation is likely to be restored upon differntiation.