



Some views on possible approaches to test and improve vegetation net primary productivity models

Gérard Dedieu, Laurent Kergoat, Philippe Maisongrande, Anne Ruimy,
Bernard Saugier, Pierre-Louis Frison, Sophie Moulin, Éric Mougin, N.
Mognard-Campbell

► To cite this version:

Gérard Dedieu, Laurent Kergoat, Philippe Maisongrande, Anne Ruimy, Bernard Saugier, et al.. Some views on possible approaches to test and improve vegetation net primary productivity models. 21th assembly, European Geophysical Society (EGS). DEU., May 1996, La Hague, Netherlands. 1 p. hal-02841143

HAL Id: hal-02841143

<https://hal.inrae.fr/hal-02841143>

Submitted on 7 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

EGS May 1996

G. Dedieu, L. Kergoat, P. Maisongrande, A. Ruimy*, B. Saugier*, P.L. Frison, S. Moulin, E. Mougou, and N. Mognard-Campbell

Centre d'Etudes Spatiales de la Biosphère (CESBIO)
18 avenue Edouard Belin, 31055 Toulouse Cedex, France

*Laboratoire d'Ecologie Végétale

U.R.A. 1492, Université de Paris Sud, Centre d'Orsay, Bat362, 91405 Orsay Cedex, France

Global NPP model intercomparisons performed under the auspices of IGBP have shown that large discrepancies exist between model results such as NPP and above all intermediate prognostic variables (e.g. light use efficiency, autotrophic respiration). The origin of these discrepancies probably arise from different treatments of processes, owing to the large gaps that exist in our understanding of biological phenomena. Our capability to estimate biospheric carbon fluxes and budget under current and mainly future environmental conditions is therefore questionable. In this study, we used both parametric, satellite driven, and mechanistic, global, uncalibrated, NPP models. We will first present and discuss the comparison of model results with in-situ measurements, satellite data and CO₂ atmospheric concentration. Interannual variabilities have been analysed. We will then present the potential of multispectral satellite data assimilation to derive model parameters. Finally, a possible strategy to make models more reliable is proposed. It relies on multiscale observations of interannual variabilities.