

Hypoxylon mammatum toxins: possible involvement in canker development on aspen

Isabelle Genetet, Jean Pinon, B. Bodo, S. Rebuffat

► To cite this version:

Isabelle Genetet, Jean Pinon, B. Bodo, S. Rebuffat. Hypoxylon mammatum toxins: possible involvement in canker development on aspen. International symposium Forest Tree Physiology, Sep 1988, Nancy, France. hal-02855223

HAL Id: hal-02855223 https://hal.inrae.fr/hal-02855223

Submitted on 8 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Hypoxylon mammatum toxins: possible involvement in canker development on aspen

I. Genetet¹, J. Pinon¹, B. Bodo² and S. Rebuffat²

¹ Laboratoire de Pathologie Forestière, INRACRF, Champenoux F-54280, Seichamps, and

² Muséum National d'Histoire Naturelle, Chimie Appliquée aux Corps Organisés, CNRS UA401, 63, rue Bulfon, F-75231 Paris Cedex 05, France

Introduction

Hypoxylon mammatum (Wahl.) Miller causes a stem canker on aspen (Populus tremula) and on some poplars of the Tacahamaca section (Pinon, 1976). The disease is characterized by a flattened sunken surface with a yellow orange margin. H. mammatum prevents host callusing. Hubbes (1964) found that diffusible substances from H. mammatum agar cultures inhibited callus formation in wounds on aspen bark. Schipper (1978) and Stermer et al., (1984) confirmed the possibility that H. mammatum could produce toxins. These compounds can be isolated by partitioning into various organic solvents and chromatography. Culture age was also supposed to affect the kind and amount of metabolites produced (Stermer et al., 1984). We have isolated and characterized 2 groups of toxins from culture filtrate (Bodo et al., 1987). Optimum secretion was achieved within 6 wk of still culture on wort medium. A technique has been developed to isolate the toxins from the filtrate by adsorption onto a neutral resin (Amberlite XAD 4). Two groups of toxins were separated by elution of the XAD resin with a methanol/water gradient: hymatoxins first eluted (about 40 mg/l of filtrate) and then neutral metabolites (a few mg/l). The chemical structures of these substances were determined by spectrometric methods (MS, IR, 1D and 2D NMR). Hymatoxins are unusual diterpene sulfates with a molecular mass of about 400. The other toxin group is constituted of trihydroxytetralones.

To provide a clearer understanding of the pathotoxic features of *H. mammatum*, search for and characterization of *in vivo* toxic metabolites were undertaken and the data are reported herein.

Materials and Methods

Purification procedure

Young aspen trees were obtained from breeding programs (Lemoine, 1973). The trees were planted and inoculated with mycelium in 1981 (Pinon *et al.*, 1988). After 6 yr, the *H. mammatum* trees were cut. Wood and bark samples from both healthy and infected areas were collected. Each sample was freeze-dried, ground and then kept at -20° C. 100 g of each sample were extracted in a soxhlet apparatus with methanol for 6 h. The extracts were concentrated and an aliquot of each sample was tested for its toxicity using a leaf bioassay. The active extracts were partially purified by chromatography on Sephadex LH-20. The fractions obtained were reduced to dryness under vacuum.

Leaf bioassay

Throughout the purification procedure, samples were bioassayed by determining their effects on the leaves according to Pinon (1984). The leaves were taken from clones obtained from *in vitro* cultures and grown in a greenhouse. After cutting, the leaf petiole was placed in small tubes containing the fraction to test. 5 or

10 leaves were taken for each assay. Acetone (1%) was used as control.

Results and Discussion

When tested at a concentration of 1 mg/ml, extracts from healthy wood and bark samples had no visible effects, whereas those from infected wood and bark samples induced necrosis. However, the response was not as large with the wood extracts as compared to those of bark.



Fig. 1. In vitro and in vivo sensitivity of some clones to toxins.

Each toxic extract was separated into 9 fractions on Sephadex LH-20. They were concentrated to dryness and 100 mg of the concentrates were dissolved in 1 ml of acetone, then completed to 100 ml with distilled water. The fractions were bioassayed against 4 aspen clones which had been selected for their different behaviors under artificial inoculation: clones 717. 1.2, 712.1 and 706.8 were susceptible and clone 710.23 was quite tolerant.

The 4 clones showed the same sensitivity to hymatoxins and to the toxic extracts from infected trees: no necrosis was observed with clones 717.1.2 and 712.1, whereas leaf necrosis was observed with clones 706.8 and 710.23 (Fig. 1).

Fractions 5 and 6 of the infected bark extract were always necrotic for these last two clones, but, in addition, fraction 1 was toxic for clone 706.8 and fraction 4 for clone 710.23. From all these results, it appears that 2 groups of toxins are present in the bark extract: one is eluted in the first fractions and the second in the middle fractions of Sephadex LH-20 chromatography. When analyzed by thin–layer chromatography, both toxic fractions were still too complex to enable identification of hymatoxins.

The most toxic fractions from the infected wood extracts were fractions 6 and 7. Each of these toxic fractions was subjected to silica gel chromatography in order to obtain simpler fractions, allowing the search for the *in vivo* toxins (hymatoxins, tetralones).

Most of the studies on toxins involved in *H. mammatum* canker were carried out with cultures of this fungus on synthetic medium. The precise role of the metabolites thus isolated in pathogenesis must be investigated. Our preliminary results

demonstrated the presence of toxic substances in infected bark and wood from a freshly cut tree. These compounds have no effect on the clone known for its resistance to *in vitro* toxins.

Preliminary chemical examination of the complex toxic extract from infected bark does not enable us to conclude as to the presence or absence of hymatoxins and trihydroxytetralones.

References

Bodo B., Davoust D., Lecommandeur D., Rebuffat S., Genetet I. & Pinon J. (1987) Hymatoxin A, a diterpene sulfate phytotoxin of *Hypoxylon mammatum*, parasite of aspen. *Tetrahedron Lett.* 28, 2355-2358

Hubbes M. (1964) New facts on host-parasite relationships in the *Hypoxylon* canker of aspen. *Can. J. Bot.* 42, 1489-1494

Lemoine M. (1973) Amélioration des peupliers de la section *Leuce* sur sols hydromorphes. Thèse, Université de Nancy I, France

Pinon J. (1976) Une menace grave pour les trembles alpins: le chancre à *Hypoxylon mam-matum. Rev. For. Fr.* 28, 31-34

Pinon J. (1984) Propriétés biologiques de la toxine d'*Hypoxylon mammatum*, parasite des peupliers de la section *Leuce. Rev. Cytol. Biol. Végét. Bot.* 7, 271-277

Pinon J., Genetet I., Bodo B. & Rebuffat S. (1988) Apport des vitrocultures à l'étude des métabolites secondaires d'origine fongique agissant sur le cambium des peupliers. 32^e Colloque de la Société française de Phytopathologie, Angers, 14–15 mai, ACTA, in press

Schipper A.L. Jr. (1978) A *Hypoxylon mammatum* toxin responsible for canker formation in quaking aspen. *Phytopathology* 68, 868-872

Stermer B.A., Scheffer R.P. & Hart J.H. (1984) Isolation of toxins of *Hypoxylon mammatum* and demonstration of some toxic effects on selected clones of *Populus tremuloides*. *Phytopathology* 74, 654-658