



# Hydrolysis of native wheat starch by an alpha-amylase from *Bacillus subtilis*

Paul Colonna, Alain Buleon, Christiane Mercier

## ► To cite this version:

Paul Colonna, Alain Buleon, Christiane Mercier. Hydrolysis of native wheat starch by an alpha-amylase from *Bacillus subtilis*. 72. annual meeting of the American Association of Cereal Chemists, Oct 1987, Nashville, TN, United States. hal-02855923

**HAL Id: hal-02855923**

**<https://hal.inrae.fr/hal-02855923>**

Submitted on 8 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

AACC 72nd Annual Meeting  
October 31-November 5, 1987, Opryland Hotel, Nashville, TN

---

Inventaire Pub  
N° 392

**73** Hydrolysis of native wheat starch by an alpha-amylase from *Bacillus subtilis*. P.Colonna , A, Buleon and Christiane Mercier , Institut National de la Recherche Agronomique, Rue de la Géraudière, 44072 Nantes Cédex , France.

Suspensions (2%) of native starch granules from wheat have been degraded by an alpha-amylase from *Bacillus subtilis*. Crystalline fractions have been prepared either by crystallization from short chain amylose or mild acid hydrolysis of native wheat starch. Complete (98%) depolymerization into small soluble oligosaccharides is obtained within 24h at high enzyme concentrations (above 189 microg. protein/mL). Electron microscopy, particle size analysis, differential scanning calorimetry and X-ray diffractometry reveal that enzymic degradation occurs granule-by-granule: the enzyme does not preferentially hydrolyze the more amorphous regions of the granules. Intrinsic viscosity, gel permeation chromatography on Sepharose CL-2B show off that the remaining material has a macromolecular structure close to the native starch: there is no preferential hydrolysis of either amylose or amylopectin. In contrast, acid erodes simultaneously all the granules, leading to a splitting into small crystalline particles. *De novo* crystalline fractions are completely degraded by alpha-amylase. These results support evidence for an active disentanglement of chains, carried out by the different subsites of alpha-amylase molecules. A simple mathematical treatment is proposed in order to explain the results of the kinetics.